

**Yu. I. Logachev, G. A. Bazilevskaya, E. V. Vashenyuk, E. I. Daibog, V. N. Ishkov,
L. L. Lazutin, L. I. Miroshnichenko, M. N. Nazarova, I. E. Petrenko, A.G Stupishin,
G. M. Surova, O. S. Yakovchuk**

CATALOG

**of Solar Proton Events in the 23rd Cycle of
Solar Activity (1996 – 2008)**

Moscow – 2016

Skobel'tsyn Institute of Nuclear Physics, Moscow State University (SINP), Moscow, Russia
Lebedev Physical Institute, Russian Academy of Sciences (LPI), Moscow, Russia
Polar Geophysical Institute, Kola Science Center, Russian Academy of Sciences (PGI), Apatity, Russia
Pushkov Institute of Terrestrial Magnetism, Ionosphere and Radio Wave Propagation, Russian Academy of Sciences (IZMIRAN), Moscow, Troitsk, Russia
Geophysical Center, Russian Academy of Sciences (GC RAS), Moscow, Russia
Saint Petersburg State University (SPbU), St. Petersburg, Russia
Fedorov Institute of Applied Geophysics (IAG), Moscow, Russia

List of Authors:

Yu. I. Logachev (SINP), **G. A. Bazilevskaya** (LPI), **E. V. Vashenyuk** (PGI), **E. I. Daibog** (SINP),
V. N. Ishkov (IZMIRAN, GC RAS), **L. L. Lazutin** (SINP), **L. I. Miroshnichenko** (IZMIRAN),
M. N. Nazarova † (IAG), **I. E. Petrenko** (IAG), **A. G. Stupishin** (SPbU), **G. M. Surova** (SINP),
O. S. Yakovchuk (SINP)

Edited by **Yu. I. Logachev**

Catalog

of Solar Proton Events in the 23rd Cycle of Solar Activity (1996 – 2008)

Moscow – 2016

Contents

	Page
Editor's preface	4
General Description of the Catalog	5
Introduction	5
Description of Layout for a separate solar proton event	6
Description of the particle flux Tables	8
Description of the concomitant solar phenomena	9
Conclusion	10
Reference list to the Catalog Description	10
Supplement 1. Spacecraft and instruments with data for Catalog	11
Supplement 2. Determination of source increase of proton fluxes in the solar energetic particle events	15
Supplement 3. Determination of the relativistic solar proton fluxes and energy spectrum from the neutron monitor data	17
Supplement 4. Determination of the quasi maximal energy of protons in an event . .	19
Supplement 5. List of symbols and abbreviations on the phenomena and events used in the Catalog materials	20
Reference list for all events of 1997–2006	23
Events of 1997–2006	
Events in 1997	34
Events in 1998	49
Events in 1999	120
Events in 2000	169
Events in 2001	270
Events in 2002	412
Events in 2003	525
Events in 2004	583
Events in 2005	632
Events in 2006	714
Summary Table of Solar Proton Events in the 23rd Cycle of Solar Activity (1997–2006)	736

Editor's preface

This Catalog of solar proton events (SPE) in the 23rd cycle of solar activity (from June 1996 till December 2008) is a continuation of the previous issues [1–5] which described SPE from 1970*) [<http://www.wdcb.ru/stp/data/SPE/>]. In the Catalog the data on SPE with the maximum flux of protons with energy $E_p > 10$ MeV, $J_p \geq 1 \text{ cm}^{-2} \cdot \text{s}^{-1} \cdot \text{sr}^{-1}$ (pfu), are collected and systematized. There were 142 such events in the solar cycle 23. The last event, presented in the Catalog refers to December 2006, since there were no events with such proton fluxes during the last two years of the solar cycle 23.

In addition to the data on the solar proton fluxes as observed by several spacecraft the Catalog gives information on the particle sources and concomitant X-ray, optic, and radio emission. Integral energy spectrum in the maximum of the intensity-time profile (so called time of maximum spectrum) is presented for each SPE. Some events with a complicated intensity-time profile have 2 or even 3 maxima. In this case the spectra for each maximum are given.

The derived energy spectra are used for finding an energy value where the solar proton flux is equal to 0.1 of the galactic cosmic ray flux. We consider this value as a certain proxy for a maximum energy of solar protons in a given event and call it the Quasimaximal Energy. Such definition provides a kind of standard approach to value of maximum energy in a given SPE.

Unlike previous issues of the Catalog each event is now illustrated by the plots of temporal behavior of X-ray, electron and protons fluxes, solar wind speed, strength of interplanetary magnetic field, and the Dst- index during the time interval around this event. Each event begins with a short legend followed by the detailed information.

The scientists from different institutions who provided a wide coverage of the SPE description on the base of solar and space physics, geophysics, and cosmic ray physics concerning problems of charged particle acceleration on and near the Sun and particle propagation in space and the Earth's magnetosphere are among the authors of the Catalog.

Experience with the previous issues proved the Catalogs to be very useful in statistical investigations of SPE, search for regularities in the energy spectra, mechanisms of solar particle acceleration and propagation. Information on the energy spectrum of solar protons is necessary for estimation and forecast of radiation level in the inner heliosphere. The Catalog will be useful for research of the particle penetration into the Earth's magnetosphere, polar cap absorption, radio wave propagation and other problems of solar-terrestrial coupling.

This work was supported by the Russian Foundation for Basic Research, grant № 13-02-00612.

The authors of the Catalog thank professor M.I. Panasyuk, the Director of SINP, professor V.D. Kuznetsov, the Director of IZMIRAN, Professor V.B. Lapshin, the Director of IAG, and doctor V.S. Makhmutov, Head of Laboratory of Solar Physics and Cosmic Rays of LPI, for their support and interest. Thanks are due to many colleagues for helpful discussions and constructive criticism. The authors are grateful to N.A. Sergeyeva and L.P. Zabarinskaya from the World Data Center for Solar-Terrestrial Physics, GC RAS, for technical editing of the Catalog.

*) References are given after General Description of the Catalog

General Description of the Catalog

Introduction

The data on the charged particle fluxes of solar origin in the solar cycle 23 (April 1996 – December 2008), registered at 1 AU by different spacecraft, balloons and by ground-based neutron monitors are presented in the Catalog. In 1996 and during the minimum of solar activity in 2007 – 2009 there were no events, which correspond to the previously mentioned conditions relevant for including in the Catalog, therefore only the data from 1997 to 2006 are presented in the Catalog.

A solar proton event is defined as a flux enhancement with the maximum in the intensity-time (time of maximum) profile for >10 MeV protons $J_p(>10 \text{ MeV}) > 1 \text{ pfu}$, where $1 \text{ pfu} = 1 \text{ cm}^{-2} \cdot \text{s}^{-1} \cdot \text{sr}^{-1}$. As a rule the TOM flux estimation is based on the hourly averaged data of the ESP instrument of the geostationary GOES satellites. In the case of the GOES data are absent or the Earth's magnetosphere is strongly disturbed the data of other spacecraft instruments compiled in the Catalog are used. Along with the isolated enhancements of solar proton fluxes the Catalog incorporates the SPE with complicated time-intensity profiles with several intensity maxima belonging to the same event and originating usually from the same active region. For such events the energy spectra were derived for the most prominent maxima, usually for 2 and rarely for 3 or more. The identical information for each selected maximum is given in the Tables as described below.

The spacecraft and instrumentation are briefly described in **Supplement 1**.

One of the most important problems in the study of SPE is identification of its source – the explosive energy release which generated the energetic protons. Here, we consider the dynamics of electromagnetic radiation in all observable ranges accompanying the explosive energy release in the limited regions of the solar atmosphere. Depending on the local magnetic field strength it can be solar flares or coronal mass ejections (CME). If SPE occurs on the back side of the Sun the CME and radio emission are still observed. Identification of the SPE sources is a long-standing problem, therefore the methods and criteria used for this purpose were described in all previous issues of the Catalog [1–5]. Nevertheless keeping in mind importance of the point, a slightly modified procedure is given in **Supplement 2 “Identification of the solar proton sources”**.

The main source of information on the relativistic solar cosmic rays are the ground-based installations – neutron monitors which record the SPE of very high energy where the particles reaching the ground level were generated. Such events are called the Ground Level Enhancements (GLE). There were 16 such events (#55 – 70) during the 23rd cycle of solar activity.

Conversion from the neutron monitor records to the proton fluxes outside the magnetosphere is a complicated process which accounts for several mechanisms of particle propagation in the Earth's magnetosphere and atmosphere. It requires solution of an inverse problem, i.e. retrieving of the energy spectrum of solar protons in space from the data of neutron monitors. More sophisticated approaches to modeling of particle transport are under development. This problem is considered in more detail in the **Supplement 3 “Retrieving of the relativistic solar proton fluxes and energy spectrum from the neutron monitor data”**.

Note that since 2006 the relativistic particle measurement is accomplished onboard the orbital spectrometer PAMELA which observes in the near-Earth space fluxes of protons and nuclei with energies from 80 MeV to several GeV. The Catalog presents the PAMELA data on the GLE–70 in 2006.

With the aim to estimate the maximal energy of accelerated protons a new parameter is introduced in the Catalog, i.e. **the quasi maximal energy of protons in a given event, E_{qm}** , which describes the spectrum in the range of highest energies recorded. A new parameter is a first approach to real maximal energy of protons observed in a given event. **The quasi maximal energy is defined as the energy of parity between the flux of solar protons and of the 0.1 fraction of**

the galactic cosmic ray (GCR) protons. The method of evaluation of the quasi maximal energy of solar protons is described in the **Supplement 4**.

Description of Layout for a separate solar proton event

The description of each event begins from the event title, which contains the date of its start in a following way:

YYYY.MM.DD – (YYYY-DOY). For example, the first event of the Catalog, which started at November 4, 1997, is named as **1997.11.04 – (1997-308)**. Indication of DOY is useful for comparison with other reference data. The number in the right end of this line corresponds to the event numeration started from the first event in 1970 according to the Catalog of solar proton events 1970 – 1979 [1].

After that the LEGEND is given, which includes some general information about the event. An example of the legend and the explanation of its every line are given below.

Event 2004.07.25 – (2004-207)

№ 450

Particle event: To(Ep>10MeV) – 25d17^h

Tmax₁(Ep>10MeV) – 25d21^h, Jmax₁(Ep>10MeV) – 27 /cm².s.sr

Tmax₂(Ep>10MeV) – 26d23^h, Jmax₂(Ep>10MeV) – 430 /cm².s.sr

Duration of the event – 4 days

Quasimaximal energy of protons in the event – Eqm(max₁) = 140 MeV

– Eqm(max₂) = 155 MeV

Sources: ● solar flare 25d14^h19^m, M1.1/1F, N08W33, AR10652

Ø solar flare 26d17^h23^m, M1.1/2N, N03W45, AR10652

Main X-ray burst 1-8 Å: onset – 25d14^h19^m, max – 25d15^h14^m, Φ = 0.065 J/m²

● CME: 25d14^h54^m; V = 1333 km/s; Δφ = 360°; dA = 204°

▲ SC 26d22^h49^m;

The following information is given:

Event 2004.07.25 – (2004-207)

№ 450

Date of the event as YYYY.MM.DD – (YYYY-DOY) and its number

The first line:

Particle event: To(Ep>10MeV) – 25d17^h – the day and hour of the first arrival of >10 MeV proton as observed by the ESP instrument onboard the geostationary GOES spacecraft or defined from other spacecraft presented in the Catalog. Accuracy of To is within 1 hour.

More accurate determination of the event start even using data with higher time resolution, is difficult and may be erroneous because of variability in the background count rates of instruments.

The second (a) and (b) lines:

(a) Tmax₁(Ep>10MeV) – 25d21^h, Jmax₁(Ep>10MeV) – 27 /cm².s.sr

(b) Tmax₂(Ep>10MeV) – 26d23^h, Jmax₂(Ep>10MeV) – 430 /cm².s.sr

A day and time of >10 MeV proton flux maximum (two maxima in this event) determined from all available data presented in the Catalog for a given event and the values of these fluxes usually as observed by the GOES. There were two maxima in this event, so two lines (a) and (b) are provided.

In the Catalog, all flux values are hourly averaged, and time is UT. The accuracy of Tmax is 1 hour. The accuracy of Jmax (Ep > 10 MeV) is not worse than ±10%. The Tmax is determined with rather low accuracy because of using hourly averaged values of particle fluxes during the whole event.

The third line gives the event total duration including all selected maxima in case of a complicated event. In this event:

Duration of the event – 4 days.

The last line contains the Eqm values, i.e. quasi maximal energy of protons. Eqm is determined for each maximum in this event and it is calculated from the integral energy spectrum of protons using a procedure described in the **Supplement 4**. In this event

Quasimaximal energy of protons in the event – $\text{Eqm}(\text{max}_1) = 140 \text{ MeV}$

– $\text{Eqm}(\text{max}_2) = 155 \text{ MeV}$

The Eqm values are found with accuracy $\pm 10\%$.

The second part of the LEGEND contains information on sources of particles observed near the Earth. The following is for **the Event 2004.07.25 – (2004-207)**:

Sources: ● solar flare 25d14^h19^m, M1.1/1F, N08W33, AR10652

Ø solar flare 26d17^h23^m, M1.1/2N, N03W45, AR10652

Main X-ray burst 1-8 Å: onset – 25d14^h19^m, max – 25d15^h14^m, $\Phi = 0.065 \text{ J/m}^2$

● CME: 25d14^h54^m; $V = 1333 \text{ km/s}$; $\Delta\phi = 360^\circ$; $dA = 204^\circ$

▲ SC 26d22^h49^m;

In the case the particle source was a flare, the first line gives the following flare properties: Time of maximum in the H α (25d14^h19^m) line – day, hour and minutes, the X-ray burst class and optical flare importance (M1.1/1F), coordinates (N08W33) and the number of the active region where the flare occurred (AR10652). If the optical importance goes ahead of X-ray burst class (e.g., 2N/M1.6), this means that the onset of the optical flare was recorded earlier than onset of the X-ray burst.

At the beginning of the line a symbol is placed indicative of confidence of the flare association to this event.

●,■ – the flare is a certain source of the observed protons;

⊙,□ – the flare is a probable source of the observed protons;

○ – the flare is a possible but for some reason doubtful source of the observed protons;

Ø – the flare is not the main source of the observed protons but contributed, or might have contributed to the proton flux.

The following symbols are used in the case when a flare was not identified:

□ – a flare (or flare associated activity) beyond the western or eastern limb of the Sun;

◇ – solar activity on the solar disk not associated with flares, modulation effects in the interplanetary space.

Procedure for identification of the solar proton sources is given in the **Supplement 2**.

In the Part **Sources** the following information on the soft X-ray burst in the range of 1–8 Å (12.4–1.55 keV) as measured by GOES is presented: the time of onset and maximum of the burst (in this event 25d14^h19^m и 25d15^h14^m), Φ – the integral of energy flux from the burst onset till the half of maximum level ($\Phi = 0.065 \text{ J/m}^2$).

Further the information on CME, if any, contains the time of CME first appearance in the coronagraph field of view (in this event 25d10^h54^m), the CME speed (1333 km/s), angular width near the Sun ($\Delta\phi = 360^\circ$) and the central position angle of CME as observed by the LASCO-SOHO coronagraph – ($dA = 204^\circ$).

If one or several storm sudden commencements (SC) took place their day and time are indicated on the last line after the symbols ▲ or Δ: ▲ stands for an SC that affected the particle flux, Δ stands for SC that did not change the intensity-time profile.

After the Legend, the overview plots concerning concomitant phenomena on the Sun and in the near-Earth space are presented. From top to bottom the following data are plotted: 1 min flux values of the soft X-rays as measured by GOES; hourly averaged fluxes of electrons as measured by EPHIN/SOHO and protons with energies $E_p > 10$, > 30 and > 100 MeV (GOES data), solar wind velocity, IMF strength and Dst-index.

The plots cover longer time than duration of event considered. The time interval related to given event is slightly shadowed.

Intensity-time profiles of proton and electron fluxes describe the solar situation, type of solar proton event, (i.e. the single, multiple or complex), number of maxima in the complex event, contribution of shocks into the flux of particles, Forbush decreases in the solar particles and the first insight of the proton spectrum in the range of 10 – 100 MeV. Concomitant bursts of the soft X-ray radiation give evidence of a flare as a possible source of the event basing on its power.

After the overview figures, the time profiles of the proton fluxes are presented in larger scale as measured by several instruments in various energy ranges. Upper panels give the integral fluxes recorded by spacecraft ACE and GOES, lower panel gives the differential fluxes recorded onboard SOHO. These spacecraft situated inside the magnetosphere on the geostationary orbit (GOES) and at the libration point L1 (ACE and SOHO) give the most full information about particles before, during and after start of the event.

After that the integral energy spectra at the time(s) of maximal intensity of the event is(are) depicted. The integral spectra were compiled with account of all energy channels contributed to the event. The differential particle flux, i.e. recorded in limiter energy intervals, were recalculated to the integral values. In the case of several intensity maxima in a given event the spectra were compiled for each maximum separately.

The high energy part of the spectrum was fitted by a power-law function ($J_p \sim E^{-\gamma}$), shown in the figure with the power law index γ for each spectrum. Because of significant scatter of results taken from various spacecraft this procedure gives γ with rather low accuracy, not higher than $\pm 10\%$.

Unfortunately, only scarce space observations in the range of 200 – 600 MeV were fulfilled during the 23rd cycle of solar activity. That makes difficult the spectrum determination in the event of medium power. This gap was partly filled with the balloon data, while the neutron monitor observations were utilized for powerful events.

Retrieving of the proton flux outside the Earth's magnetosphere from the neutron monitor count rate enhancement in percent of the background count rate before the event is a complicated task which is solved specially for each event. Note that with the aim of avoiding of the strong anisotropy effect the so called delayed component of solar protons determined from the neutron monitor network [10] was used. For weak GLE we took the maximum enhancement recorded.

Description of the particle flux Tables

After the spectra figures there are the particle flux Tables for a given event. The Tables contain the particle flux values at various energies in the maximum of the intensity-time profile (for each maximum) as recorded by different spacecraft. Usual conventional designations for spacecraft are used, BALLOONS stand for the observation in the stratosphere, NM means the ground-based neutron monitor measurements. The instrument description is given in **Supplement 1**.

All designations in the Tables are commonly accepted. Note that if the maximum occurred in the same day as the event start then only time without the day is indicated, if the maximum was next day or later then the day is indicated. For example, 20^h means that maximum occurred on the day of

the event start, 07d08^h – the maximum occurred next day or later. In the case of two or more maxima they are separated by the slash symbol “/”, for example, 11^h/21^h means that both maxima were on the same day, 21^h/07d08^h – the second maximum was next day or later.

The flux values in the maximum of the time profile are given without any corrections of the original data but the background is subtracted. The accuracy and of data presentation in the Tables is not worse than 10%. In the case of two or more maxima the flux values are separated by the symbol “/” in accordance with the times of maxima. The event duration is measured in days. Note that this parameter shows strong variations depending on the instrument and the proton energy. The duration is measured with accuracy not worse than ± 1 day.

Description of the concomitant solar phenomena

The following are materials on the observed solar flare phenomena, sources of proton events parameters and characteristics of the flare event are tabulated with numerical values and a graph (graphs) of its frequency spectrum is presented. Since the conditions of their registration differ for different events, tables are also differ for different flare events. Below the list of all data in all Catalog events is presented.

As in all previous Catalogs [1–5], all tables begin with the title line, which shows the date, the degree of reliability of identification (●, ■, ☉, □, ○, ◻ or Ø) of solar flare event, the source of the corresponding proton event, the active region (AR) according to NOAA US solar service and a sequence number of proton event.

All designation listed in the table of parameters are commonly accepted in the solar physics, the details can be found in the descriptive volumes of monthly bulletin «Solar Geophysical Data» (SGD) [6]. Data on the soft X-ray bursts are from GOES measurements. Main data on hard X-rays and gamma from 1996 to 2010 were taken from catalogs of solar flares [<http://www.ssl.berkeley.edu/~moka/rhessi/catalogs.html>] and [7, 8]. Information from CORONAS-F on hard X-ray bursts is from [9]. CME data were taken from the Catalog of CMEs LASCO (SOHO) [http://cdaw.gsfc.nasa.gov/CME_list/].

Information in the Tables of the Catalog is organized by blocks, including all observed diapasons of electromagnetic radiation (optics, soft X-rays, gamma and radio radiation), dynamical phenomena of H α line and of continuous spectrum (WL – in white light).

In each block of data meanings of Table columns slightly differ for different events. Each section of the Table is separated by an empty narrow line. Absence of some data or whole lines in the Tables indicates absence of corresponding phenomena or observations in considered event.

Information is grouped by columns (1 – 8) in the next way:

- Optical diapason data: (1) – spectral H α line, (2) – wave length, (3 – 5) – times of maximum and the end of the event, (6) – heliocoordinates, localization of a flare on the visible solar disk, (7) –optical importance of the flare event, (8) – code of the flare, characterizing the structure of the event according to SGD. Few next lines present dynamical phenomena in H α line (EPL, LPS, DSF, SPY, BSL) in a case of their observation during a flare event.
- Soft X-ray data: (1 and 2) – diapason of energy and units of measurement, (3 – 5) times of start, maximum and end of the event, (7) – radiation flux in flare maximum (importance/class of the burst), (8) – integral soft X-ray flux in joules per square meter J/m².
- The same information on hard X-ray and/or gamma radiation.
- Data on radio emission of fixed frequency (1) – frequency, (2) – units of measurements, (3 – 5) – times of start, maximum and end of radio event, (6) – code (see below explanations), (7) – logarithm of flux in units 10⁻²² W m⁻² s⁻¹ (more detailed see in previous Catalogs descriptions).
- Characteristics of dynamical radio burst spectrum: (1) – kind, frequency diapason in MHz, (3 and 5) – times of start and end, (6) – type, (7) – importance. Type of dynamical spectrum (DS) of

the meter component and features of radio burst are from [6, 10] and [Solar Radio Burst: <ftp://ftp.ngdc.noaa.gov/STP/space-weather/solar-data/solar-features/solar-radio/radio-bursts/reports/spectral-listings>].

- Data on solar neutrons: (1) – ⁰n indicates that solar neutrons were recorded in a given proton event; (7 – 8) – neutron monitor(s) and/or spacecraft detected the neutrons according to [7, 8].
- Data on coronal mass ejection (CME): (2) – continuous spectrum (WL), (3) – time of the first appearance in sight of coronagraph LASCO/C2 κ/a SOHO, (4) – median value of the speed V (km/s) of radial CME propagation in sight of coronagraph, (5) – acceleration, km/s², can be positive, negative or close to zero, (6) – Δφ, angular size (degrees), (7) – dA, positional angle (degrees). It's necessary to note that absence of CME data in the table doesn't mean absence of CME in the event. It is possible that during considered event there were no observations on coronagraph («gap»), or behind the limb localization of the flare did not contribute to the possibility of its registration.

After the Tables, describing solar phenomena accompanying SPE, plots of radio spectrum in the decimeter and centimeter range of wavelengths are presented. Such a spectrum allows to estimate and potentially identify patterns that are typical of flare events.

A more detailed description of all the symbols shown in the tables is in **Supplement 5**.

Information about the event is completed by the short list of references (not complete). Detailed information on publications is presented in the general publications list at the end of Catalog.

Conclusion

The Catalog contains the data on 142 solar energetic particle events occurred in the 23rd cycle of solar activity (1996 – 2008). There is digital and graphic information on the fluxes and energy spectra of solar protons in the near-Earth space as well as plots of the intensity-time profiles for all the events. Comprehensive description is presented of concomitant solar phenomena, optical flares, radio and X-ray emission, characteristics of the flaring active regions, flares etc.

The Catalog is intended for a wide range of professionals working in the field of solar-terrestrial physics, experts in solar physics and, radiation safety of space flight, and people interested in the problems of the impact of solar activity on the atmosphere and biosphere of the Earth.

Reference list to the Catalog Description

1. Akiniyan S.T., Bazilevskaya G.A., Ishkov V.N. et al. Catalog of solar proton events 1970 – 1979 // Edited by Yu.I. Logachev. Moscow, IZMIRAN, 1982. 184 p. http://www.wdcb.ru/stp/data/SPE/SPE_1970-1979.pdf
2. Bazilevskaya G.A., E.V. Vashenyuk, V.N. Ishkov et al. Catalogue of energy spectra of solar proton events 1970 – 1979 // Edited by Yu.I. Logachev. Moscow, IZMIRAN, 1986. 234 p. [http://www.wdcb.ru/stp/data/SPE/SPE_1970-1979_\(SPECTRA\).pdf](http://www.wdcb.ru/stp/data/SPE/SPE_1970-1979_(SPECTRA).pdf)
3. Bazilevskaya G.A., E.V. Vashenyuk, V.N. Ishkov et al. Solar Proton Events. Catalogue 1980 – 1986. Data on particles and electromagnetic emission // Edited by Yu.I. Logachev. Materials of the World Data Center B, Soviet Geophysical Committee of the Academy of Sciences of the USSR, Moscow, 1989. 160 p. http://www.wdcb.ru/stp/data/SPE/SPE_1980-1986.pdf
4. Bazilevskaya G.A., E.V. Vashenyuk, V.N. Ishkov et al. Solar Proton Events. Catalogue 1980 – 1986. Plots of the time profiles and energetic spectra of protons, synoptic charts and schemes of sunspot groups // Edited by Yu.I. Logachev. Materials of the World Data Center B, Soviet Geophysical Committee of the Academy of Sciences of the USSR, Moscow, 1990. 204 p. [http://www.wdcb.ru/stp/data/SPE/SPE_1980-1986_\(SPECTRA\).pdf](http://www.wdcb.ru/stp/data/SPE/SPE_1980-1986_(SPECTRA).pdf)

5. Sladkova A.I., G.A. Bazilevskaya, V.N. Ishkov et al. Catalogue of solar proton events 1987 – 1997. // Edited by Yu.I. Logachev. Moscow University Press, Moscow, 1998. 246 p. http://www.wdcb.ru/stp/data/SPE/SPE_1987-1996.pdf
6. Solar Geophysical Data. National Geophys. Data Center, Boulder, Colorado, USA, 1996–2006. ftp://ftp.ngdc.noaa.gov/STP/SOLAR_DATA/SGD_PDFversion
7. Muraki Y. Solar neutron events that have been found in solar cycle 23 // Proceedings of the Second International Symposium “Solar Extreme Events” (SEE-2005), Nor-Amberd, Armenia, 26-30 September 2005.
8. Xiao Xia Yu, Hong Lu, Guan Ting Chen et al. Detection of solar neutron events and their theoretical approach // New Astronomy. 2015. V. 39. P. 25–35.
9. Kuznetsov S.N., Kurt V.G., Myagkova I.N. et al. Gamma-ray emission and neutrons from solar flares recorded by the SONG instrument in 2001 – 2004 // Solar System Research. 2006. V. 40. No. 2. P. 104–110. DOI: [10.1134/S0038094606020031](https://doi.org/10.1134/S0038094606020031)
10. Vashenyuk E.V., Balabin Yu.V., Gvozdevsky B.B. Features of relativistic solar proton spectra derived from ground level enhancement events (GLE) modeling // Astrophys. Space Sci. Trans. 2011. V. 7. P. 459–463. doi:10.5194/astra-7-459-2011. <http://www.astrophys-space-sci-trans.net/7/459/2011/astra-7-459-2011.html>

Supplement 1

Spacecraft and instruments with data for Catalog

GOES (Geostationary Operational Environmental Satellite system) are the satellites operating since 1974 on the geostationary orbit. The Catalog incorporates information from the semi-conductor telescopes **EPS** of the **GOES-9** and **GOES-10** satellites. The telescope **EPS** registers solar protons on 6 integrated channels with energy >5, >10, >30, >50, >60 and >100 MeV. Geometric factors for all the channels are given in the Table “The space crafts and instruments” (see below). Detailed description of the **GOES** satellites and instrumentation can be found in the GOES Data Book [<http://goes.gsfc.nasa.gov/text/databook/section05.pdf>].

METEOR – is common name of meteorological satellites with polar circular sun-synchronous orbits with altitude of 1000 km. Data from three different detectors were used in Catalog: **CBM** have Geiger counters with different screens providing proton on four integral channels (>5, >15, >25 and >40 MeV), scintillator **BP** registered protons with energy >90 MeV and cherenkov detector **ChD** registered protons with energy >600 MeV. Data for year 1997 were obtained on board of **METEOR-2**, all others on **METEOR-3M**. **METEOR** description and database are given at [<http://smdc.sinp.msu.ru/index.py?nav=meteor3m>].

CORONAS-F – low altitude polar orbiter (500 km) operates from August 2001 to May 2005. General description of the **CORONAS-F** mission is given in [<http://coronas.izmiran.ru/F/>]. In Catalogue there were used data of the **MKJI** spectrometer described in: Kuznetsov S.N., Bogomolov A.V., Galkin V.I. et al. Scientific set of instruments “Solar Cosmic Rays” // The Coronas-F Space Mission. Key Results for Solar Terrestrial Physics. Ed. V. Kuznetsov. Astrophysics and Space Science Library, V. 400, Springer, Berlin Heidelberg, 2014. P. 289–299. doi:10.1007/978-3-642-39268-9-12. <http://www.springer.com/us/book/9783642392672>. Proton data registered in four differential channels 1 – 5 MeV, 14 – 26, 26 – 50 and 50 – 90 MeV

are given in a site [<http://smdc.sinp.msu.ru/index.py?nav=coronasf>]. Solar proton fluxes measured by **CORONAS-F** in the polar cap were given in Catalogue recalculated to the integral fluxes >1, >14, >26 and >50 MeV.

POES (Polar Orbiting Environmental Satellites – project of the National Oceanic and Atmospheric Administration – NOAA) is a series of satellites on polar circular sun-synchronous orbits. Altitudes of all **POES** satellites usually range from 700 to 850 kilometers, with orbital periods from 98 to 103 minutes. For the Catalogue **POES-15** and **POES-16** measurements by the **MEPED** (Medium Energy Proton and Electron Detector) were used. Energy interval of the **MEPED** is 0.024 – 140 MeV. Detailed description of **MEPED** is given at [<http://www.ngdc.noaa.gov/stp/satellite/poes/>].

IMP-8 (Interplanetary Monitoring Platform). In the catalog **CPME** (Charged Particles Measurement Experiment) – (Johns Hopkins University, Applied Physics Laboratory – JHU/APL), [http://sd-www.jhuapl.edu/IMP/imp_index.html#cpme].

ACE (Advanced Composition Explorer). ACE description and the list of experiments can be found at [<http://www.srl.caltech.edu/ACE/>]. The results of measurements of proton fluxes $E > 10$ MeV and > 30 MeV from instrument **SIS** (The Solar Isotope Spectrometer) in the Catalog were used. See data at the [http://www.srl.caltech.edu/ACE/CRIS_SIS/].

SOHO (Solar and Heliospheric Observatory). Information about **SOHO**: [http://sohowww.nascom.nasa.gov/about/docs/SOHO_Fact_Sheet.pdf]. Data of two instruments from at **SOHO** (**LION** and **EPHIN**) were used in the Catalog.

– The instrument **LION** (Low Energy ION and Electron Instrument) is capable to measure particle spectra in the range 44 keV to 6 MeV for protons. Two **LION** channels with proton energies 0.75 – 2 MeV and 2 – 6 MeV were used. **LION** data are at [<http://sohowww.nascom.nasa.gov/>] and [<http://www.ieap.uni-kiel.de/et/ag-heber/costep/>].

– **EPHIN** (The Electron Proton Helium INstrument). For the Catalog data of proton channels 4 – 8 MeV; 8 – 25; 25 – 41 and 41 – 53 MeV (25 – 53 MeV – after a year of the mission) were used, as well as INTEGRAL: > 10 MeV (electrons) + > 50 MeV (protons). **EPHIN** data are at [<http://sohowww.nascom.nasa.gov/>] and [<http://www.ieap.uni-kiel.de/et/ag-heber/costep/>].

STEREO (Solar TERrestrial RELations Observatory). Descriptions of **STEREO A** and **B** are at [<http://stereo-ssc.nascom.nasa.gov/>] and in: *The STEREO Mission*. Ed. C.T. Russell. Springer Science & Business Media, BV. 2008. DOI: 10.1007/978-0-387-09649-0_14. For the Catalog data from the **HET** (High Energy Telescope) instruments were used. Instruments **HET** registered protons in energy interval from 13.6 to 100 MeV, which were united into three energy intervals: 13.6 – 29.5 MeV, 29.5 – 60 MeV and 60 – 100 MeV. For obtaining energy spectra the authors recalculated data of these three united **HET** intervals into integral fluxes $J_p (>13.6, >29.5, >60 \text{ MeV})$ [<http://stereo-ssc.nascom.nasa.gov/>].

PAMELA (Payload for Antimatter-Matter Exploration and Light Nuclei Astrophysics): Magnetic spectrometer **PAMELA** is an international project being performed by collaboration of Italy, Russia, Germany and Sweden. The apparatus was launched in a quasi-polar orbit with inclination of 70.4° and height of 300 – 600 km on 15 June 2006. Period of orbital rotation is 90 min. Main tasks of the **PAMELA** experiment are study of antimatter and measurement of galactic cosmic rays in a wide energy interval. During the solar energetic particles (SEP) events, **PAMELA** is capable for the first time to measure the energy spectrum of SEPs in the range from ~ 80 MeV up to several GeV. Previously, observation of so high energy SEPs was possible only with the ground-based installations (neutron monitors). In the solar cycle 23 **PAMELA** registered only events in December

2006. For details see: [Adriani O. Barbarino G.C., Bazilevskaya G.A. et al. Observations of the 2006 December 13 and 14 solar particle events in the 80 MeV – 3 GeV range from space with the PAMELA detector // The Astrophysical Journal. 2011. V. 742. No 2. doi:10.1088/0004-637X/742/2/102.](#)

The balloons. The Catalogue utilizes results of the balloon SEP measurements in the stratosphere at the Murmansk region (Mu, cutoff rigidity $R_c = 0.6$ GV), Moscow region (Mo, $R_c = 2.4$ GV) and the Mirny observatory, Antarctica, (Mi, $R_c = 0.03$ GV). A device lifted on balloon is a telescope composed of two Geiger counters. It returns count rates of the upper counter and the telescope. Energy of solar protons is found from the particle absorption in the atmosphere. Geometric factors of both channels depend on the SEP energy spectrum. Integral fluxes of solar protons vs. energy on the atmospheric boundary are derived from these observations and presented in the Catalogue. For details see: [Bazilevskaya G.A., Makhmutov V.S., Stozhkov Y.I. et al. Solar proton events recorded in the stratosphere during cosmic ray balloon observations in 1957 – 2008 // Advances in Space Research. 2010. V. 45. Iss. 5. P. 603–613.](#)

The Neutron Monitor (NM). Till the end of 2006, high energy tail of the SEP energy spectrum could only be recorded with the ground-based installations, mainly NM. Such SEP events are conventionally called Ground Level Enhancements (GLE). Due to the geomagnetic rigidity cutoff the worldwide network of NM comprising a few tens of installations may be considered as a unique device with a huge geometric factor and good angular resolution. Neutron monitors record the secondary nucleons, mainly neutrons, generated in the atmosphere by the SEPs with energy above 500 MeV. Rather complicated calculations are needed which account for particle motion in the Earth's magnetosphere and the cascade generation of secondary particles in the atmosphere. Therefore resulting energy spectra are model dependent. The Catalogue gives the solar proton fluxes derived from the NM observations according to a model developed with participation of one of the Catalogue's authors. The model is adopted by the international project Neutron Monitor Database [<http://www.nmdb.eu/>]. For details see: [Vashenyuk E.V., Balabin Yu.V., Gvozdevsky B.B. Features of relativistic solar proton spectra derived from ground level enhancement events \(GLE\) modeling // Astrophys. Space Sci. Trans. 2011. V. 7. P. 459–463. doi:10.5194/astra-7-459-2011. <http://www.astrophys-space-sci-trans.net/7/459/2011/astra-7-459-2011.html>.](#) It must be noted that some increases of NM count rates can be caused not only by protons, but also by solar neutrons. This is observed by low latitude NMs located at large heights. Absence of neutron geomagnetic cutting and less layer of absorbing atmosphere let energetic neutrons reach monitors. These neutrons similar to protons create secondary neutrons in the matter, surrounding counters, which are registered by neutron counters after their deceleration.

Geometrical factors (G) of all instruments and their backgrounds (under quiet Sun conditions) are shown in the Table «**The Spacecrafts and Instruments**» and «**The Balloons and Neutron Monitors**».

The Spacecrafts and Instruments

Spacecraft	Instruments	E, ΔE , MeV	G, $\text{cm}^2 \cdot \text{sr}$	Background, $1/\text{cm}^2 \cdot \text{s} \cdot \text{sr}$	Work on 23-rd cycle SA
GOES - 9, 10	EPS	>5	>0.06	0.25	1997-2009
	EPS	>10	>0.06	0.15	
	EPS	>30	>0.25	0.08	
	EPS	>50	>0.25	0.06	
	EPS	>60	>0.25	0.05	
	EPS	>100	>0.25	0.03	

METEOR – 2, METEOR – 3M	CBM	>5	6.3	1.1	1997-1998,
	CBM	>15	6.3	1.1	2002-2005
	CBM	>25	6.3	1.1	
	CBM	>40	6.3	1.1	
	BP	>90	94	0.20	
	ChD	>600	314	0.19	
POES - 15, 16	MEPED	>0.24	0.06	40	1998-2009
	MEPED	>0.8	0.06	33	
	MEPED	>2.5	0.06	32	
	MEPED	>6.9	0.06	30	
	MEPED	>16	1.6	0.5	
	MEPED	>36	1.6	0.5	
	MEPED	>70	1.6	0.3	
	MEPED	>140	1.6	0.27	
CORONAS-F	MKJI	1 – 5	0.5	0.02	2001-2005
	MKJI	14 – 26	30	0.03	
	MKJI	26 – 50	2	0.05	
	MKJI	50 – 90	2	0.06	
IMP-8	CPME	1 – 2	1.51	0.02	1997-2001
	CPME	2 – 4.6	1.51	0.02	
	CPME	4.6 – 15	1.51	0.03	
	CPME	15 – 25	0.32	0.03	
	CPME	25 – 48	0.32	0.1	
	CPME	48 – 96	0.32	0.1	
	CPME	>1	1.51	0.9	
	CPME	>4	1.51	0.8	
	CPME	>10	1.51	0.75	
	CPME	>30	0.32	0.7	
	CPME	>60	0.32	0.6	
ACE	SIS	>10	≈40	0.8	1997-2009
	SIS	>30	≈40	0.5	
SOHO	LION	0.75 – 2	0.32	0.03	1997-2009
	LION	2 – 6	0.32	0.03	
	EPHIN	4 – 8	5.1	0.0004	
	EPHIN	8 – 25	5.1	0.0002	
	EPHIN	25 – 53	5.1	0.0002	
	EPHIN	>50	5.1	0.16	
STEREO-A,-B	HET	13.6 – 100	0.61	0.04 – 0.02	2006-2009
PAMELA	PAMELA	>200	20	The flux of GCR	2006-2009

The Balloons and Neutron Monitors

Instrument	Geographic observation point	E, MeV	G, sm ² ·sr	Observation time	Comments
BALLOONS	Mo Mi Mu	F(h)	~ 30 – 90 ^{*)}	1997-2009	h – altitude of observation point
NM Network	Moscow, Apatity, Mirnyi and anothers	F(R)		1997-2009	R – cutoff rigidity of protons for this NM

^{*)} Geometrical factors depend on proton energy spectrum (see [Bazilevskaya G.A., Makhmutov V.S., Stozhkov Y.I. et al. Solar proton events recorded in the stratosphere during cosmic ray balloon observations in 1957 – 2008 // Advances in Space Research. 2010. V. 45. Iss. 5. P. 603–613](#)).

Supplement 2

Determination of source increase of proton fluxes in the solar energetic particle events

Solar cosmic ray event (SCR), or Solar energetic particle (SEP) event, in the Catalog - solar proton event (SPE), is a population of energetic particles observed in the interplanetary space as a result of the explosive energy release on the Sun, mostly of solar flares accompanied by a specific set of electromagnetic radiation and coronal mass ejections (CMEs). Their relative role is a central problem of charged particle acceleration in SPEs during many years.

Since solar protons can come in the near-Earth space even from far behind the limb flare events, sometimes it's impossible to find the proper optical flash as a source of SPE. In many cases identification of the flares – sources of particles is unfortunately rather subjective. For progress in this area it is necessary to identify as precise as possible both a flare and a CME – main sources of energetic particles in the majority of SEPs. In our Catalog this subjectivity is minimized due to complex accounting for reliable statistical regularities, none of which, however, was supposed to be crucial. Three types of data were considered:

- charge-particle fluxes, their intensity-time profiles, and the energy spectra;
- electromagnetic radiation of flares in the X-ray, optical and radio ranges, their spectral and time characteristics, etc;
- magnetic field structure and evolution of the active regions on the Sun, their flare activity. [[Laurenza et al., 2007](#)].

The first type of data allows preliminary assessment of localization and time interval of a flare event, which could be a source of the proton event. The rapid (during hours) increase of the E>10 MeV proton flux up to a maximum, a hard spectrum in 10 – 100 MeV energy interval (integral spectral index $\gamma \leq 3$) more likely indicate, that a flare event took place at the Western hemisphere of the visible solar disk. Slow increase (>10 hours), wide maximum, as a rule, softer spectrum ($\gamma \geq 3$) more likely indicate the source at the Eastern hemisphere of the visible solar disk [[e.g., Lario, 2005; Laurenza et al., 2007](#)].

In this Catalog (for the 23rd SA cycle) data on dynamic features, such as solar filament eruption, limb eruptions and other phenomena in the optical diapason, were used as well. Most informative are coronal mass ejections (CMEs), the main characteristics of which are given in the Catalog. If a proton

burst event happens far behind the limb (usually the Western one), its only evidence is just existence of a CME of “halo or partial halo III” type [Park et al., 2012].

First of all, when analyzing the electromagnetic radiation of a flare event importance of the soft X-ray burst ($1 - 8 \text{ \AA}$, $E_x = 1 - 12.5 \text{ keV}$) is taken into consideration. On the visible solar disk (longitudes $E70 \div W70$) it must exceed the M1 value being usually the long lasting (> 1 hour) flares of middle importance ($M < 5$), with integral radiation flux $J/m^2 \geq 5 \cdot 10^{-2} J/m^2$. If solar flares of small X-ray importance are located on the visible solar disk, then they are accompanied by a whole range of dynamic phenomena, both in optics and radio frequencies. Data on the $H\alpha$ hydrogen line ($\lambda = 6563 \text{ \AA}$) give an accurate localization of the flare and indirectly speak about energetics and time evolution of the flare process.

A significant diagnostic factor is the radio emission of a flare event: the proton flares are usually accompanied by dynamic radio bursts of types II and/or IV, by rather intensive radio emission at centimeter ($\sim 9 \text{ GHz}$) and meter ($\sim 245 \text{ MHz}$) wave lengths and a relatively low flux density in the decimeter diapason. Duration of a microwave burst rise is usually ≥ 5 minutes and its frequency spectrum has an U-shape [Akinyan et al., 1980].

Important additional factors in the identification of a proton flux increase with a certain flare event are characteristics of active regions (ARs) or of a set of active regions (SARs), which are considered as sources of particles of the event. Proton flare events happen in the ARs, where a rapid emergence of new large magnetic fluxes occurs, converting a simple group of spots into a complex. In such ARs the flare energy release occurs usually in a series of flares of large and medium power during the limited time interval [Ishkov, 2003]. Particularly must be noted the flare activity in a SAR, consisting of two or more adjacent spot groups with common magnetic field. A big flare event usually involves the main components of the SAR, and its magnetic structure is likely to contribute to generation of large fluxes of protons.

A special class of increases of proton fluxes is caused by behind the limb flare events. The following main signatures indicate the relation of the increase to the behind the limb flare:

- presence of a high-speed wide-spread CME and/or the meter type II and/or type IV radio bursts under the absence of suitable flare events on the visible hemisphere;
- recent (within 4 days) outlet behind the western limb of a flare-active AR with a complex magnetic configuration and likely forthcoming emergence of a new magnetic flux, or being in the period of development of large flare events.
- the expected (up to 3 days) emergence of an AR from behind the eastern limb of the visible disk of the Sun, which has been active during the previous solar rotation and departed the western limb in the full development.

In case of the presence of these features it is concluded that the source of the flare event was at the invisible side of the Sun.

If proton increase is not associated with any flare activity, then it is necessary to conclude that its source is unknown. This applies mainly to low-power events.

For all events in the Catalog the reliability of the SEP source, based on the above approaches, is specified. The degree of confidence of the SEP association with the source is denoted by the corresponding symbols, which are explained in the Catalog description and are the same as in [Dodson, Hedeman, 1975].

References

Akinyan S.T., Fomichyov V.V., Chertok I.M. Results of quantitative diagnostics of proton flares according to a radio burst data for the control period of 1970 – 1977 // *Geoagnetism and Aeronomy*. 1980 V. 20. No. 3. P.385–390. (In Russian).

Dodson H.W., Hedeman E.R. Experimental comprehensive solar flare indices for certain flares, 1970 – 1974 // World Data Center A for Solar-Terrestrial Physics. Report UAG-52. Boulder, NOAA. 1975. 27 pp.

Ishkov V.N. Short term forecast of solar geoeffective flare event // Proc. of ISCS 2003 “Solar Variability as Input to the Earth’s Environment”, Tatranska Lomnitsa, Slovakia (ESA SP-535, Sept. 2003). 2003. V. 535. P. 539–540.

Lario D. Advances in modeling gradual solar energetic particle events // Advances Space Res. 2005. V. 36. P. 2279–2288. doi:10.1016/j.asr.2005.07.081
http://www.am.ub.edu/~blai/articles/AdvSpaceRes_36_12_2279-2288.pdf.

Laurenza M., Hewitt J., Cliver E.W., et al. Solar energetic proton events and soft X-ray flares // 20th ECRS Proceedings, September 5 – 8, 2006, Lisbon (Portugal). 2007.
<http://www.lip.pt/events/2006/ecrs/proc/ecrs06-s1-34.pdf>

Park J., Moon Y.-J., and Gopalswamy N. Dependence of solar proton events on their associated activities: Coronal mass ejection parameters // J. Geophys. Res. 2012. V. 117. A08108. doi:[10.1029/2011JA017477](https://doi.org/10.1029/2011JA017477).

Supplement 3

Determination of the relativistic solar proton fluxes and energy spectrum from the neutron monitor data

The main source of information on relativistic solar cosmic rays (SCR) is the ground-based installations recording so called ground-level enhancements (GLE). The only exception is an orbital spectrometer PAMELA which measures fluxes of protons and nuclei with energies above 80 MeV in the near-Earth space. Up to now PAMELA has detected the last GLE of the solar cycle 23 and the only GLE of the solar cycle 24 [Adriani et al., 2011, Bazilevskaya et al., 2013].

16 GLEs (Nos. 55 – 70) were observed in the course of the 23rd cycle of solar activity. For transition from an enhancement of the count rate of a ground-based installation to the SCR intensity on the atmospheric boundary one needs to know the response function of the instrument. Almost all information on relativistic SCR is taken from the data of neutron monitors. A number of techniques for retrieving the SCR energy spectrum utilize the world-wide network of neutron monitors as a single giant particle detector. [Shea and Smart, 1982; Cramp et al., 1997; Bieber et al., 2004; Plainaki et al., 2007; Vashenyuk et al., 2006].

Relativistic particles quickly reach the Earth's orbit, therefore the SCR fluxes usually demonstrate a strong pitch-angle anisotropy during the initial phase of the event (fast SCR component). That leads to significant difference in the records of neutron monitors with equal geomagnetic rigidity cutoffs but different asymptotic directions of arriving particles [Vashenyuk et al., 2006]. For the anisotropic phase of an event the method gives the flux value of particles coming from the most favorable direction. In the later, isotropic, phase of an event the SCR fluxes detected by different neutron monitors are uniquely dependent on the geomagnetic rigidity cutoff (delayed SCR component). The Catalog presents the SCR fluxes and spectra in the maximum of the intensity-time profile, usually not earlier than ~30 minutes after the first particle arrival, so we believe that the delayed component is more relevant to this condition although a correct approach would account for an integral of the fast component over pitch angle distribution. [Vashenyuk et al., 2011] give parameters of differential energy spectra obtained from the data of the neutron monitor network for

the following GLEs: 55 (06.11.1997), 59 (14.07.2000), 60 (15.04.2001), 61 (18.04.2001), 65 (28.10.2003), 67 (02.11.2003), 69 (20.01.2005) and 70 (13.12.2006).

In the Catalog the results are given of integrating of the delayed SCR component in the range from 1000 MeV up to E_c [Vashenyuk et al., 2006], where E_c is the energy of geomagnetic cutoff of the neutron monitor with maximum E_c [www.nmdb.eu]. The GLEs 56 – 58 (02 05 1998, 06 5 1998, 24 08 1998), 62 – 64 (04 11 2001, 26 12 2001, 24 08 2002), 66 (29 10 2003) and 68 (17 01 2005) were recorded just by a few neutron monitors and could not be processed with the usual technique of [Vashenyuk et al., 2006]. A treatment procedure for these events was taken from [Caballero-Lopez and Moraal, 2012]. Correspondence between the counting rate relative enhancement $dN(R_c, t)/N(R_c, t)$, of a neutron monitor with the geomagnetic cutoff R_c and the solar proton intensity at the atmospheric boundary $I(R, t)$ is given by a formula [Dorman, 1957]:

$$\frac{dN}{N}(R_c, t) = \frac{1}{N(R_c, t)} \int_{R_c}^{R_m} I(R, t) m(R) dR,$$

where $m(R)$ is the atmospheric yield function fitted by [Caballero-Lopez and Moraal, 2012].

The count rate of a neutron monitor $N(R_c, t)$ was taken from the latitude dependence of the neutron monitor count rates in the maximum of solar activity given in the same paper [Caballero-Lopez and Moraal, 2012]. A power-law differential energy spectrum in the range of 350 – 510 MeV measured by the HEPAD device onboard the GOES spacecraft was taken as a first approach. If the calculated dN/N was inconsistent with the observed value, the spectral index was increased until the calculated dN/N coincided with the observed one within accuracy of 0.5%. From the estimated spectrum the SCR intensity $I(>500 \text{ MeV})$ was found. It should be noted that in a weak GLE the dN/N values observed just by a few installations may differ several times for different neutron monitors probably due to contribution of the anisotropic component. Therefore the $I(>500 \text{ MeV})$ values presented in the Catalog for the GLEs 56 – 58, 62 – 64 and 66 are valid within a factor 5.

References

- Adriani O., Barbarino G.C., Bazilevskaya G.A. et al. Observations of the 2006 December 13 and 14 solar particle events in the 80 MeV n^{-1} – 3 GeV n^{-1} range from space with the PAMELA detector // *Astrophys. J.* 2011. V. 742. No. 2. doi:10.1088/0004-637X/742/2/102.
- Bazilevskaya G.A., Mayorov A.G., Mikhailov V.V. for the PAMELA Collaboration. Comparison of solar energetic particle events observed by PAMELA experiment and by other instruments in 2006 – 2012 // *Proc. ICRC 33. Rio de Janeiro. 2013. Paper 332.* <http://www.cbpf.br/~icrc2013/papers/icrc2013-0332.pdf>.
- Bieber J.W., Evenson P., Dröge W. et al. Spaceship Earth observations of the Easter 2001 Solar particle event // *Astrophys. J.* 2004. V. 601. No. 1. L103–L106. <http://dx.doi.org/10.1086/381801>.
- Caballero-Lopez R.A., Moraal H. Cosmic-ray yield and response functions in the atmosphere // *J. Geophys. Res.* 2012. V. 117. Iss. A12103. DOI: 10.1029/2012JA017794.
- Cramp J.L., Duldig M.L., Flückiger E.O. et al. The October 22, 1989, solar cosmic ray enhancement: An analysis of the anisotropy and spectral characteristics // *J. Geophys. Res.* 1997. V. 102. Iss. A11. 24237–24248. DOI: 10.1029/97JA01947.
- Dorman L.I. Cosmic Ray Variations. Washington, DC: US Department of Defence, Ohio Air-Force Base, 1957.
- Plainaki C., Belov A., Eroshenko E. et al. Modeling ground level enhancements: Event of 20 January 2005 // *J. Geophys. Res.* 2007. V. 112. Iss. A4. A04102. DOI: 10.1029/2006JA011926.

Shea M.A., and Smart D.F. Possible evidence for a rigidity-dependent release of relativistic protons from the solar corona // Space Sci. Rev. 1982. V. 32. P. 251–271.
<http://adsabs.harvard.edu/full/1982SSRv...32..251S>

Smart D.F., Shea M.A. Comment on the use of GOES solar proton data and spectra in solar proton dose calculations // Radiation Measurements. 1999. V. 30. P. 327–3359.
https://www.ngdc.noaa.gov/stp/satellite/goes/doc/goes_particles_smartshea.pdf

Vashenyuk E.V., Balabin Yu.V., Gvozdevskii B.B. et al. Relativistic solar protons in the event of January 20, 2005: Model studies // Geomagn. Aeron. 2006. V. 46. Iss. 4. P. 424–429. DOI: 10.1134/S0016793206040037.

Vashenyuk E.V., Balabin Yu.V., Gvozdevsky B.B. Features of relativistic solar proton spectra derived from ground level enhancement events (GLE) modeling // Astrophys. Space Sci. Trans. 2011. V. 7. P. 459–463. doi:10.5194/astra-7-459-2011. <http://www.astrophys-space-sci-trans.net/7/459/2011/astra-7-459-2011.html>

Supplement 4

Determination of the quasi maximal energy of protons in an event

One of parameters of a SCR event describing its power could be the maximum energy of particles observed in this event. However, a direct assessment of the maximum energy is almost impossible since there is no guarantee that particles with higher energy were not generated in this event. So, any adopted value of the maximal energy would be conventional. One should agree to accept a certain critical value of the proton flux, J_c , which will determine an upper energy for a given event. The energy value at which the proton flux J_c is observed in a given event can be conventionally considered as a maximal energy. We call this value the quasi maximal energy, E_{qm} .

Galactic cosmic rays (GCR) permanently available in space may serve as a natural critical flux value. Energy spectrum of GCR is rather stable, and its modulation by solar activity at energy below 10 GeV is well studied. Making use of GCR allows one to determine the maximal energy (actually, E_{qm}) of particles accelerated on the Sun. Such a determination is visual, it enables one to relate the SCR flux to the well-known GCR flux and roughly estimate power of solar phenomena connected with a given SEP event.

It appeared that devices that are used to detect solar protons can measure much lower proton flux than the flux of the GCR protons at the same energy. In order to approach the actual “maximal” energy we take only 10 % of the GCR flux rather than the total GCR flux for the E_{qm} determination. Since the fluxes of GCR with energy $<500 - 1000$ MeV are subject to almost two-fold variation in the 11-year solar cycle, the GCR flux in the maximum of solar activity is adopted as a fiducial one. Thus, the quasi maximal energy is determined as energy at which the proton flux is equal to fraction of 0.1 of the GCR protons. According to definition of E_{qm} , a following procedure for the E_{qm} value finding is implemented. The point of crossing of two integral spectra is found, namely the spectrum of the 0.1 fraction of the GCR flux and the SCR spectrum in a given event. At this energy $J_p(E_{SCR}) = 0.1 \cdot J_p(E_{GCR}) = J_p(E_{qm})$. The E_{qm} value for each incorporated event is given in the Catalog.

Let us note some characteristics relating to the E_{qm} of the solar proton events presented in the Catalog. First of all, it refers to the proton energy spectrum of nearly all events well fitted by a power-law function $J(>E) \sim E^{-\gamma}$ in the energy range of 10 – 100 MeV. For more powerful events the spectrum extends up to higher energies however it naturally becomes steeper with the energy growing. For example it can be fitted as $J(>E) \sim E^{-\gamma} \cdot \exp(-E/E_0)$. Unfortunately, we were not able to

fit the spectra by this function in real events of the solar cycle 23 because of paucity data for >100 MeV solar protons.

In order to account for an energy spectrum steepening while determine a E_{qm} value we used lower limits of the >100 MeV proton flux from the EPS (GOES) device or of the >1000 MeV proton flux retrieved from neutron monitor observations. In all events of the solar cycle 23 the flux values in these channels were not lower than $J_p(>100 \text{ MeV}) = 4 \cdot 10^{-3} \text{ pfu}$ and $J_p(>1000 \text{ MeV}) = 10^{-3} \text{ pfu}$. Therefore the absence of a signal in these channels meant that the actual SCR fluxes were $J_p(>100 \text{ MeV}) < 4 \cdot 10^{-3} \text{ pfu}$ or $J_p(>1000 \text{ MeV}) < 10^{-3} \text{ pfu}$. For determination of E_{qm} value we built the power-law energy spectrum using 2 points, namely $J_p(>E^*)$ where E^* was the highest actually observed energy and $J_p(>100 \text{ MeV}) = 4 \cdot 10^{-3} \text{ pfu}$ (if $E^* < 100 \text{ MeV}$) or $J_p(>1000 \text{ MeV}) = 10^{-3} \text{ pfu}$ (for $100 < E^* < 1000 \text{ MeV}$). Since in reality the >100 MeV (or >1000 MeV) protons were not detected in a given event the real spectrum was steeper than calculated ($\gamma_{\text{real}} > \gamma^*$), and the calculated E_{qm} is somewhat overestimated $E_{qm}(\text{real}) < E_{qm}(\text{calculated})$. The values of γ^* are not presented in the Catalog.

In case of the Ground Level Enhancements (GLE) observed not only by the high-latitude stations with the cutoff rigidity $\sim 1 \text{ GV}$ the best approach to a real maximal energy is the cutoff of a station at lowest latitude among all sites recorded the event. This cutoff energy is higher than E_{qm} as determined via standard procedure. Only 8 such GLE happened in the cycle 23. For uniformity of result presentation the E_{qm} values as determined via standard procedure are given in the Catalog for these events.

Thus defined the value of E_{qm} gives an idea of the power of a SEP event and enables one to have additional parameter for comparison of SEP characteristics.

▪ Supplement 5

List of symbols and abbreviations on the phenomena and events used in the Catalog materials

1. Designations in the legend of events, the overview plots and the tables of fluxes of protons

To – the time of the event beginning;

Tmax – the time of the event maximum;

Ep – the energy of solar protons;

Eqm – quasimaximal energy of protons in the event

Jmax – proton flux maximum;

Φ – integrated X-ray flux from start to $\frac{1}{2} \text{ max}$ – J/m^2 ;

pfu – particle flux unit ($1 \text{ pfu} = 1 \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$);

●, ■ – the association is certain;

⊙, □ – the association is probable;

○ – the association is possible but for some reason open to doubt;

∅ – the flare is not the main source, but contributed (or might have contributed) to the observed proton fluxes;

□ – flare-associated process (or activity) beyond the west or east limb;

◇ – active regions on the solar disk or modulation effects in the interplanetary space.

SC – suddenly commencement, the arrival of interplanetary shock into the environment space

▲ – a particle flux change associated with a SC;
Δ – SC was not affected time profiles of particle fluxes;
CME – coronal mass ejection;
V – CME median velocity – km/s;
Accel – the acceleration of CME – km/s²;
Δφ – CME angular width (degrees) near the Sun;
dA – the position angle of the first appearance of CME – the principal angle, counterclockwise from North;
gap – no observations;
V_{sw} – solar wind velocity – km/s;
B – magnetic field value in solar wind – nT;
Dst – the index of geomagnetic activity.
S/C – spacecraft
SCR – solar cosmic rays;
GLE – Ground Level Enhancement;
Location – heliographic coordinates of the events;
WL – white-light emission;

2. Designations of optical and X-ray radiations

The H α -flare characteristics as encoded in Latin characters – similarity to the Solar Geophysical Data:

A – eruptive prominence at the distance < 90° from central meridian;
B – probably the end of a more important flare;
D – bright point
E – two or more bright points;
F – several eruptive centers;
G – no visible spots in the flare neighborhood ;
H – flare accompanied by a high speed dark filament;
K – several intensity maxima;
L – existing filaments show signs of sudden activity;
M – white-light flare;
N – continuous spectrum shows effects of polarization;
O – observations have been made in Calcium II lines H or K;
P – flare shows emission in the line Helium D₃;
Q – flare shows the emission of Balmer continuum;
R – marked asymmetry in H α line suggests ejection of high velocity material;
S – brightness follows disappearance of filament in the same position;
U – two bright branches, parallel or converging;
V – occurrence of an explosive phase: important and abrupt expansion in about a minute with or without important intensity increase;
W – great increase in area after time of maximum intensity);
X – unusually wide H α line;
Y – system of loop-type prominences;
Z – major sunspot umbra covered by flare emission.

Soft X-ray burst class in the range of 1 – 8 Å (1.6 – 12.7 keV)

[<http://legacy-www.swpc.noaa.gov/weekly/index.html>]:

Classification	Peak Flux Range (0.1-0.8 nm)	
	mks system (W m^{-2})	cgs system ($\text{erg cm}^{-2}\text{s}^{-1}$)
A	$\Phi < 10^{-7}$	$\Phi < 10^{-4}$
B	$10^{-7} \leq \Phi < 10^{-6}$	$10^{-4} \leq \Phi < 10^{-3}$
C	$10^{-6} \leq \Phi < 10^{-5}$	$10^{-3} \leq \Phi < 10^{-2}$
M	$10^{-5} \leq \Phi < 10^{-4}$	$10^{-2} \leq \Phi < 10^{-1}$
X	$10^{-4} \leq \Phi$	$10^{-1} \leq \Phi$

The optical classification, observed in $\text{H}\alpha$ ($\lambda = 656.3 \text{ nm}$)

[<http://legacy-www.swpc.noaa.gov/weekly/index.html>]. Importance is the corrected area of the flare in heliographic square degrees at maximum brightness, observed in the $\text{H}\alpha$ line:

S – Subflare (area ≤ 2.0 squared deg.)

1 – Importance 1 ($2.1 \leq \text{area} \leq 5.1$ squared deg.)

2 – Importance 2 ($5.2 \leq \text{area} \leq 12.4$ squared deg.)

3 – Importance 3 ($12.5 \leq \text{area} \leq 24.7$ squared deg.)

4 – Importance 4 (area ≤ 24.8 squared deg.)

Brightness is the relative maximum brightness of flare in $\text{H}\alpha$:

F – faint;

N – normal;

B – bright

$\text{H}\alpha$ (6563 Å) dynamics phenomena:

EPL – eruptive prominence on the limb;

LPS – loop prominence system = arches flare filament system);

DSF – filament disappearance

SPY – flare spray – rapid gigantic ejection on the limb close to a flare location;

BSL – bright surge on the limb.

3. Designations of radio-bursts

The type of dynamic spectrum (DS) of the meter component radio burst:

[[Illustrated Glossary for Solar and Solar-Terrestrial Physics](#). Eds. A. Bruzek, C.J. Durrant. Springer, Netherlands, 1977. DOI: 10.1007/978-94-010-1245-4] and [<ftp://ftp.ngdc.noaa.gov/STP/space-weather/solar-data/solar-features/solar-radio/radio-bursts/reports/spectral-listings>]

DS I – noise storm bursts;

DS II – slow drift bursts;

DS III – fast drift bursts;

DS IV – prolonged broadband continuum with fine structure and slow frequency drift;

DS V – continuous broadband continuum in the meter wavelength range (normally following DS III);

CONT – broadband, smooth continuum DS IV;

DCIM – decimeter bursts defined by very fast drift spikes or groups of spikes with very high degree of polarization;

UNCLF – unclassified activity in the radio range.

The encoded spectral type of radio bursts in the decimeter and centimeter wavelength range:

P5 – means that the spectrum shows a peak at 5 GHz. P5(2.3) means that log of the maximum flux at 5 GHz is 2.3 (the maximum density of the flux is 200 units);

- 1/9** – means that the flux density is lowest at 1 GHz and rises up to 9 GHz; no information is available at frequencies below 0.6 GHz and above 9 GHz);
- 0.6\9** – means that the flux density rises toward high frequencies from 0.6 GHz to 9 GHz;
- 0.6/9** – means that the flux density falls toward high frequencies from 0.6 GHz to 9 GHz;
- U2 P7** – means that the flux density is minimal at 2 GHz and probably peaks at 7 GHz;
- 3–9** – means a flat frequency spectrum between 3 and 9 GHz.

If the element of designation is indicated in the brackets (for example [P5]), it means that this parameter is not determined reliably enough because of incompleteness or discrepancy of initial data.

4. The sources of data, used in Catalog

SGD – Solar Geophysical Data

PRF –The Preliminary Report and Forecast of Solar Geophysical Data [<http://legacy-www.swpc.noaa.gov/weekly/index.html>].

Reference list for all events of 1997-2006

Achard P., O. Adriani, M. Aguilar-Benitez et al. (The L3 collaboration). The solar flare of the 14th of July 2000 (L3+C detector results) // *Astronomy & Astrophysics*. **2006**. V. 456. No. 1. P. 351-357. doi: 10.1051/0004-6361:20065252.

Adriani O., G.C. Barbarino, G.A. Bazilevskaya et al. (Collaboration PAMELA). Observations of the 2006 December 13 and 14 solar particle events in the MeV n^{-1} – 3 GeV n^{-1} range from space with the PAMELA detector // *Astrophys. J.* **2011**. V. 742. No. 2. 11 pp. doi:10.1088/0004-637X/742/2/102.

Agueda N., R. Vainio, D. Lario, and B. Sanahuja. Solar near-relativistic electron observations as a proof of a back-scatter region beyond 1 AU during the 2000 February 18 event // *Astronomy & Astrophysics*. **2010**. V. 519. A36. 7 pp. doi: 10.1051/0004-6361/2009139634.

Al-Thoyaib S.S. Observation of the interplanetary and solar parameters in major solar-energetic particle fluxes // *J. King Saud Univ.* **2005**. V. 18. Science (1). P. 19–34, Riyadh (1426H./2005).

Arkhangelskaja I.V., A.I. Arkhangel'sky, E.V. Troitskaya, L.I. Miroshnichenko. The investigation of powerful solar flares characteristics by analysis of excited states of ^{12}C and various neutrons capture lines // *Adv. Space Res.* **2009**. V. 43. No. 4. P. 594–599.

Belov A.V., E.V. Vashenyuk, E.A. Eroshenko, V.V. Pchelkin. The increase of cosmic rays and solar wind disturbance structure on May 2, 1998 // *Solar System Research*. **2000**. V. 34. No. 2. P. 169–172. (In Russian).

Belov A.V., J.W. Bieber, E.A. Eroshenko, P. Evenson, B.B. Gvozdevsky, V.V. Pchelkin, R. Pyle, E.V. Vashenyuk, and V.G. Yanke. The “Bastille Day” GLE 14 July 2000 as observed by the worldwide neutron monitor network // *Proc. 27th Int. Cosmic Ray Conf. Hamburg, Germany*, **2001**. V. 8. P. 3446–3449.

Bieber J.W., W. Dröge, P.A. Evenson, R. Pyle, D. Ruffolo, U. Pinsook, P. Tooprakai, M. Rujiwarodom, T. Khumlumlert, and S. Krucker. Energetic particle observations during the 2000 July solar event // *Astrophys. J.* **2002**. V. 567. P. 622–634.

Bieber J.W., P. Evenson, W. Dröge, R. Pyle, D. Ruffolo, M. Rujiwarodom, P. Tooprakai, and T. Khumlumlert. Spaceship Earth observations of the easter 2001 solar particle event // *Astrophys. J.* **2004**. V. 601. P. L103–L106.

Bieber J.W., J. Clem, P. Evenson, R. Pyle, D. Ruffolo, A. Sáiz. Relativistic solar neutrons and protons on 28 October 2003 // *Geophys. Res. Lett.* **2005**. V. 32. L03S02. doi:10.1029/2004GL021492, 2005.

Bieber J.W., J. Clem, P. Evenson, R. Pyle, M. Duldig, J. Humble, D. Ruffolo, M. Rujiwarodom, and A. Sáiz. Largest GLE in half a century: Neutron monitor observations of the January 20, 2005 event // *Proc. 29th Int. Cosmic Ray Conf., Pune, India*, **2005**. V. 1. P. 237–240.

Bieber J.W., J. Clem, P. Evenson, R. Pyle, D. Ruffolo, A. Saiz, and M. Wechakama. A Maverick GLE: The relativistic solar particle event of December 13, 2006 // *Proc. 30th Int. Cosmic Ray Conf., Merida, Mexico*, **2008**. V. 1. P. 217–220.

Bieber J.W., Clem J., Evenson P., Pyle R., Sáiz A., and Ruffolo D. Giant ground level enhancement of relativistic solar protons on 2005 January 20. I. Spaceship Earth observations // *Astrophys. J.* **2013**. V. 771. 92 (13 pp). doi:10.1088/0004-637X/771/2/922,3.

Bombardieri D.J., M.L. Duldig, K.J. Michael, and J.E. Humble. Relativistic proton production during the 2000 July 14 solar event: The case for multiple source mechanisms // *Astrophys. J.* **2006**. V. 644. P. 565–574.

Bombardieri D.J., M.L. Duldig, K.J. Michael, and J.E. Humble. Relativistic proton production during the 2001 April 15 solar event // *Astrophys. J.* **2007**. V. 665. P. 813–823.

Bombardieri D.J., M.L. Duldig, J.E. Humble, and K.J. Michael. An improved model for relativistic solar particle acceleration applied to the 2005 January 20 and earlier events // *Astrophys. J.* **2008**. V. 682. P. 1315–1327.

Caroubalos C., C.E. Alissandrakis, A. Hillaris, A. Nindos, P. Tsitsipis, X. Moussas, J.-L. Bougeret, K. Bouratzis, G. Dumas, G. Kanellakis, A. Kontogeorgos, D. Maroulis, N. Patavalis, C. Perche, J. Polygiannakis, and P. Preka-Paradema. ARTHEMIS IV radio observations of the 14 July 2000 large solar event // *Solar Phys.* **2001**. V. 204. P. 167–179.

Chernetsky V.A., M.A. Livshits, L.K. Kashapova et al. Observations of X-ray flares on July 14, 2005 with Mars and near-Earth orbits // *Proc. of the Russian conference “God astronomii: Solnechnaya i solnechno-zemnaya fizika – 2009”, 5–11 July 2009. GAO RAN, Pulkovo, SPb.*, **2009**. P. 451–454. (In Russian). http://www.gao.spb.ru/russian/publ-s/conf_2009/conf_2009.pdf

Cohen C.M.S., A.C. Cummings, R.A. Leske, R.A. Mewaldt, E.C. Stone, B.L. Dougherty, M.E. Wiedenbeck, E.R. Christian, T.T. von Rosenvinge. Inferred charge state of high energy solar particles from the Solar Isotope Spectrometer on ACE // *Geophys. Res. Lett.* **1999**. V. 26. No. 2. P. 149–152.

Cohen C.M.S., G.M. Mason, R.A. Mewaldt, E.E. Cholle, E.R. Christian, A.C. Cummings, M.I. Desai, A.W. Labrador, R.A. Leske, E.C. Stone, T.T. von Rosenvinge and M.E. Wiedenbeck. Time dependent composition in the December 2006 SEP events // *Proc. 31st Int. Cosmic Ray Conf., (ID_icrc1305), Lodz, Poland*. **2009**. <http://icrc2009.uni.lodz.pl/proc/html/>

Damian A.I., M. Storini, M. Santee, S. Wang. Atmospheric impact of SEP events during last years of solar cycle 23 highlighted by MLS OH radicals // *HERPA Conference 2009 (High Energy Particle Precipitation in the Atmosphere)*. **2009**.

Danilova O.A., M.I. Tyasto, E.V. Vashenyuk, B.B. Gvozdevsky, H. Kananen, P. Tanskanen. The GLE of May 2, 1998: An effect of disturbed magnetosphere on solar cosmic rays // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA, 1999*. V. 6. P. 399–402.

Danilova O.A., M.I. Tyasto, E.V. Vashenyuk, B.B. Gvozdevsky, H. Kananen, P. Tanskanen. Magnetosphere response to a ground-based increase of SCR on May 2, 1998 // *Geomagnetism and Aeronomy*. **2002**. V. 42. No. 1. P. 32–35. (In Russian).

De Simone N., O. Adriani, G.C. Barbarino et al. (for Pamela collaboration). Study of protons of solar origin in the events of 13 and 14 December 2006 with Pamela detector // *Proc. 31st Int. Cosmic Ray Conf., Lodz, Poland. 2009*. <http://icrc2009.uni.lodz.pl/proc/html/>

Dietrich W. and C. Lopate. Measurements of iron reach SEP events using the University of Chicago IMP-8 instrument // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA, 1999*. V. 6. P. 71–74.

Dietrich W. and C. Lopate. Determination of the ionic charge states of SEPs using the University of Chicago IMP-8 instrument // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA, 1999*. V. 6. P. 91–94.

Ding L., on behalf of the L3 Collaboration. Search for possible enhancement in the flux of high energy muons due to the solar flare of 14 July 2000 with the L3+Cosmics Muon Spectrometer // *Proc. 27th Int. Cosmic Ray Conf., Hamburg, Germany. 2001*. P. 3372–3374.

Dmitriev A.V., H.-C. Yeh, J.-K. Chao, I.S. Veselovsky, S.-Y. Su, and C.C. Fu. Top-side ionosphere response to extreme solar events // *Ann. Geophys.* **2006**. V. 24. P. 1469–1477.

Duldig M.L., D.J. Bombardieri, and J.E. Humble. Further fine time resolution analysis of the Bastille Day 2000 GLE // *Proc. 28th Int. Cosmic Ray Conf., Japan, Tsukuba. 2003*. V. 6. P. 3389–3392.

El-Borie M.A. Major solar-energetic particle fluxes: I. Comparison with the associated ground level enhancements of cosmic rays // *Astroparticle Physics*. **2003a**. V. 9. No. 4. P. 549–558.

El-Borie M.A. Major solar-energetic particle fluxes: II. Comparison of the interplanetary parameters between the three largest high-energy peak flux events 19-20/10/1989, 14/7/2000 and 9/11/2000 // *Astroparticle Physics*. **2003b**. V. 19. No. 5. P. 667–677.

Falcone A.D. for Milagro collaboration. Detection of 6 November 1997 Ground Level Event by Milagrito // *CP510, The Fifth Compton Symposium*, edited by M.L. McConnell and J.M. Ryan. American Institute of Physics, **2000**. P. 574–578.

Gan W.Q. On both the time histories of the 0.511 MeV line and 2.223 MeV line from the X4.8 flare of 23 July 2002 observed with RHESSI // *Solar Phys.* **2004**. V. 219. P. 279–287.

Grechnev V.V., V.G. Kurt, I.M. Chertok *et al.* (in all 12 authors). The extreme solar event of 20 January 2005: Properties of the flare and origination of energetic particles // *Solar Phys.* **2008**. V. 252. P. 149–177.

Heber B., A. Struminsky, I. Zimovets, R. Müller-Mellin, R. Gomez-Herrero, A. Klassen, R. Wimmer-Schweingruber, C. Steigies, W. Dröge, O. Malandraki and R. Marsden. Observations of the December 2006 particle events at high latitudes with the KET aboard ULYSSES // *Proc. 30th Int. Cosmic Ray Conf., Merida, Mexico. 2008*. V. 1. P. 217–220.

Hoshida T., Y. Matsubara, Y. Muraki et al. (in all 70 authors of Tibet Collaboration). A possible observation of solar neutrons in association with November 28th, 1998

flare // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA, 1999*. V. 6. P. 38–41.
<http://www.swpc.noaa.gov/ftpdir/indices/SPE.txt>

Iles R.H., J.B.L. Jones, G.C. Taylor, J.B. Blake, R.D. Bentley, R. Hunter, L.K. Harra, and A.J. Coates. Effect of solar energetic particle (SEP) events on the radiation exposure levels to aircraft passengers and crew: Case study of 14 July 2000 SEP event // *J. Geophys. Res.* **2004**. V. 109. A11103. doi: 1029/2003JA010343.

Kallenrode M.-B. and E.W. Cliver. Rogue events: Observational aspects // *Proc. 27th Int. Cosmic Ray Conf., Hamburg, Germany, 2001*. P. 3314–3317.

Karpov S.N., E.V. Vashenyuk, V.I. Volchenko, Z.M. Karpova, L.I. Miroshnichenko, V.B. Petkov, P.S. Striganov, and A.F. Yanin. Muon intensity bursts with energy above 200 GeV during GLEs of cycles 21-23 of solar activity // *Bulletin of the Russian Academy of Sciences: Physics*. **2005**. V. 69. No. 6. P. 800–803.

Kichigin G.N., L.I. Miroshnichenko, V.I. Sidorov, S.A. Yazev. Features of a large solar event on July 23, 2002: model of the accelerated particles source // *Proc. of the Russian conference "Solar and Solar-Terrestrial Physics – 2010"*. SPb., Pulkovo, **2010**. P. 201–204. (In Russian).
http://www.gao.spb.ru/russian/publ-s/conf_2010/conf_2010.pdf

Kichigin G.N., L.I. Miroshnichenko, V.I. Sidorov, S.A. Yazev. Gamma-flares at the Sun: Dynamics of accelerated ions in eruptive and post-eruptive coronal arches // *Solnechno-Zemnaya Fizika (Solar-Terrestrial Physics)*, Irkutsk. **2012**. Iss. 21. P. 79–91. (In Russian).
http://ru.iszf.irk.ru/images/c/c4/Kichigin_13_21.pdf

Kichigin G.N., L.I. Miroshnichenko, V.I. Sidorov, and S.A. Yazev. Dynamics of accelerated ions in coronal loops and model of a gamma-ray Source // *Plasma Physics Reports*. **2014**. V. 40. No. 3. P. 178–193. © Pleiades Publishing, Ltd. ISSN 1063-780X.

Krucker S. and R.P. Lin. Two classes of solar proton events derived from onset time analysis // *Astrophys. J. Lett.* **2000**. V. 542. P. L61–L64.

Kuwabara T., K. Munakata, S. Yasue, C. Kato, S. Akahane, M. Koyama, J.W. Bieber, P. Evenson, R. Pyle, Z. Fujii, M. Tokumaru, M. Kojima, K. Marubashi, M.L. Duldig, J.E. Humble, M.R. Silva, N.B. Trivedi, W.D. Gonzalez, and N.J. Schuch. Geometry of an interplanetary CME on October 29, 2003 deduced from cosmic rays // *Geophys. Res. Lett.* **2004**. V. 31. L19803. doi:10.1029/2004GL020803.

Kuwabara T., J.W. Bieber, J. Clem, P. Evenson, and R. Pyle. Development of a ground level enhancement alarm system based upon neutron monitors // *Space Weather*. **2006**. V. 4. S10001. doi: 10.1029/2006SW000223.

Kuwabara T., J.W. Bieber, J. Clem, P. Evenson, R. Pyle, K. Munakata, S. Yasue, C. Kato, S. Akahane, M. Koyama, Z. Fujii, M.L. Duldig, J.E. Humble, R. Silva, N.B. Trivedi, W.D. Gonzalez, and N.J. Schuch. Real-time cosmic ray monitoring system for space weather // *Space Weather*. **2006**. V. 4. S08001. doi:10.1029/2005SW000204.

Kuzhevskij B.M., L.I. Miroshnichenko, and E.V. Troitskaia. Derivation of density profiles in the solar atmosphere by the 2.223 MeV line data for the 6 November 1997 flare // *Proc. 27th Int. Cosmic Ray Conf., Germany, Hamburg. Invited Rapporteur and Highlight Papers, 2001*. P. 285–288.

Kuzhevskij B.M., L.I. Miroshnichenko, and E.V. Troitskaia. Derivation of density profiles in the solar atmosphere by the 2.223 MeV line data from two flares // *Bulletin of the Russian Academy of Sciences: Physics*. **2002**. V. 66. No. 11. P. 1673–1676.

Kuzhevskij B.M., W.Q. Gan, and L.I. Miroshnichenko. The role of nuclei-nuclei interactions in the production of gamma-ray lines in solar flares // *Chin. J. Astron. Astrophys.* **2005**. V. 5. No. 3. P. 295–301.

Kuzhevskij B.M., L.I. Miroshnichenko, and E.V. Troitskaja. Gamma-ray radiation with energy of 2.223 MeV and the density distribution in the solar atmosphere during flares // *Astronomy Reports*. **2005**. V. 49. No. 7. P. 566–577. (In English).

Kuzhevskij B.M., L.I. Miroshnichenko, E.V. Troitskaia, and W. Q. Gan. Plasma density and energy spectrum of accelerated particles according to data on gamma-ray lines for three flares // *Bulletin of the Russian Academy of Sciences: Physics*. **2005**. V. 69. No. 6. P. 772–775.

Kuznetsov S.N., V.G. Kurt, B.Yu. Yushkov, I.N. Myagkova, K. Kudela, J.Kaššovicov'a, and M. Slivka. Proton acceleration during 20 January 2005 solar flare: CORONAS-F observations of high-energy emission and GLE // *Contrib. Astron. Obs. Skalnat'e Pleso*. **2006**. V. 36. P. 85–92.

Kuznetsov S.N., B.Yu. Yushkov, and K. Kudela. Measurement of the spectrum of relativistic protons from solar flares on October 28 and November 2, 2003 onboard the CORONAS-F satellite // *Cosmic Research*. **2007**. V. 45. No. 4. P. 373–375.

Lario D., R.G. Marsden, T.R. Sanderson, M. Maksimovic, A. Balogh, R.J. Forsyth, R.P. Lin, and J.T. Gosling. Ulysses and WIND particle observations of the November 1997 solar events // *Geophys. Res. Lett.* **1998**. V. 25. No. 18. P. 3469–3472.

Lario D., R.B. Decker, and A. Aran. Solar energetic particle intensities above the streaming limit // *CP1039, Particle Acceleration and Transport in the Heliosphere and Beyond – 7th Annual Astrophysics Conference*, edited by G. Li, Q. Hu, O. Verkhoglyadova, R.P. Lin, and J. Luhmann. American Institute of Physics. **2008**. P. 156–161.

Lario D., A. Aran, R.B. Decker. Major solar energetic particle events of solar cycles 22 and 23: Intensities close to the streaming limit // *Solar Phys.* **2009**. V. 260. P. 407–421. doi: 10.1007/s11207-009-9463-1.

Le G.M. The observational comparison analysis between the SEP events on 2000 July 14 and 2003 October 28 // *Proc. 29th Int. Cosmic Ray Conf., Pune, India*, **2005** (poster, not published?).

Le G.M., Yu.H. Tang and Y.B. Han. Solar energetic particle event of 2005 January 20: Release times and possible sources // *Chin. J. Astron. Astrophys.* **2006**. V. 6. No. 6. P. 751–758.

Lepping R.P., D.B. Berdichevsky, L.F. Burlaga, A.J. Lazarus, J. Kasper, M.D. Desch, C.-C. Wu, D.V. Reames, H.J. Singer, C.W. Smith and K.L. Ackerson. The Bastille Day magnetic clouds and upstream shocks: Near-Earth interplanetary observations // *Solar Phys.* **2001**. V. 204. P. 287–305.

Leske R.A., C.M.S. Cohen, A.C. Cummings, R.A. Mewaldt, E.C. Stone, B.L. Dougherty, M.E. Wiedenbeck, E.R. Christian, T.T. von Rosenvinge. Unusual isotopic composition of solar energetic particles observed in the November 6, 1997 event // *Geophys. Res. Lett.* **1999**. V. 26. No. 2. P. 153–156.

Leske R.A., R.A. Mewaldt, A.C. Cummings, E.C. Stone and T.T. von Rosenvinge. The ionic charge composition at high energies in large solar energetic particle events in solar cycle 23 //

CP598, Solar and Galactic Composition, edited by R.F. Wimmer-Schweingruber, American Institute of Physics. **2001**. P. 171–176.

Li C., Y.H. Tang, Y. Dai, W.G. Zong, and C. Fang. The acceleration characteristics of solar energetic particles in the 2000 July 14 event // *Astronomy & Astrophysics*. **2007**. V. 461. P. 1115–1119. doi: 10.1051/0004-6361:20065754.

Li C., Y. Dai, J.-C. Vial, C.J. Owen, S.A. Matthews, Y.H. Tang, C. Fang, F.N. Fazakerley. Solar source of energetic particles in interplanetary space during the 2006 December 13 event // *Astronomy & Astrophysics*. **2009**. V. 503. P. 1013–1021. doi: 10.1051/0004-6361/200911986.

Luhmann J.G., S.A. Ledvina, D. Odstrcil, M.J. Owens, X.-P. Zhao, Yang Liu, Pete Riley. Cone model-based SEP event calculations for applications to multipoint observations // *Adv. Space Res.* **2010**. V. 46 (1). P. 1–21.

Ma Yu.Q., on behalf of the L3 Collaboration. Search for a muon flux enhancement during the solar flare of 14 July 2000 with the L3+C data // *Proc. 28th Int. Cosmic Ray Conf., Japan, Tsukuba*, **2003**. V. 6. P. 3393–3396.

Malandraki O.E., N. Agueda, A. Papaioannou, K.-L. Klein, E. Valtonen, B. Heber, W. Dröge, H. Aurass, A. Nindos, N. Vilmer, B. Sanahuja, A. Kouloumvakos, S. Braune, P. Preka-Papadema, K. Tziotziou, C. Hamadache, J. Kiener, V. Tatischeff, E. Riihonen, Y. Kartavykh, R. Rodríguez-Gasén, R. Vainio. Scientific analysis within SEPServer – New perspectives in solar energetic particle research: The case study of the 13 July 2005 event // *Solar Phys.* **2012**. V. 281. P. 333–352. doi: 10.1007/s11207-012-0164-9.

Mason G.M., C.M.S. Cohen, A.C. Cummings, J.R. Dwyer, R.E. Gold, S.M. Krimigis, R.A. Leske, J.E. Mazur, R.A. Mewaldt, E. Möbius, M. Popecki, E.C. Stone, T.T. von Rosenvinge, and M.E. Wiedenbeck. Particle acceleration and sources in the November 1997 solar energetic particle events // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA*, **1999a**. V. 6. P. 115–118.

Mason G.M., C.M.S. Cohen, A.C. Cummings, J.R. Dwyer, R.E. Gold, S.M. Krimigis, R.A. Leske, J.E. Mazur, R.A. Mewaldt, E. Möbius, M. Popecki, E.C. Stone, T.T. von Rosenvinge, and M.E. Wiedenbeck. Particle acceleration and sources in the November 1997 solar energetic particle events // *Geophys. Res. Lett.* **1999b**. V. 26. No. 2. P. 141–144.

Matsubara Y., Y. Muraki, T. Sako et al. (in all 15 authors). Possible observation of solar neutrons on November 6, 1997 at Chacaltaya // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA*, **1999**. V. 6. P. 46–49.

Matsubara Y., Y. Muraki, T. Sako, K. Watanabe, S. Masuda, T. Sakai, S. Shibata, E. Flückiger, R. Butikofer, A. Chilingarian, G. Hovsepyan, Y. H. Tan, T. Yuda, M. Ohnishi, H. Tsuchiya, Y. Katayose, R. Ogasawara, Y. Mizumoto, M. Nakagiri, A. Miyashita, A. Velarde, R. Ticona, N. Martinic. Search for solar neutrons associated with proton flares in solar cycle 23 // *29th Int. Cosmic Ray Conf., Pune, India*, **2005**. V. 1. P. 17–20.

Maurchev E.A., Yu.V. Balabin, E.V. Vashenyuk, B.B. Gvozdevsky. Transport of solar protons through the atmosphere during GLE // *23rd European Cosmic Ray Symposium (and 32nd Russian Cosmic Ray Conference)*. IOP Publishing, *Journal of Physics: Conference Series* 012200. **2013**. V. 409. doi: 10.1088/1742-6596/409/1/012200.

Mazur J.E., G.M. Mason, M.D. Looper, R.A. Leske, and R.A. Mewaldt. Charge states of solar energetic particles using the geomagnetic cutoff technique: SAMPEX measurements in the 6 November 1997 solar particle event // *Geophys. Res. Lett.* **1999**. V. 26. No. 2. P. 173–176.

- Miroshnichenko L.I. High-energy cutoff for solar cosmic rays by the data of large non-standard detectors // *Bulletin of the Russian Academy of Sciences: Physics*. **2003**. V. 67. No. 4. P. 462–464.
- Miroshnichenko L.I. High-energy cutoff for solar cosmic rays by the data of large non-standard detectors // *Proc. XII Int. School "Particles and Cosmology"* (Eds: V.A. Matveev, V.A. Rubakov, Kh.S. Nirov), Moscow, INR of RAS. **2004**. P. 96–100.
- Miroshnichenko L.I. and S.N. Karpov. Cosmophysical factors and registration of rare events at the Baksan underground scintillation telescope // *Geomagnetism and Aeronomy*. **2004**. V. 44. No. 5. P. 601–606.
- Miroshnichenko L.I., K.-L. Klein, G. Trottet, P. Lantos, E.V. Vashenyuk, and Yu.V. Balabin. Electron acceleration and relativistic nucleon production in the 2003 October 28 solar event // *Adv. Space Res.* **2005**. V. 35. No. 10. P. 1864–1870.
- Miroshnichenko L.I., K.-L. Klein, G. Trottet, P. Lantos, E.V. Vashenyuk, Yu.V. Balabin, and B.B. Gvozdevsky. Relativistic nucleon and electron production in the 2003 October 28 solar event // *J. Geophys. Res.* **2005**. V. 110. No. 9. A09S08. (13 pp.). doi: 10.1029/2004JA010936.
- Miroshnichenko L.I. and J. Perez-Peraza. Astrophysical aspects in the studies of solar cosmic rays // *Internat. J. Modern Physics*. **2008**. V. 23. No. 1. P. 1–141.
- Miroshnichenko L.I., E.V. Vashenyuk, and J. Perez-Peraza. Two-component concept for ground level enhancements of solar cosmic rays: solar and interplanetary aspects // *Bulletin of the Russian Academy of Sciences: Physics*. **2009**. V. 73. No. 3. P. 297–300.
- Miroshnichenko L.I. and W.Q. Gan. Particle Acceleration and Gamma-Rays in Solar Flares: Recent Observations and New Modeling // *Adv. Space Res.* **2012**. V. 50. P. 736–756. doi: 10.1016/j.asr.2012.04.024.
- Miroshnichenko L.I., J.A. Pérez-Peraza, V.M. Velasco-Herrera, J. Zapotitla, and E.V. Vashenyuk. Oscillations of Galactic Cosmic Rays and Solar Indices before the Arrival of Relativistic Solar Protons // *Geomagnetism and Aeronomy*. **2012**. V. 52. No. 5. P. 547–560.
- Miroshnichenko L.I., E.V. Vashenyuk, J.A. Perez-Peraza. Solar Cosmic Rays: 70 Years of Ground-Based Observations // *Geomagnetism and Aeronomy*. **2013**. V. 53. No. 5. P. 541–560.
- Miroshnichenko L.I., V.I. Sidorov, Yu.V. Kuz'minykh. Solar proton event on December 14, 2006: Generation and registration of relativistic particles? // *Proc. of the Russian conference "Solar and Solar-Terrestrial Physics – 2012"* (VAK – 2013). SPb., Pulkovo, **2013**. P. 149–152. (In Russian). http://www.gao.spb.ru/russian/publ-s/conf_2013/conf_2013.pdf
- Möbius E., M. Popecki, B. Klecker, L.M. Kistler, A. Bogdanov, A.B. Galvin, D. Heitzler, D. Hovestadt, E.J. Lund, D. Morris. Energy dependence of the ionic charge state distribution during the November 1997 solar energetic particle event // *Geophys. Res. Lett.* **1999**. V. 26. No. 2. P. 145–148.
- Mu J., J.S. Wang, H.J. Zhao, J.P. Dun. A method for prediction the maximum flux of solar proton events // *Chinese J. Geophysics*. **2010**. V. 53. No. 5. P. 704–709.
- Mulligan T., J.B. Blake, R.A. Mewaldt. Unusual solar energetic particle fluxes at 1 AU within an interplanetary CME // *Proc. 30th Int. Cosmic Ray Conf., Merida, Mexico*. **2008**. V. 1. P. 179–182.
- Nitta N.V., E.W. Cliver, A.J. Tylka, and P. Smit. Source regions of major solar energetic particle events // *Proc. 28th Int. Cosmic Ray Conf., Tsukuba, Japan*. **2003**. V. 6. P. 3363–3366.

Nonaka T., Y. Hayashi, N. Ito, S. Kawakami, T. Matsuyama, A. Oshima, H. Tanaka, T. Yoshikoshi, S. K. Gupta, A. Jain, S. Karthikeyan, P. K. Mohanty, S.D. Morris, B.S. Rao, K.C. Ravindran, K. Sivaprasad, B.V. Sreekantan, S.C. Tonwar, and K. Viswanathan. Did the 28 October 2003 solar flare accelerate protons to ≥ 20 GeV? A study of the subsequent Forbush decrease with the GRAPES-3 tracking muon telescope // *Phys. Rev.* **2006**. D 74. 052003, 1550-7998. doi: 10.1103/PhysRevD.74.052003.

Perez-Peraza J., A. Gallegos-Cruz, E.V. Vashenyuk, and L.I. Miroshnichenko. Efficiency for RSP acceleration in the 14.07.2000 and 15.04.2001 events // *Proc. 28th Int. Cosmic Ray Conf., Japan, Tsukuba*. **2003**. V. 6. P. 3327-3330.

Perez-Peraza J., A. Gallegos-Cruz, E.V. Vashenyuk, Yu.V. Balabin, and L.I. Miroshnichenko. Relativistic proton production at the Sun in the October 28th, 2003 event // *Adv. Space Res.* **2006**. V. 38. No. 3. P. 418-424.

Perez-Peraza J., A. Gallegos-Cruz, L.I. Miroshnichenko, and E.V. Vashenyuk. Relativistic proton production at the Sun in the January 20, 2005 event // *Adv. Space Res.* **2008**. V. 41. No. 6. P. 947-954.

Perez-Peraza J., A. Gallegos-Cruz, E.V. Vashenyuk, L.I. Miroshnichenko, and Yu.V. Balabin. Impulsive, stochastic, and shock wave acceleration of relativistic protons in large solar events of 1989 September 29, 2000 July 14, 2003 October 28, and 2005 January 20 // *Astrophys. J.* **2009**. V. 695. No. 2. P. 865-873.

Perez-Peraza J.A., V.M. Velasco-Herrera, J. Zapotitla, L.I. Miroshnichenko, E.V. Vashenyuk. Search of periodicities in galactic space beams, solar spots and coronal index before arrival of relativistic protons from the Sun // *Bulletin of the Russian Academy of Sciences: Physics*. **2011**. V. 75. No. 6. P. 816-818. (In Russian).

Plainaki C., A. Belov, E. Eroshenko, V. Kurt, H. Mavromichalaki, V. Yanke. Unexpected burst of solar activity recorded by neutron monitors during October–November 2003 // *Adv. Space Res.* **2005**. V. 35. P. 691-696.

Quack M., M.-B. Kallenrode, M. von König, K. Künzi, J. Burrows, B. Heber, and E. Wolff. Ground level events and consequences for stratospheric chemistry // *Proc. 27th Int. Cosmic Ray Conf., Hamburg, Germany*, **2001**. P. 4023-4026.

Rawat R., S. Alex, and G.S. Lakhina. Low-latitude geomagnetic signatures during major solar energetic events of solar cycle 23 // *Ann. Geophys.* **2006**. V. 24. P. 3569-3583.

Reames D.V., C.K. Ng, A.J. Tylka. Energy-dependent ionization states of shock-accelerated particles in the solar corona // *Geophys. Res. Lett.* **1999**. V. 26. No. 24. P. 3585-3588.

Reames D.V., C.K. Ng, and A.J. Tylka. Initial time dependence of abundances in solar energetic particle events // *Astrophys. J. Lett.* **2000**. V. 531. P. L83-L86.

von Rosenvinge T.T., I.G. Richardson, D.V. Reames, C.M.S. Cohen, A.C. Cummings, R.A. Leske, R.A. Mewaldt, E.C. Stone, M.E. Wiedenbeck. The solar energetic particle event of 14 December 2006 // *Solar Phys.* **2009**. V. 256. P. 443-462. doi: 10.1007/s11207-009-9353-6.

Roussev I.I., I.V. Sokolov, T.G. Forbes, T.I. Gombosi, M.A. Lee, and J.I. Sakai. A numerical model of a coronal mass ejection: Shock development with implications for the acceleration of GeV protons // *Astrophys. J. Lett.* **2004**. V. 605. P. L73-L76.

Ryan J.M. for Milargo collaboration. Detection of 6 November 1997 ground level by Milagrito // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA*, **1999**. V. 6. P. 378-381.

Share G.H., R.J. Murphy, A.J. Tylka, R.A. Schwartz, M. Yoshimori, K. Suga, S. Nakayama, and H. Takeda. Gamma-ray line observations of the 2000 July 14 flare and SEP impact on the Earth // *Solar Phys.* **2001**. V. 204. P. 43–55.

Skoug R.M., S.J. Bame, W.C. Feldman, J.T. Gosling, D.J. McComas, J.T. Steinberg, R.L. Tokar, P. Riley, L.F. Burlaga, N.F. Ness, and C.W. Smith. A prolonged He⁺ enhancement within a coronal mass ejection in the solar wind // *Geophys. Res. Lett.* **1999**. V. 26. No. 2. P. 161–164.

Slocum P.L., E.C. Stone, R.A. Leske, E.R. Christian, C.M.S. Cohen, A.C. Cummings, M.I. Desai, J.R. Dwyer, G.M. Mason, J.E. Mazur, R.A. Mewaldt, T.T. von Rosenvinge, and M.E. Wiedenbeck. Elemental fractionation in small solar particle events // *Astrophys. J.* **2003**. V. 594. P. 592–604.

Smith C.W., N.F. Ness, L.F. Burlaga, R.M. Skoug, D.J. McComas, T.H. Zurbuchen, G. Gloeckler, D.K. Haggerty, R.E. Gold, M.I. Desai, G.M. Mason, J. E. Mazur, J.R. Dwyer, M.A. Popecki, E. Möbius, C.M.S. Cohen, and R.A. Leske. ACE observations of the Bastille Day 2000 interplanetary disturbances // *Solar Phys.* **2001**. V. 204. P. 229–254.

Somov B.V. Plasma astrophysics (2 parts): 1. Fundamentals and practice, Second Edition, 498 p. 2. Reconnection and flares, Second Edition, 504 p. Springer, SBM, New York, **2012**.

Stovpyuk M.F. and V.M. Ostryakov. Plasma parameters for the 6 November 1997 SEP event derived from the charge-consistent acceleration model // *Astron. Astrophys. Trans.* **2001**. V. 20. No. 4. P. 667–675. doi: 10.1080/10556790108221139.

Struminsky A.B. Prolonged release of > 100 MeV protons from local radiation belts of the Sun // *Proc. 26th Annual Seminar “Physics of Auroral Phenomena”, Apatity, Russia*, **2003**. P. 162–165.

Struminsky A. On possibility of prolonged two-step production of high-energy neutrons during the solar flare on 28 October 2003 // *Proc. 29th Int. Cosmic Ray Conf., Pune, India*, **2005**. V. 1. P. 45–48.

Struminsky A. Variations of solar proton spectrum during the ground level enhancement of 2005 January 20 // *Proc. 29th Int. Cosmic Ray Conf., Pune, India*, **2005**. V. 1. P. 201–204.

Struminsky A.B. Multiple proton acceleration on the Sun and free proton propagation to the Earth on January 20, 2005 // *Astronomy Letters*. **2006**. V. 32. No. 10. P. 767–776. (In Russian).

Struminsky A.B. Intensity enhancements of 200–2000 MeV protons in 1997–2005 as measured by the *KET/ULYSSES* // *Cosmic Research*. **2007**. V. 45. No. 4. P. 1. (In Russian). doi: 10.1134/S0010952507040119.

Struminsky A., B. Heber, R. Gomez-Herrero, A. Klassen. Modulation of proton fluxes at ~5 AU during the largest SEP events of 2005 // *Proc. 30th Int. Cosmic Ray Conf., Merida, Mexico*, **2008**. V. 1. P. 131–134.

Struminsky A. and I. Zimovets. On estimates of first solar proton arrival, CD-ROM, ECRS-2008, Koshice, Slovakia, **2008**.

Struminsky A.B., I.V. Zimovec. To an assessment of arrival time of the first relativistic solar protons to the Earth // *Bulletin of the Russian Academy of Sciences: Physics*. **2009**. V. 73. No. 3. P. 332–335. (In Russian).

Struminsky A.B. The dynamics and spatial distribution of solar cosmic rays in the heliosphere. Abstract thesis Dr. Sci. Moscow, IKI RAS. Moskva, IKI RAN, **2011**. 42 p. (In Russian).

- Tang Y.H., Y. Dai. Origin of coronal and interplanetary shock and particle acceleration of a flare/CME event // *Adv. Space Res.* **2003**. V. 32. No. 12. P. 2609–2612.
- Timofeev V.E. and S.A. Starodubtsev. The solar flare of April 20, 1998 by aboard Interball-2 measurements // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA, 1999*. V. 6. P. 196–199.
- Timofeev V.E. and S.A. Starodubtsev. Solar energetic particle events at the rise phase of the 23rd solar activity cycle registered aboard the spacecraft Interball-2 // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA, 1999*. V. 6. P. 200–203.
- Tonwar S., on behalf of the L3 collaboration. Variations in cosmic ray intensity observed with the L3+Cosmics shower array detectors and the intense solar flare on 14 July 2000 // *Proc. 27th Int. Cosmic Ray Conf., Hamburg, Germany, 2001*. P. 3375–3378.
- Troitskaia E.V., W.Q. Gan, B.M. Kuzhevskij, and L.I. Miroshnichenko. Solar plasma density and spectrum of accelerated particles derived from the 2.223 MeV line of a solar flare // *Solar Phys.* **2007**. V. 242. Nos. 1-2 (May 2007). P. 87–99. doi: 10.1007/s11207-007-0281z.
- Troitskaia E.V., I.V. Arkhangelskaja, L.I. Miroshnichenko, and A.I. Arkhangelsky. Study of the 28 October 2003 and 20 January 2005 solar flares by means of 2.223 MeV gamma-emission line // *Adv. Space Res.* **2009**. V. 43. No. 4. P. 547–552.
- Tylka A.J., O.E. Malandraki, G. Dorrian, Y.-K. Ko, R. Marsden, C.K. Ng, C. Tranquille. Initial Fe/O enhancements in large, gradual, solar energetic particle events: Observations from *Wind* and *Ulysses* // *Solar Phys.* **2013**. V. 285. P. 251–267. doi: 10.1007/s11207-012-0064-z.
- Vashenyuk E.V., B.B. Gvozdevsky, V.V. Pchelkin, I.G. Usoskin, K. Mursula, and G.A. Kovaltsov. The ground-level enhancement of 14 July 2000: Explaining the difference between near-by neutron monitors at Apatity and Oulu // *Proc. 27th Int. Cosmic Ray Conf., Hamburg, Germany, 2001*. P. 3383–3386.
- Vashenyuk E.V., L.I. Miroshnichenko, Yu.V. Balabin, and B.B. Gvozdevsky. Relativistic solar cosmic rays dynamics during October-November 2003 events: Modeling study // *Bulletin of the Russian Academy of Sciences: Physics.* **2005**. V. 69. No. 6. P. 808–811.
- Vashenyuk E.V., Yu.V. Balabin, J. Perez-Peraza, A. Gallegos-Cruz, and L.I. Miroshnichenko. Some features of the sources of relativistic particles at the Sun in the solar cycles 21-23 // *Adv. Space Res.* **2006**. V. 38. No. 3. P. 411–417.
- Vashenyuk E.V., Yu.V. Balabin, B.B. Gvozdevsky, and L.I. Miroshnichenko. Characteristics of relativistic SCR in large events at the Earth's surface in 1956-2005 // *Bulletin of the Russian Academy of Sciences: Physics.* **2007**. V. 71. No. 7. P. 968–971.
- Velinov P., A. Mishev. Comparison of ionization effect in the atmosphere of the Earth due to GLE 65 and GLE 69 // *23rd European Cosmic Ray Symposium (and 32nd Russian Cosmic Ray Conference)*. IOP Publishing, *Journal of Physics: Conference Series*. **2013**. V. 409. 012211. doi:10.1088/1742-6596/409/1/012211.
- Verkhoglyadova O.P., G. Li, G.P. Zank, Q.Hu, C.M.S. Cohen, R.A. Mewaldt, G.M. Mason, D.K. Haggerty, T.T. von Rosenvinge, and M.D. Looper. Understanding large SEP events with the PATH code: Modeling of the 13 December 2006 SEP event // *J. Geophys. Res.* **2010**. V. 115. A12103. doi: 10.1029/2010JA015615.
- Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk. The dynamics of the energy spectra of solar proton events in the 23rd solar cycle // *Solar System Research.* **2012**. V. 46. No. 3. P. 235–258. (In Russian).

- Wan W., L. Liu, H. Yuan, B. Ning, S. Zhang. The GPS measured SITEC caused by the very intense solar flare on July 14, 2000 // *Adv. Space Res.* **2005**. V. 36. P. 2465–2469.
- Wang R.G. Did the 2000 July 14 solar flare accelerate protons to ≥ 40 GeV? // *Astroparticle Physics*. **2009**. V. 31. P. 149–155.
- Wang R.G. and J.X. Wang. Spectra and solar energetic protons over 20 GeV in Bastille Day event // *Astroparticle Physics*. **2006**. V. 25. P. 41–46.
- Wang X. and Y. Yan. A CME-driven shock analysis of the 14-Dec-2006 SEP event // *Res. Astron. Astrophys.* V. 0. No. 0. P. 000–000. Accepted on 7 May **2012**. Available at <http://www.raa-journal.org>, <http://www.iop.org/journals/raa>. <http://arxiv.org/abs/1201.5960>
- Watari S., M. Kunitake, and T. Watanabe. The Bastille Day (14 July 2000) event in historical large Sun–Earth connection events // *Solar Phys.* **2001**. V. 204. P. 423–436.
- Yoshimori M., A. Shiozawa, and K. Suga. Solar gamma-ray flares in the 23rd solar maximum // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA, 1999*. V. 6. P. 1–49.
- Yoshimori M., A. Shiozawa, and K. Suga. Photospheric ^3He to ^1H abundance ratio derived from gamma-ray line observations // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA, 1999*. V. 6. P. 5–8.
- Yoshimori M., A. Shiozawa, and K. Suga. Low-FIP to high-FIP gamma-ray line ratio in an impulsive flare on 6 November 1997 // *Proc. 26th Int. Cosmic Ray Conf., Salt Lake City, USA, 1999*. V. 6. P. 30–33.
- Yoshimori M., K. Suga, and A. Shiozawa. Yohkoh observations of solar gamma-ray flare on November 6, 1997 // *Adv. Space Res.* **2000**. V. 25(8). P. 1801–1804.
- Yurchyshyn V., H. Wang, V. Abramenko, T.J. Spirock, S. Krucker. Magnetic field, $\text{H}\alpha$, and RHESSI observations of the 2002 July 23 gamma-ray flare // *Astrophys. J.* **2004**. V. 605. P. 546–553.
- Zhao H., G. Zhu, S. Wang, Y. Gao, Z. Liu. Relativistic electron flux enhancement at synchronous orbit during SEP event on July 14, 2000 // *Science in China (Series D)*. **2002**. V. 45. No. 5. P. 399–408.

Events in 1997

			Page
1.	Event 1997.11.04 – (1997-308) № 335		35
2.	Event 1997.11.06 – (1997-310) - GLE № 336		40
3.	Event 1997.11.13 – (1997-317) № 337		45

Particle event: To($E_p > 10$ MeV) – 04d07^h

Tmax₁($E_p > 10$ MeV) – 04d11^h, Jmax₁($E_p > 10$ MeV) – 66 /cm².s.sr

Tmax₂($E_p > 10$ MeV) – 05d02^h, Jmax₂($E_p > 10$ MeV) – 17.5 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 470 MeV


– Eqm₂ = 320 MeV

Sources: • solar flare 04d05^h52^m, X2.1/3B, S14W34, AR8100



Main burst X-ray 1–8 Å: onset – 04d05^h52^m, max – 04d05^h58^m, $\Phi = 0.056$ J/m²

CME: 04d06^h10^m, V = 0785 km/s, $\Delta\phi = 360^\circ$, dA = 243°

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1997 November 04



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1997 November 04

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-9						
EPS	>5	07 ^h	11 ^h /5d03 ^h	89/40	2d	
EPS	>10	07 ^h	11 ^h /5d02 ^h	66/17.5	2d	
EPS	>30	07 ^h	11 ^h /5d02 ^h	19.1/2.6	2d	
EPS	>50	07 ^h	09 ^h /5d02 ^h	8.6/ 1	2d	
EPS	>60	07 ^h	09 ^h /5d02 ^h	6.0/0.5	2d	
EPS	>100	07 ^h	09 ^h /5d02 ^h	2.3/0.15	2d	
METEOR-2						
CBM	>5	-	- /<5d01 ^h	- />52	-	
CBM	>15	-	- /<22 ^h	- />13	-	
CBM	>25	-	- /<21 ^h	- />3.4	-	
CBM	>40	-	- /<20 ^h	- />2.4	-	
BP	>90	06 ^h	09 ^h / -	1.7/ -	21h	
ChD	>600	06 ^h	08 ^h / -	0.07/ -	13h	
IMP-8						
CPME	>1	>06 ^h	13 ^h /5d06 ^h	105/244	2d	
CPME	>4	>06 ^h	13 ^h /5d04 ^h	56/46	2d	
CPME	>10	>06 ^h	12 ^h /5d03 ^h	29/14	2d	
CPME	>30	>06 ^h	<10 ^h /5d02 ^h	10/2.4	2d	
CPME	>60	>06 ^h	<10 ^h /5d02 ^h	5.3/1.2	2d	

ACE						
SIS	>10	05 ^h	09 ^h / -	88/35	2d	
SIS	>30	05 ^h	08 ^h / -	38/8.2	2d	
SOHO						
EPHIN (INT)	>50	06 ^h	11 ^h /05d03 ^h	8/0.3	2d	

Differential fluxes of protons for the event of 1997 November 04

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	06 ^h	18 ^h /05d21 ^h	17,9/371	2d	
CPME	2-4.6	06 ^h	14 ^h /05d14 ^h	15.1/42.1	2d	
CPME	4.6-15	06 ^h	14 ^h 05d00 ^h	2.9/3	2d	
CPME	15-25	06 ^h	13 ^h /05d00 ^h	1.1/0.6	-	
CPME	25-48	06 ^h	11 ^h / -	0.2/ -	-	
CPME	48-96	06 ^h	11 ^h / -	0.07/ -	-	
CPME	96-145	06 ^h	11 ^h / -	0.05/ -	-	
CPME	145-440	06 ^h	11 ^h / -	0.005/ -	-	
SOHO						
LION	0.75-2	08 ^h	14 ^h /05d02 ^h	103/261	2d	
LION	2-6	08 ^h	14 ^h /05d02 ^h	32/46	2d	
EPHIN	4-8	06 ^h	12 ^h /05d03 ^h	12.3/9.3	2d	
EPHIN	8-25	06 ^h	11 ^h /05d03 ^h	3/1.1	2d	
EPHIN	25-41	06 ^h	10 ^h /05d03 ^h	0.8/0,07	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References

Lario D., R.G. Marsden, T.R. Sanderson et al., 1998.
Mason G.M., C.M.S. Cohen, A.C. Cummings et al., 1999a.
Mason G.M., C.M.S. Cohen, A.C. Cummings et al., 1999b.
Möbius E., M. Popecki, B. Klecker et al., 1999.
Leske R.A., R.A. Mewaldt, A.C. Cummings et al., 2001.
Luhmann J.G., S.A. Ledvina, D. Odstrcil et al., 2010.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1997 November 04

1997 November 04


•

AR 8100

To event 335

H _α	6563 Å	0554	0559	0648	S14 W34	3B	CF
1 – 12	keV	0552	0558	0648		X2.1	5.6E-2
52.7 – 92.8	keV	<060958	~063020	064148		9	HXT Y

15.4	GHz	0556.0	0557.0	0607.0	U1.4 / 15	3.20	
8.8	GHz	0555.0	0557.0	0602.0		3.00	
5	GHz	0555.0	0557.0	0604.0		2.90	
2.7	GHz	0555.0	0557.0	0604.0		2.84	
1.4	GHz	0556.0	0557.0	0604.0		2.70	
610	MHz	0556.0	0557.0	0610.0		2.89	
410	MHz	0556.0	0558.0	0611.0		3.87	
245	MHz	0555.0	0558.0	0611.0		4.08	
DS II	30-230	0558		0607	SH	3	
DS II	30-50	0559		0606		3	
DS II	18-60	0600		0607	FN	3	
DS II	18-34	0607		0615	FN	3	
DS II	25-75	0608		0617	SH	3	
DS II	50-73	0608		0611		3	
DS IV	20-200	0606		>0709	FS	3	
DS III	25-410	0556		0558	G	3	
DS III	100-1200	0557		0602	G	2	
DS III	2000-4315	0817		0818	B	1	
DS V	35-85	0556		0614		3	
DS DCIM	1000-1525	0817		0822	G	1	
CME	WL	0610	0785km/s	-22.1 km/s ²	360°	243°	



Particle event: To($E_p > 10$ MeV) – 06d13^h

Tmax($E_p > 10$ MeV) – 7d02^h, Jmax($E_p > 10$ MeV) – 430 /cm².s.sr

Duration of the event – 7 days

Quasimaximal energy of protons in the event – $E_{qm} = 2900$ MeV


Sources: • solar flare <06d11^h22^m, 2B/X9.4, S18W63, AR8100

Main burst X-ray 1-8 Å: onset – 06d11^h49^m, max – 06d11^h55^m, Jmax = 0.036 J/m²



CME: 06d12^h10^m, V = 1556 km/s, $\Delta\phi = 360^\circ$, dA = 262°

▲ SC 06d22^h52^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1997 November 06



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1997 November 06

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES 9						
EPS	>5	13 ^h	7d07 ^h	704	7d	
EPS	>10	13 ^h	7d02 ^h	432	7d	
EPS	>30	13 ^h	21 ^h	187	7d	
EPS	>50	13 ^h	18 ^h	119	4d	
EPS	>60	13 ^h	18 ^h	95	4d	
EPS	>100	13 ^h	16 ^h	49	3d	
METEOR-2						
CBM	>5	<13 ^h	7d00 ^h	2360	7d	
CBM	>15	<13 ^h	7d00 ^h	740	7d	
CBM	>25	<13 ^h	7d00 ^h	315	6d	
CBM	>40	<13 ^h	7d00 ^h	119	4d	
BP	>90	11 ^h	-	-	-	
ChD	>600	11 ^h	-	-	-	
IMP-8						
CPME	>1	12 ^h	7d07 ^h	1840	6d	
CPME	>4	12 ^h	7d06 ^h	620	6d	
CPME	>10	11 ^h	23 ^h	348	6d	
CPME	>30	11 ^h	18 ^h	165	6d	
CPME	>60	11 ^h	18 ^h	110	6d	
ACE						
SIS	>10	12 ^h	21 ^h	670	6d	
SIS	>30	12 ^h	18 ^h	370	6d	
SOHO						
EPHIN (INT)	>50	13 ^h	18 ^h	84	6d	

BALLOONS						
Mo	>1630		16 ^h	0.023		
NM						
Network	>433		14 ^h	4.6		
Network	>1000		14 ^h	0.23		
Network	>2000		14 ^h	0.019		
Network	>3700		14 ^h	0.002		

Differential fluxes of protons for the event of 1997 November 06

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2		07d09 ^h	700	6d	
CPME	2-4.6		07d07 ^h	250	6d	
CPME	4.6-15		07d01 ^h	30	6d	
CPME	15-25		07d01 ^h	11	6d	
CPME	25-48		07d00 ^h	3	6d	
CPME	48-96		07d00 ^h	0.8	6d	
CPME	96-145		07d01 ^h	0.5	6d	
CPME	145-440		07d01 ^h	0.12	6d	
SOHO						
LION	0.75-2	19 ^h	07d06 ^h	1560	6d	
LION	2-6	19 ^h	07d06 ^h	392	6d	
EPHIN	4-8	19 ^h	07d06 ^h	97	6d	
EPHIN	8-25	19 ^h	07d02 ^h	16	6d	
EPHIN	25-41	19 ^h	07d02 ^h	2.8	6d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References

- Lario D., R.G. Marsden, T.R. Sanderson et al., 1998.
Mason G.M., C.M.S. Cohen, A.C. Cummings et al., 1999a.
Mason G.M., C.M.S. Cohen, A.C. Cummings et al., 1999b.
Möbius E., M. Popecki, B. Klecker et al., 1999.
Mazur J.E., G.M. Mason, M.D. Looper et al., 1999.
Yoshimori M., A. Shiozawa, and K. Suga, 1999.
Reames D.V., C.K. Ng, A.J. Tylka, 1999.
Matsubara Y., Y. Muraki, T. Sako et al., 1999.
Dietrich W. and C. Lopate., 1999.
Cohen C.M.S., A.C. Cummings, R.A. Leske et al., 1999.
Leske R.A., C.M.S. Cohen, A.C. Cummings et al., 1999.
Ryan J.M. for Milargo collaboration, 1999.
Yoshimori M., K. Suga, and A. Shiozawa, 2000.
Leske R.A., R.A. Mewaldt, A.C. Cummings et al., 2001.
Stovpyuk M.F. and V.M. Ostryakov, 2001
Kuzhevskij B.M., L.I. Miroshnichenko, and E.V. Troitskaia, 2002.
Nitta N.V., E.W. Cliver, A.J. Tylka et al., 2003.
Struminsky A.B., 2003.
Kuzhevskij B.M., L.I. Miroshnichenko, and E.V. Troitskaja, 2005.
Kuzhevskij B.M., L.I. Miroshnichenko, E.V. Troitskaia et al., 2005.
Wang R.G. and J.X. Wang, 2006.

Miroshnichenko L.I. and J. Perez-Peraza, 2008.
 Mu J., J.S. Wang, H.J. Zhao et al., 2010.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
 1997 November 06**

1997	November 06	•			AR 8100	To event 336	
H α	6563 Å	<1122	1156	1244	S18 W63	2B	FH
1 – 12	keV	1149	1155	1201		X9.4	3.6E-1
53 – 93	keV	1151:12	~1154	1213		1E+9	HXT2 Y
15.4	GHz	1151	1153	1209	U1.4 / 15	4.04	
8.8	GHz	1151	1153	1212		4	
5	GHz	1151	1153	1210		3.79	
2.7	GHz	1151	1153	1216		3.53	
1.4	GHz	1152	1153	1219		3.3	
610	MHz	1151	1154	1217		3.7	
410	MHz	1149	1155	1216		3.58	
245	MHz	1149	1153	1220		4.73	
DS II	45-245	1153		1216		2	
DS IV	30-80	1152		1728		3	
DS IV	35-85	1152		1446		3	
DS IV	40-800	1152		~1418	FS	3	
DS III	40-325	1134		1141	G	3	
DS III	40-800	1151		1152	G	3	
DS III	45-270	1151		1159	GG	3	
DS CONT	45-270	1152		>1308		3	
DS DCIM	2000-4295	1151		1220	GG	3	
DS DCIM	1000-2000	1152		1217	GG	3	
DS DCIM	800-1000	1152		1216		3	
CME	WL	1210	1556 km/s	-44.1km/s	360°	262°	



Particle event: To($E_p > 10$ MeV) – 13d23^h

Tmax($E_p > 10$ MeV) – 14d04^h, Jmax ($E_p > 10$ MeV) – 1.3 /cm².s.sr


Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm} = 170$ MeV



Sources: ☐ back side solar flare event < 13d22^h26^m, AR8100, 5d behind W-limb

CME: 13d22^h26^m, $V = 546$ km/s, $\Delta\phi = 288^\circ$; $dA = 310^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1997 November 13



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1997 November 13

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-9						
EPS	>5	23 ^h	14d04 ^h	2.9	2d	
EPS	>10	23 ^h	14d04 ^h	1.3	2d	
EPS	>30	23 ^h	14d00 ^h	0.3	2d	
EPS	>50	23 ^h	14d00 ^h	0.12	2d	
EPS	>60	-	14d00 ^h	0.08	2d	
EPS	>100	-	22 ^h	0.03	2d	
METEOR-2						
CBM	>5	21 ^h	22 ^h	23,7	2d	
CBM	>15	22 ^h	14d00 ^h	0.95	1d	
CBM	>25	22 ^h	14d00 ^h	0.17	1d	
CBM	>40	-	-	-	-	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
IMP-8						
CPME	>1	23 ^h	14d06 ^h	12.5	2d	
CPME	>4	23 ^h	14d03 ^h	2.5	2d	
CPME	>10	23 ^h	14d02 ^h	1.1	2d	
CPME	>30	23 ^h	14d01 ^h	0.3	2d	
CPME	>60	22 ^h	14d00 ^h	0.15	2d	
ACE						
SIS	>10	21 ^h	14d02 ^h	2.5	2d	
SIS	>30	22 ^h	14d00 ^h	0.5	2d	
SOHO						
EPHIN (INT)	>50	23 ^h	14d00 ^h	0.06	-	

Differential fluxes of protons for the event of 1997 November 13

S/c, instrument	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	23 ^h	14d11 ^h	8	~3d	
CPME	2-4.6	23 ^h	14d06 ^h	2.2	~3d	
CPME	4.6-15	21 ^h	14d02 ^h	1.8	~3d	
CPME	15-25	22 ^h	14d01 ^h	0.05	~3d	
CPME	25-48	18 ^h	14d01 ^h	0.01	~3d	
CPME	48-96	22 ^h	23 ^h	0.004	~3d	
CPME	96-145	20 ^h	14d01 ^h	0.003	~3d	
CPME	145-440	21 ^h	23 ^h	0.002	~3d	
SOHO						
LION	0.75-2	14d01 ^h	14d06 ^h	11	~3d	1.1
LION	2-6	23 ^h	14d05 ^h	2.5	~3d	0.2
EPHIN	4-8	23 ^h	14d06 ^h	0.6	~3.5d	<<
EPHIN	8-25	23 ^h	14d01 ^h	0.1	~3.5d	<<
EPHIN	25-41	23 ^h	14d00 ^h	0.01	~3.5d	<<
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	- “ -

References

Lario D., R.G. Marsden, T.R. Sanderson et al., 1998.

Krucker S. and R.P. Lin, 2000.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1997 November 13

1997 November 13



AR 8100

To event 337

Electromagnetic and corpuscular radiation it was not observed							
CME	WL	2226	546km/s	-7.4km/s	288°	310°	

Events in 1998

			Page
1.	Event 1998.04.20 – (1998-110)	№ 338	50
2.	Event 1998.04.30 – (1998-120)	№ 339	54
3.	Event 1998.05.02 – (1998-122)	№ 340	59
4.	Event 1998.05.06 – (1998-126) – GLE	№ 341	64
5.	Event 1998.05.09 – (1998-129)	№ 342	69
6.	Event 1998.06.16 – (1998-167)	№ 343	74
7.	Event 1998.08.22 – (1998-234)	№ 344	78
8.	Event 1998.08.24 – (1998-236)	№ 345	82
9.	Event 1998.09.23 – (1998-266)	№ 346	86
10.	Event 1998.09.30 – (1998-273)	№ 347	90
11.	Event 1998.10.18 – (1998-291)	№ 348	94
12.	Event 1998.11.06 – (1998-310)	№ 349	98
13.	Event 1998.11.07 – (1998-311)	№ 350	103
14.	Event 1998.11.14 – (1998-318)	№ 351	107
15.	Event 1998.11.22 – (1998-326)	№ 352	111
16.	Event 1998.11.24 – (1998-328)	№ 353	116

Particle event: To(Ep>10 MeV) – 20d11^h

Tmax₁(Ep>10 MeV) – 21d06^h, Jmax₁(Ep>10 MeV) – 860 /cm².s.sr

Tmax₂(Ep>10 MeV) – 21d12^h, Jmax₂(Ep>10 MeV) – 1600 /cm².s.sr

Duration of the event – 7 days

Quasimaximal energy of protons in the event – Eqm₁ = 440 MeV

– Eqm₂ = 600 MeV


Sources: ■ solar flare 20d09^h38^m, M1.4/EPL, s20w90, AR8194 ~2 days behind the W-limb.

Main burst X-ray 1-8 Å: onset – 20d09^h38^m, max – 20d10^h21^m, $\Phi = 0.061 \text{ J/m}^2$



CME: 20d10^h07^m, V = 1863 km/s, $\Delta\phi = 165^\circ$; dA = 284°;

ΔSC 23d18^h25^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 April 20



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 April 20

S/c, instruments	Ep, MeV	To	Tmax	Jmax /cm ² .s.sr	Dura- tion	Comments
GOES-9						
EPS	>5	11 ^h	21d06 ^h /21d12 ^h	1040/2120	7d	
EPS	>10	11 ^h	21d06 ^h /21d12 ^h	860/1600	7d	
EPS	>30	11 ^h	21d06 ^h /21d12 ^h	202/405	5d	
EPS	>50	11 ^h	21d06 ^h /21d12 ^h	47/108	4d	
EPS	>60	11 ^h	21d03 ^h /21d12 ^h	16.3/45	4d	
EPS	>100	11 ^h	21d00 ^h /21d12 ^h	2.1/6.6	3d	
METEOR-2						
CBM	>5	11 ^h	- /21d14 ^h	- /19040	7.5d	
CBM	>15	11 ^h	- /21d14 ^h	- /1640	5.5d	
CBM	>25	11 ^h	- /21d14 ^h	- /207	4.2d	
CBM	>40	11 ^h	- /21d14 ^h	- /90	3.4d	
BP	>90	11 ^h	- /21d12 ^h	- /2.5	2.2d	
ChD	>600	-	- /21d12 ^h	- /0.43	>6 h	
IMP-8						
CPME	>1	10 ^h	21d06 ^h /21d13 ^h	1360/2620	6d	
CPME	>4	10 ^h	21d06 ^h /21d13 ^h	1000/1910	6d	
CPME	>10	10 ^h	21d06 ^h /21d13 ^h	726/1300	6d	
CPME	>30	10 ^h	21d06 ^h /21d13 ^h	227/375	5d	
CPME	>60	10 ^h	21d06 ^h /21d13 ^h	86/150	4d	
ACE						
SIS	>10	10 ^h	21d03 ^h /21d10 ^h	725/1223	6d	
SIS	>30	10 ^h	21d03 ^h /21d11 ^h	220/351	5d	

SOHO						
EPHIN (INT)	>50	11 ^h	21d04 ^h /21d12 ^h	32/56	3.5d	
BALLOONS						
Mi	>122		21d(06 ^h -06 ^h)	4.1		
Mi	>149		21d(06 ^h -06 ^h)	1.41		
Mi	>206		21d(06 ^h -06 ^h)	0.34		

Differential fluxes of protons for the event of 1998 April 20

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	11 ^h	21 ^h /21d13 ^h	144/269	6d	
CPME	2-4.6	11 ^h	21d06 ^h /21d13 ^h	89/203	6d	
CPME	4.6-15	11 ^h	21d06 ^h /21d13 ^h	33/72	7d	
CPME	15-25	11 ^h	21d06 ^h /21d13 ^h	33/60.6	7d	
CPME	25-48	11 ^h	21d06 ^h /21d13 ^h	7.5/11.5	7d	
CPME	48-96	11 ^h	21d06 ^h /21d13 ^h	1.1/1.9	7d	
CPME	96-145	11 ^h	21d06 ^h /21d13 ^h	0.9/1.71	7d	
CPME	145-440	11 ^h	21d06 ^h /21d11 ^h	0.09/0.125	7d	
SOHO						
LION	0.75-2		21d05 ^h /21d11 ^h	1670/2540	6d	
LION	2-6		21d04 ^h /21d11 ^h	261/406	6d	
EPHIN	4-8	11 ^h	21d04 ^h /21d11 ^h	43/66	7d	
EPHIN	8-25	11 ^h	21d04 ^h /21d11 ^h	21/29	7d	
EPHIN	25-41	11 ^h	21d04 ^h /21d12 ^h	5.5/6	7d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References

Timofeev V.E. and S.A. Starodubtsev, 1999.

Dietrich W. and C. Lopate, 1999.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 April 20

1998	April 20	■	AR8194	To event 338			
Hα	6563 Å	No	Flare	Patrol			
EPL	6563 Å	<0931		>1000	S43 W90	3	
1 – 12	keV	0938	1021	1118		M1.4	6.1E-2
23 – 33	keV	095035	095035	095037		71	HXT Y
DS II	40-60	0956		~0959	UE		
DS III	140-170	0937		0937	G		
DS III	110-140	0950		0955	G		
DS III	40-65	1007		1012	G		
DS CONT	45-55	0957		0958			
DS CONT	55-90	1003		~1012			
DS UNCLF	110-140	0955		1001			
CME	WL	1007	1863 km/s	43.5 km/s	165°	284°	

Particle event: To(Ep>10 MeV) – 30d02^h

Tmax₁(Ep>10 MeV) – 30d15^h, Jmax (Ep>10 MeV) – 1.3 /cm².s.sr

Tmax₂(Ep>10 MeV) – 01d15^h, Jmax (Ep>10 MeV) – 1.2 /cm².s.sr *)

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 65 MeV

– Eqm₂ = 75 MeV

*) Data from IMP-8 (CPME)


Sources: • solar flare 29d16^h00^m, M6.8/3B, S16E22, AR8210

Main burst X-ray 1-8 Å: onset – 29d16^h00^m, max – 29d16^h37^m, Φ = 0.1 J/m²



CME: 30d16^h58^m, V = 1374 km/s, Δφ = 360°, dA = 336°

ΔSC 01d21^h56^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 April 30



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 April 30

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm2.s.sr	Dura- tion	Comments
GOES-9						
EPS	>5	01 ^h	15 ^h / -	16/ -	>2d	
EPS	>10	01 ^h	15 ^h / -	1.3/ -	>2d	
EPS	>30	-	15 ^h / -	0.07/ -	-	
EPS	>50	-	15 ^h / -	0.03/ -	-	
EPS	>60	-	15 ^h / -	0.01/ -	-	
EPS	>100	-	15 ^h / -	0.004/ -	-	
IMP-8						
CPME	>1	01 ^h	15 ^h /01d21 ^h	1210/1050	2.5d	
CPME	>4	01 ^h	15 ^h /01d19 ^h	38/11.9	2.5d	
CPME	>10	01 ^h	15 ^h /01d15 ^h	2.4/1.2	2.5d	
CPME	>30	01 ^h	17 ^h /01d09 ^h	0.2/0.2	2.5d	
CPME	>60	01 ^h	17 ^h /01d12 ^h	0.1/0.1	2.5d	
ACE						
SIS	>10	01 ^h	16 ^h /01d12 ^h	1.24/1.25	2.5d	
SIS	>30	01 ^h	16 ^h /01d13 ^h	0.23/0.11	2.5d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 1998 April 30

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	00 ^h	15 ^h /01d21 ^h	930/1040	2.5d	
CPME	2-4.6	01 ^h	15 ^h /01d21 ^h	112/480	2.5d	
CPME	4.6-15	01 ^h	15 ^h /01d19 ^h	2,4/0,6	2.5d	
CPME	15-25	01 ^h	17 ^h /01d08 ^h	0,03/0,05	2.5d	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	01 ^h	15 ^h /01d21 ^h	1110/1360	2.5d	
LION	2-6	01 ^h	15 ^h /01d21 ^h	83/63	2.5d	
EPHIN	4-8	01 ^h	14 ^h /01d18 ^h	5.3/1.5	2.5d	
EPHIN	8-25	01 ^h	14 ^h /01d15 ^h	0.11/0.04	2.5d	
EPHIN	25-41	01 ^h	14 ^h /01d12 ^h	0.002/0.0005	2.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 April 30

1998 April 29


•

AR8210

To event 339

H _{α}	6563 Å	1605	1630	2104	S16 E22	3B	FTUZ
1 – 12	keV	1606	1637	1659		M6.8	1.0E-1
50 – 100	keV	<163927	~163931	1702		334	HXT Y
5.4	GHz	1631	1631	1632		1.72	
8.8	GHz	1630	1631	1631		1.78	
6.7	GHz	1612.7	1630	1638.9		2.24	
5	GHz	1629	1630	1636	U2.7 P5	2.32	
2.7	GHz	1629	1630	1636		2	
1.4	GHz	1619	1624	1636		2.4	

610	MHz	1624	1626	1636		2.7	
410	MHz	1624	1632	1636		2.67	
280	MHz	1623.4	1632	1710		2.94	
245	MHz	1627	1630	1636		3.15	
DS II	40-90	1622		1632	SH,H	3	
DS II	40-90	1637		1700	UE,H	3	
DS IV	40-800	1613		1702		3	
DS IV	35-85	1702		1720		3	
DS III	40-400	1607		1614	GG	3	
DS III	40-170	1700		1701	G	3	
DS V	30-80	1613		1619		2	
DS DCIM	450-750	1610		1610		2	
DS DCIM	800-2000	1613		1653	GG,FS	3	
DS DCIM	2000-4395	1613		1630	GG	2	
CME	WL	1658	1374 km/s	-44.8rm/s ²	360°	336°	



Particle event: To($E_p > 10$ MeV) – 02d14^h

Tmax($E_p > 10$ MeV) – 02d16^h, Jmax ($E_p > 10$ MeV) – 130 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm} = 800$ MeV


Sources: • solar flare 02d13^h31^m, X1.1/3B, S15W15, AR8210

Main burst X-ray 1-8 Å: onset – 02d13^h31^m, max – 02d13^h42^m, $\Phi = 0.067$ J/m²



CME: 02d14^h06^m, V = 0938 km/s, $\Delta\phi = 360^\circ$, dA = 331°

Δ SC 03d17^h43^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 May 02



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 May 02

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Duration	Comments
GOES-9						
EPS	>5	14 ^h	16 ^h	181	3d	
EPS	>10	14 ^h	16 ^h	130	3d	
EPS	>30	14 ^h	16 ^h	46	3d	
EPS	>50	14 ^h	16 ^h	21	3d	
EPS	>60	14 ^h	16 ^h	15	3d	
EPS	>100	14 ^h	15 ^h	6.8	2d	
METEOR-2						
CBM	>5	14 ^h	16 ^h	343	2.9d	
CBM	>15	14 ^h	17 ^h	87	2.4d	
CBM	>25	14 ^h	17 ^h	28	2d	
CBM	>40	14 ^h	15 ^h	23	1.9d	
BP	>90	14 ^h	15 ^h	>4.7	1.7d	
ChD	>600	14 ^h	<17 ^h	0.19	1.1d	
IMP-8						
CPME	>1	14 ^h	21 ^h	250	4d	
CPME	>4	14 ^h	19 ^h	75	3d	
CPME	>10	14 ^h	18 ^h	47	3d	
CPME	>30	14 ^h	17 ^h	22	3d	
CPME	>60	14 ^h	16 ^h	15	3d	

ACE						
SIS	>10	14 ^h	17 ^h	116	3d	
SIS	>30	14 ^h	16 ^h	47	3d	
SOHO						
EPHIN (INT)	>50	14 ^h	16 ^h	7.2	2,5d	
BALLOONS						
Mi	>116		3d(07 ^h -08 ^h)	1.1		After
Mi	>136		3d(07 ^h -08 ^h)	0.61		maximum
Mi	>183		3d(07 ^h -08 ^h)	0.26		
NM						
Network	>500		15 ^h	0.25		

Differential fluxes of protons for the event of 1998 May 02

S/c, instrument	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	14 ^h	03d04 ^h	131	3d	
CPME	2-4.6	15 ^h	21 ^h	29	3d	
CPME	4.6-15	14 ^h	18 ^h	2,9	3d	
CPME	15-25	15 ^h	20 ^h	1,4	3d	
CPME	25-48	15 ^h	17 ^h	0,8	3d	
CPME	48-96	14 ^h	18 ^h	0,5	2,8d	
CPME	96-145	14 ^h	16 ^h	0,08	2,8d	
CPME	145-440	14 ^h	15 ^h	0,03	2,8d	
SOHO						
LION	0.75-2	15 ^h	21 ^h	280	3,5d	
LION	2-6	15 ^h	21 ^h	67	4d	
EPHIN	4-8	14 ^h	20 ^h	6.5	3,5d	
EPHIN	8-25	15 ^h	19 ^h	1.3	3,5d	
EPHIN	25-41	14 ^h	16 ^h	0.22	3,5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References

Timofeev V.E. and S.A. Starodubtsev, 1999.
 Danilova O.A., M.I. Tyasto, E.V. Vashenyuk et al., 1999.
 Danilova O.A., M.I. Tyasto, E.V. Vashenyuk et al., 2002.
 Belov A.V., E.V. Vashenyuk, E.A. Eroshenko et al., 2000.
 Dietrich W. and C. Lopate, 1999.
 Skoug R.M., S.J. Bame, W.C. Feldman et al., 1999.
 Struminsky A.B., 2003.
 Roussev I. I., I.V. Sokolov, T.G. Forbes et al., 2004.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
1998 May 02**

1998 May 02 • AR8210 To event 340

H _α	6563 Å	1334	1342	1540	S15 W15	3B	FZ
1 – 12	keV	1331	1342	1351		X1.1	6.7E-2
57 – 100	keV	133422	133730	>133742		524	HXT Y
15.4	GHz	1337	1341	1513		3.11	
8.8	GHz	1337	1341	1513		3.23	
5	GHz	1335	1338	1513	U2.7 [P5]	3.26	
2.7	GHz	1337	1338	1513		3.11	
610	MHz	1334	1344	1513		4.34	
410	MHz	1334	1345	1505		4.3	
245	MHz	1333	1338	1511		4.15	
DS IV	35-85	1338		1723		3	
DS DCIM	800-2000	1334		1355	GG,FS,SP	3	
DS DCIM	2000-4395	1336		1348	GG,FS	3	
CME	WL	1406	0938 km/s	-28.8km/s ²	360°	331°	



Particle event: To($E_p > 10$ MeV) – 06d08^h

Tmax($E_p > 10$ MeV) – 06d09^h, Jmax ($E_p > 10$ MeV) – 120 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm} = 575$ MeV


Sources: • solar flare 06d07^h58^m, X2.7/1N, S15W64, AR8210

Main burst X-ray 1-8 Å: onset – 06d07^h58^m, max – 06d08^h09^m, $\Phi = 0.21$ J/m²



CME: 06d08^h29^m, $V = 1099$ km/s, $\Delta\phi = 190^\circ$, $dA = 309^\circ$;

ΔSC 08d09^h52^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 May 06



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 May 06

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Duration	Comments
GOES-9						
EPS	>5	08 ^h	10 ^h	171	2d	
EPS	>10	08 ^h	09 ^h	120	2d	
EPS	>30	08 ^h	09 ^h	30.1	2d	
EPS	>50	08 ^h	09 ^h	10.3	2d	
EPS	>60	08 ^h	09 ^h	7.0	2d	
EPS	>100	08 ^h	09 ^h	2.5	2d	
METEOR-2						
CBM	>5	08 ^h	09 ^h	414	3d	
CBM	>15	08 ^h	09 ^h	82	2d	
CBM	>25	08 ^h	09 ^h	15	1.5d	
CBM	>40	08 ^h	09 ^h	8,4	1d	
BP	>90	08 ^h	09 ^h	1,7	1d	
ChD	>600	-	-	-	-	
IMP-8						
CPME	>1	08 ^h	11 ^h	170	2.5d	
CPME	>4	08 ^h	10 ^h	146	2.5d	
CPME	>10	08 ^h	09 ^h	84	2.5d	
CPME	>30	08 ^h	09 ^h	26.9	2.5d	
CPME	>60	08 ^h	09 ^h	13.8	2.5d	
ACE						
SIS	>10	08 ^h	09 ^h	402	2.5d	
SIS	>30	08 ^h	09 ^h	103	2.5d	
SOHO						
EPHIN (INT)	>50	09 ^h	09 ^h	15,4	2d	

BALLOONS						
Mi	>116		3d(07 ^h -08 ^h)	1.1		
Mi	>136		3d(07 ^h -08 ^h)	0.61		
Mi	>183		3d(07 ^h -08 ^h)	0.26		
NM						
Network	>500		09 ^h	0.039		

Differential fluxes of protons for the event of 1998 May 06

S/c, instrument	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	11 ^h	15 ^h	187	3d	
CPME	2-4.6	08 ^h	11 ^h	27	3d	
CPME	4.6-15	08 ^h	10 ^h	9.3	3d	
CPME	15-25	08 ^h	09 ^h	4.65	3d	
CPME	25-48	08 ^h	09 ^h	0.65	3d	
CPME	48-96	08 ^h	09 ^h	0.12	3d	
CPME	96-145	08 ^h	09 ^h	0.15	3d	
CPME	145-440	08 ^h	09 ^h	0.02	2.5d	
SOHO						
LION	0,75-2	09 ^h	16 ^h	1140	3d	
LION	2-6	09 ^h	14 ^h	204	2.5d	
EPHIN	4-8	09 ^h	11 ^h	46	2.5d	
EPHIN	8-25	09 ^h	11 ^h	12.5	2.5d	
EPHIN	25-41	08 ^h	10 ^h	2,1	2.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References


Dietrich W. and C. Lopate, 1999.
Reames D.V., C.K. Ng, and A.J. Tylka, 2000.
Struminsky A.B., 2003.
Miroshnichenko L.I. and J. Perez-Peraza, 2008.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 May 06

1998 May 06 • AR8210 To event 341

H _{α}	6563 Å	0801	0805	>0918	S15 W64	1N	
1 – 12	keV	0758	0809	0820		X2.7	2.1E-1
57 – 100	keV	075728	080610	>090818		1020	HXT Y
200 – 10000	keV	0808					GRS Y

15.4	GHz	801	807	824		3.18	
8.8	GHz	801	807	822	U2.8 P8.8	3.3	
5.7	GHz	646.1	809.2	<1030.1		3.06	
2.8	GHz	711	804	957		2.68	
1.4	GHz	801	804	828		2.71	
500	MHz	800	815	905		2.92	
410	MHz	801	811	828		3.08	
245	MHz	800	806	806		4.81	
200	MHz	803	806	811		2.97	
DS II	45-270	805		819	HARM	3	
DS IV	300-1100	810		817		2	
DS III	25-160	801		804	G	3	
DS CONT	45-65	801		802		2	
DS DCIM	800-2000	800		821	GG,FS,SP	3	
DS DCIM	2000-4395	801		828	GG,FS	3	
CME	WL	0829	1099 km/s	24.5km/s ²	190°	294°	



Particle event: To($E_p > 10$ MeV) – 09d06^h

Tmax₁($E_p > 10$ MeV) – 09d13^h, Jmax₁ ($E_p > 10$ MeV) – 4.7 /cm².s.sr

Tmax₂($E_p > 10$ MeV) – 09d23^h, Jmax₂ ($E_p > 10$ MeV) – 8.9 /cm².s.sr

Duration of the event – 2.5 days

Quasimaximal energy of protons in the event – Eqm₁ = 230 MeV


– Eqm₂ = 230 MeV

Sources: ■ solar flare 09d03^h04^m, M7.7/-, s15w90*, AR8210 ~ 1 day behind the W-limb; Main burst X-ray 1-8 Å: onset – 09d03^h04^m, max – 09d03^h40^m, $\Phi = 0.11$ J/m²



CME: 09d03^h35^m, V = 2331 km/s, $\Delta\phi = 178^\circ$, dA = 262°;

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 May 09



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 May 09

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-9						
EPS	>5	06 ^h	13 ^h /23 ^h	7.3/19	2.5d	
EPS	>10	06 ^h	13 ^h /23 ^h	4.7/8.9	2.5d	
EPS	>30	06 ^h	13 ^h /22 ^h	1.3/1.4	2.5d	
EPS	>50	06 ^h	12 ^h /20 ^h	0.4/0.4	2.5d	
EPS	>60	08 ^h	12 ^h /20 ^h	0.24/0.25	2d	
EPS	>100	08 ^h	11 ^h /20 ^h	0.07/0.1	2d	
METEOR-2						
CBM	>5	<17 ^h	- /18 ^h	- /28	3d	
CBM	>15	<17 ^h	- /18 ^h	- /2.9	2d	
CBM	>25	<17 ^h	- /17 ^h	- /0.18	1d	
CBM	>40	<17 ^h	- /17 ^h	- /0.09	1d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
IMP-8						
CPME	>1	05 ^h	12 ^h /10d01 ^h	17.2/105	5d	
CPME	>4	05 ^h	12 ^h /10d01 ^h	7.2/27	3.5d	
CPME	>10	05 ^h	12 ^h /10d00 ^h	4.0/7.3	3.5d	
CPME	>30	05 ^h	12 ^h /21 ^h	1.3/1.6	3.5d	
CPME	>60	05 ^h	12 ^h /20 ^h	0.8/0.7	3.5d	
ACE						
SIS	>10	05 ^h	12 ^h /23 ^h	4.8/10.7	2d	
SIS	>30	05 ^h	12 ^h /19 ^h	1.3/1.7	2d	

SOHO						
EPHIN (INT)	>50	04 ^h	11 ^h /21 ^h	0.1/0.1	1d	

Differential fluxes of protons for the event of 1998 May 09

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	08 ^h	- /10d01 ^h	- /45	3.5d	
CPME	2-4,6	07 ^h	- /10d01 ^h	- /16	3.5d	
CPME	4,6-15	06 ^h	- /10d00 ^h	- /1.74	3d	
CPME	15-25	06 ^h	- /10d00 ^h	- /0.3	3d	
CPME	25-48	05 ^h	- /21 ^h	- /0.04	2.5d	
CPME	48-96	06 ^h	- /17 ^h	- /0.01	2.5d	
CPME	96-145	05 ^h	- /21 ^h	- /0.01	1.5d	
CPME	145-440	05 ^h	- /11 ^h	- /0.003	1.5d	
SOHO						
LION	0,75-2	05 ^h	12 ^h /10d01 ^h	13/85	4d	
LION	2-6	06 ^h	12 ^h /10d01 ^h	4/20	4d	
EPHIN	4-8	05 ^h	11 ^h /23 ^h	0.9/5	4d	
EPHIN	8-25	05 ^h	11 ^h /21 ^h	0.15/0.6	4d	
EPHIN	25-41	05 ^h	11 ^h /21 ^h	0.026/0.04	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References



Dietrich W. and C. Lopate, 1999.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 May 09

1998	May 09	■ AR8210			To event 342		
H _α	6563 Å	No Flare Patrol			s15w90*		
1 – 12.5	keV	0304	0340	0355		M7.7	1.1E-1
57 – 100	keV	031733	032641	045755		673	HXT, Y
57 – 100	keV	031733	032641	045755		243	HXT Y
8.8	GHz	0323	0323	0325		1.79	
5	GHz	0323	0323	0332		2.18	
2.7	GHz	0322	0323	0347	[P2.7]	2.38	
1.4	GHz	0322	0324	0346		2.23	
245	MHz	0323	0324	0330		2.53	
DS II	23-75	0326		0329	FN	3	
DS II	50-150	0326		0329	SH	3	
DS III	30-80	0314		0331		3	
DS III	18-310	0322		0328	G	2	

15.4	GHz	0335	0336	0339		1.72	
8.8	GHz	0335	0337	0339		1.89	
5.7	GHz	0310	0337	0353		2.1	
5	GHz	0323	0337	0347		2.28	
610	MHz	0323	0345	0352	P0.6	3.4	
410	MHz	0323	0345	0352		3.15	
DS II	20-75	0331		0341	FN	3	
DS II	40-150	0331		0342	SH	3	
DS IV	200-1000	0323		0350		2	
DS IV	30-80	0331		0413		3	
DS III	25-75	0320		0348	N	2	
CME	WL	0335	2331km/s	-140.5km/s ²	178°	275°	

* – probable localization of the flare event



Particle event: To($E_p > 10$ MeV) – 16d21^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 17\text{d}09^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 1.3 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 18\text{d}02^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 1.4 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{\text{qm}1} = 80 \text{ MeV}$


– $E_{\text{qm}2} = 75 \text{ MeV}$

Sources: ■ solar flare 16d18^h03^m, M1.0/ -, s22w90*, AR8232 ~ 1 day behind the W-limb; Main burst X-ray 1-8 Å: onset – 16d18^h03^m, max – 16d18^h42^m, $\Phi = 0.037 \text{ J/m}^2$



CME: 16d18^h27^m, $V = 1484 \text{ km/s}$, $\Delta\phi = 281^\circ$, $dA = 278^\circ$;

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 June 16



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 June 16

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Duration	Comments
GOES-9						
EPS	>5	-	17d09 ^h /18d05 ^h	2.9/3.5	3d	
EPS	>10	-	17d09 ^h /18d02 ^h	1.3/1.4	3d	
EPS	>30	-	17d09 ^h /18d02 ^h	0.15/0.22	2d	
EPS	>50	-	17d04 ^h /18d02 ^h	0.06/0.07	2d	
EPS	>60	-	17d04 ^h /18d02 ^h	0.04/0.04	-	
EPS	>100	-	-	-	-	
IMP-8						
CPME	>1	20 ^h	17d10 ^h /18d01 ^h	157/200	6d	
CPME	>4	20 ^h	17d10 ^h /18d01 ^h	5.5/6	5d	
CPME	>10	20 ^h	17d10 ^h /18d01 ^h	1.2/1.2	5d	
CPME	>30	20 ^h	17d10 ^h /18d01 ^h	0.13/0.3	2d	
CPME	>60	20 ^h	17d09 ^h /18d01 ^h	0.06/0.05	2d	
ACE						
SIS	>10	20 ^h	17d10 ^h /18d01 ^h	2.2/1.5	2d	
SIS	>30	20 ^h	17d10 ^h /18d01 ^h	0,6/0.2	2d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 1998 June 16

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	23 ^h	18d01 ^h /18d23 ^h	178/184	4.5d	
CPME	2-4.6	23 ^h	17d11 ^h /18d23 ^h	12.5/8.6	4.5d	
CPME	4.6-15	21 ^h	18d01 ^h / -	0.3/ -	4.5d	
CPME	15-25	23 ^h	17d15 ^h / -	0.05/ -	4.5d	
CPME	25-48	23 ^h	17d23 ^h / -	0.01/ -	4.5d	
CPME	48-96					
CPME	96-145					
CPME	145-440					
SOHO						
LION	0,75-2	23 ^h	17d16 ^h /18d23 ^h	330/225	5d	
LION	2-6	23 ^h	17d16 ^h /18d23 ^h	26/9.8	5d	
EPHIN	4-8	21 ^h	17d10 ^h /18d03 ^h	1.2/0.8	5d	
EPHIN	8-25	20 ^h	17d10 ^h /18d03 ^h	0.09/0.1	5d	
EPHIN	25-41	20 ^h	17d10 ^h /18d03 ^h	0.004/0.003	5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 June 16

1998

June 16



AR8232

To event 343

H α	6563 Å	No Flare					
LPS	6563 Å	1956		2202	S17 W90		
1 – 12	keV	1803	1842	1928	s22w90*	M1.0	3.7E-2
CME	WL	1827	1484 km/s	-74.7km/s ²	341°	278°	

* – probable localization of the flare event

Particle event: To(Ep>10 MeV) – 22d06^h

Tmax₁(Ep>10 MeV) – 23d00^h, Jmax₁(Ep>10 MeV) – 1.7 /cm².s.sr

Tmax₂(Ep>10 MeV) – 23d08^h, Jmax₁(Ep>10 MeV) – 1.5 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 80 MeV


– Eqm₂ = 85 MeV

Sources: • solar flare 21d23^h57^m, M9.0/2B, N42E51, AR8307


Main burst X-ray 1-8 Å: onset – 21d23^h57^m, max – 22d00^h09^m, Φ = 0.061 J/m²

CME: gap

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 August 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 August 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	06 ^h	23d00 ^h /23d08 ^h	7/7.6	2d	
EPS	>10	06 ^h	23d00 ^h /23d08 ^h	1.7/1.5	2d	
EPS	>30	06 ^h	23d00 ^h /23d09 ^h	0.14/0.1	1d	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
IMP-8						
CPME	>1	06 ^h	23d00 ^h /23d08 ^h	146/165	3d	
CPME	>4	06 ^h	23d00 ^h /23d08 ^h	13.6/14.4	3d	
CPME	>10	06 ^h	23d00 ^h /23d08 ^h	2.15/2.25	3d	
CPME	>30	06 ^h	23d00 ^h /23d08 ^h	0.16/0.2	2d	
CPME	>60	06 ^h	23d00 ^h /23d08 ^h	0.07/0.11	2d	
ACE						
SIS	>10	06 ^h	23d00 ^h /23d08 ^h	2.3/2.6	2d	
SIS	>30	06 ^h	23d01 ^h /23d08 ^h	0.3/0.45	2d	


Differential fluxes of protons for the event of 1998 August 22

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	-	- /23d08 ^h	- /110	-	
CPME	2-4.6	-	- /23d08 ^h	- /18.5	-	
CPME	4.6-15	-	- /23d07 ^h	- /0.9	-	
CPME	15-25	-	- /23d07 ^h	- /0.06	-	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO		No	data			

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 August 22

1998		August 22		•		AR8307		To event 344	
Hα	6563 Å	<0009	0016	>0140	N42 E51	2B			
1 – 12	keV	2357	0009	0016		M9.0		6.1E-2	

15.4	GHz	2359.0	0003.0	0017.0	2.8 / 15	3.20	
8.8	GHz	2359.0	0003.0	0013.0		2.89	
2.8	GHz	2358.0	0000.0	0100.0		2.19	
DS II	20-120	0005		0023	FN	3	
DS II	40-280	0005		0017	SH	3	
DS IV	30-80	0004		0129		2	
DS III	18-300	0007		0009	G	2	
CME	WL						gap



Particle event: To($E_p > 10$ MeV) – 24d23^h

Tmax₁($E_p > 10$ MeV) – 25d02^h, Jmax₁($E_p > 10$ MeV) – 96 /cm².s.sr

Tmax₂($E_p > 10$ MeV) – 26d07^h, Jmax₂($E_p > 10$ MeV) – 320 /cm².s.sr

Duration of the event – 6 days


Quasimaximal energy of protons in the event – Eqm₁ = 720 MeV

– Eqm₂ = 310 MeV


Sources: • solar flare 24d21^h48^m, 3B/X1.0, N35E09, AR8307

Main burst X-ray 1-8 Å: onset – 24d21^h50^m, max – 24d22^h12^m, $\Phi = 0.16$ J/m²

CME: gap




Time profiles of the proton fluxes for the event of 1998 August 24



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 August 24

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	23 ^h	25d02 ^h /26d07 ^h	158/1730	6d	
EPS	>10	23 ^h	25d02 ^h /26d07 ^h	96/320	6d	
EPS	>30	23 ^h	25d02 ^h /26d04 ^h	25/34	3d	
EPS	>50	23 ^h	25d02 ^h /26d04 ^h	10/6.7	2d	
EPS	>60	23 ^h	25d02 ^h /26d04 ^h	7.5/3	2d	
EPS	>100	23 ^h	25d01 ^h /26d04 ^h	3.2/0.3	2d	
METEOR-2						
CBM	>5	-	-	-	-	
CBM	>15	-	-	-	-	
CBM	>25	23 ^h	25d < 22 ^h /-	>19/ -	5d	
CBM	>40	23 ^h	25d < 21 ^h /-	>18.5/ -	3d	
BP	>90	-	25d < 21 ^h /-	>0.7/ -	-	
ChD	>600	-	-	-	-	
POES-15						
MEPED	>0.24	23 ^h	25d00 ^h /26d07 ^h	630/108600	-	
MEPED	>0.8	23 ^h	25d00 ^h /26d07 ^h	460/28000	-	
MEPED	>2.5	23 ^h	25d00 ^h /26d07 ^h	350/8000	-	
MEPED	>6.9	23 ^h	25d00 ^h /26d07 ^h	300/700	-	
MEPED	>16	23 ^h	25d00 ^h /26d07 ^h	65.1/135.1	-	
MEPED	>36	23 ^h	25d00 ^h /26d07 ^h	30/20.5	-	
MEPED	>70	23 ^h	25d00 ^h /26d07 ^h	5.23/1.25	-	
MEPED	>140	23 ^h	25d00 ^h /26d07 ^h	2.24/0.28	-	
IMP-8						
CPME	>1	>20 ^h	25d08 ^h /26d07 ^h	1180/16000	7d	
CPME	>4	>20 ^h	25d08 ^h /26d07 ^h	270/2590	7d	
CPME	>10	>20 ^h	25d08 ^h /26d07 ^h	90/455	5d	
CPME	>30	>20 ^h	25d08 ^h /26d07 ^h	20,3/50,	4d	
CPME	>60	>20 ^h	25d08 ^h /26d07 ^h	10.5/29	4d	
ACE						
SIS	>10	23 ^h	25d03 ^h / -	150/ -	-	
SIS	>30	23 ^h	25d01 ^h / -	42/ -	-	
BALLOONS						
Mi	>154		25d(08 ^h -09 ^h)/ -	2.7/ -		
Mi	>215		25d(08 ^h -09 ^h)/ -	0.8/ -		
Mi	>357		25d(08 ^h -09 ^h)/ -	0.16/ -		
Mi	>150		- /26d(07 ^h -07 ^h)	- /0.2		
Mi	>164		- /26d(07 ^h -07 ^h)	- /0.14		
Mi	>189		- /26d(07 ^h -07 ^h)	- /0.05		
NM						
Network	>500		25d01 ^h / -	0.22/ -		

Differential fluxes of protons for the event of 1998 August 24

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						

CPME	1-2	>20 ^h	25d08 ^h /26d07 ^h	479/8340	6d	
CPME	2-4.6	>20 ^h	25d08 ^h /26d07 ^h	192/2220	6d	
CPME	4.6-15	>20 ^h	25d08 ^h /26d07 ^h	17,4/179	6d	
CPME	15-25	>20 ^h	<25d04 ^h /26d07 ^h	3,4/14,7	2d	
CPME	25-48	>20 ^h	<25d04 ^h /26d07 ^h	0,5/1,0	2d	
CPME	48-96	>20 ^h	<25d04 ^h /26d07 ^h	0,14/0,6	2d	
CPME	96-145	>20 ^h	<25d04 ^h /26d07 ^h	0,09/0,12	2d	
CPME	145-440	>20 ^h	<25d04 ^h /26d07 ^h	0,01/0,003	2d	
SOHO		No	data			


References

Dietrich W. and C. Lopate, 1999.
Leske R.A., R.A. Mewaldt, .C. Cummings et al., 2001.
Struminsky A.B, 2003.
Miroshnichenko L.I. and J. Perez-Peraza. 2008.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 August 24

1998 August 24 • AR8307 To event 345

H α	6563 Å	2148	2204	>25 ^d 0108	N35 E09	3B	T
1 – 12	keV	2150	2212	2235		X1.0	1.6E-1
53 – 93	keV	<2218	~2218	2248		894	HXT Y
15.4	GHz	2200.0	2203.0	2302.0		3.08	
8.8	GHz	2200.0	2203.0	2302.0		3.32	
5	GHz	2159.0	2203.0	2231.0	U0.6 P5	3.69	
2.8	GHz	2150.0	2205.0	>2232.0		3.43	
1.4	GHz	2157.0	2210.0	2231.0		2.97	
610	MHz	2159.0	2203.0	2235.0		2.46	
410	MHz	2158.0	2203.0	2236.0		2.70	
245	MHz	2158.0	2203.0	2302.0		2.92	
DS II	18-140	2203		2218	SH,H	3	
DS IV	20-200	2203		>2400	FS	2	
DS III	70-700	2158		2205	G	2	
CME	WL						gap



Particle event: To($E_p > 10$ MeV) – 23d13^h

Tmax($E_p > 10$ MeV) – 25d01^h, Jmax ($E_p > 10$ MeV) – 22 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm} = 75$ MeV


Sources: ● solar flare 23d06^h40^m, M7.1/3B, N19E09, AR8340

Main burst X-ray 1-8 Å: onset – 23d06^h40^m, max – 23d07^h13^m, $\Phi = 0.12$ J/m²


CME: gap

▲ SC 24d23^h45^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 September 23



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 September 23


S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	13 ^h	25d01 ^h	324	3d	
EPS	>10	13 ^h	25d01 ^h	22	3d	
EPS	>30	-	-	-	-	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-15						
MEPED	>0.24		25d01 ^h	43280		
MEPED	>0.8		25d01 ^h	6880		
MEPED	>2.5		25d01 ^h	1290		
MEPED	>6.9		25d01 ^h	40		
MEPED	>16		25d01 ^h	1.9		
MEPED	>36		-	-		
MEPED	>70		-	-		
MEPED	>140		-	-		
IMP-8						
CPME	>1	13 ^h	25d01 ^h	14100	6d	
CPME	>4	13 ^h	25d01 ^h	826	6d	
CPME	>10	13 ^h	25d01 ^h	29	6d	
CPME	>30	13 ^h	25d01 ^h	1.6	6d	
CPME	>60	13 ^h	25d01 ^h	1.4	6d	
ACE						
SIS	>10	13 ^h	25d00 ^h	21	3d	
SIS	>30	13 ^h	24d19 ^h	0.36	3d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 1998 September 23

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	24d15 ^h	25d00 ^h	$9.4 \cdot 10^3$	7d	
CPME	2-4.6	24d15 ^h	25d01 ^h	$1.6 \cdot 10^3$	7d	
CPME	4.6-15	24d16 ^h	25d01 ^h	54.8	7d	
CPME	15-25	24d17 ^h	25d01 ^h	0.4	-	
CPME	25-48	24d18 ^h	26d00 ^h	0.007	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO		No	data			

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
1998 September 23**

1998	September 23	•	AR8340	To event 346			
H α	6563 Å	0643	0701	0937	N19E09	3B	CEFHU
1 – 12	keV	0640	0713	0731		M7.1	1.2E-1
53 – 93	keV	>070434	~070436	>112234		697	HXT Y
15.4	GHz	0652.0	0656.0	0723.0		3.00	
8.8	GHz	0651.0	0656.0	0741.0		3.18	
5	GHz	0649.0	0656.0	0755.0		3.15	
2.8	GHz	0620.0	0701.0	0809.0	[U2.8 P8.8]	3.05	
1.4	GHz	0647.0	0654.0	0743.0		3.15	
610	MHz	0652.0	0656.0	0747.0		3.26	
410	MHz	0652.0	0709.0	0748.0		3.32	
245	MHz	0655.0	0711.0	0747.0		3.53	
204	MHz	0653.5	0714.0	0724.5		2.70	
DS II	40-280	0656		~0719	SH,H	3	
DS II	18-50	0704		0714	FN	2	
DS IV	30-900	0653		>0725		2	
DS III	200-800	0722		0723	B	2	
DS DCIM	800-2000	0643		0737	GG,SP	3	
DS DCIM	2000-4395	0717		0742	GG	2	
410	MHz	0814.0	0818.0	0820.0		3.11	
245	MHz	0815.0	0818.0	0820.0		3.57	
DS DCIM	800-2000	0801		0835	GG	3	
CME	WL						gap



Particle event: To($E_p > 10$ MeV) – 30d14^h

Tmax($E_p > 10$ MeV) – 30d23^h, Jmax ($E_p > 10$ MeV) – 785 /cm².s.sr

Duration of the event – 3 days


Quasimaximal energy of protons in the event – $E_{qm} = 600$ MeV

Sources: • solar flare 30d13^h08^m, M2.8/2N, N23W78, AR8340


Main burst X-ray 1-8Å: onset – 30d13^h08^m, max – 30d13^h50^m, $\Phi = 0.11$ J/m²

CME: gap

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 September 30



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 September 30

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	14 ^h	01d03 ^h	1620	3d	
EPS	>10	14 ^h	23 ^h	785	3d	
EPS	>30	14 ^h	20 ^h	90	3d	
EPS	>50	14 ^h	19 ^h	24	2d	
EPS	>60	14 ^h	19 ^h	12.5	2d	
EPS	>100	14 ^h	19 ^h	2.7	1d	
POES-15						
MEPED	>0.24	-	21 ^h	3050	-	
MEPED	>0.8	-	20 ^h	2970	-	
MEPED	>2.5	-	19 ^h	2500	-	
MEPED	>6.9	-	18 ^h	1590	-	
MEPED	>16	-	18 ^h	480	-	
MEPED	>36	-	17 ^h	97	-	
MEPED	>70	-	17 ^h	4.8	-	
MEPED	>140	-	16 ^h	1.15	-	
IMP-8						
CPME	>1	14 ^h	21 ^h	1500	5d	
CPME	>4	14 ^h	21 ^h	910	5d	
CPME	>10	14 ^h	21 ^h	460	5d	
CPME	>30	14 ^h	20 ^h	95	3d	
CPME	>60	14 ^h	20 ^h	36	3d	
ACE						
SIS	>10	14 ^h	20 ^h	685	3d	
SIS	>30	14 ^h	19 ^h	144	3d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 1998 September 30

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	15 ^h	01d15 ^h	2000	~7d	
CPME	2-4.6	15 ^h	01d15 ^h	603	~7d	
CPME	4.6-15	15 ^h	21 ^h	48	~7d	
CPME	15-25	15 ^h	21 ^h	23	~4d	
CPME	25-48	15 ^h	20 ^h	3.1	~3d	
CPME	48-96	15 ^h	19 ^h	0.37	~2d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO		Data	absent			


References

Dietrich W. and C. Lopate, 1999.
Reames D.V., C.K. Ng, and A.J. Tylka, 2000.
Leske R.A., R.A. Mewaldt, C. Cummings et al., 2001.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
1998 September 30**

1998 September 30 • AR8340 To event 347

H α	6563 Å	1402	1434	1542	N23 W78	2N	FY
1 – 12	keV	1308	1350	1448		M2.8	1.1E–1
57-100	keV	<1314	~1343	>1344		354	HXT Y
15.4	GHz	1327.0	1331.0	0000.0		2.04	
8.8	GHz	1325.0	1331.0	1350.0		2.49	
5	GHz	1317.0	1334.0	1405.0	U0.4 P2.7 - 5	2.65	
2.7	GHz	1312.0	1330.0	1405.0		2.69	
1.4	GHz	1317.0	1332.0	1405.0		2.46	
610	MHz	1312.0	1317.0	0000.0		2.26	
410	MHz	1312.0	1317.0	1410.0		1.81	
245	MHz	1310.0	~1329.0	1405.0		2.54	
245	MHz	1251.0	1329.0	0000.0		2.54	
DS II	35-85	1322		1330		3	
DS IV	40-800	1310		~1500		3	
DS III	40-90	1307		1312	G	2	
DS V	30-63	1308		1315		2	
DS DCIM	800-2000	1308		1355	GG	3	
DS DCIM	2000-4400	1309		1355	GG	3	
CME	WL						gap



Particle event: To(Ep>10 MeV) – 18d22^h

Tmax₁(Ep>10 MeV) – 19d02^h Jmax₁(Ep>10 MeV) – 1.8 /cm².s.sr

Tmax₂(Ep>10 MeV) – 19d06^h Jmax₂(Ep>10 MeV) – 2.3 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 140 MeV

– Eqm₂ = 80 MeV

Sources: o DSF 15d<10^h05^m, N19E10, 27°


o solar flare 18d01^h38^m, M2.4/2B, N16W53, AR8358

Main burst X-ray 1-8Å: onset – 18d01^h38^m, max – 18d01^h45^m, Φ = 0.001 J/m²



CME: 15d10^h05^m, V = 199 km/s, Δφ = 360°, dA = 264°

▲ SC 18d19^h52^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 October 18



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 October 18

S/c, instrument	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	22 ^h	19d02 ^h /19d09 ^h	2.8/5.3	2d	
EPS	>10	22 ^h	19d02 ^h /19d06 ^h	1.8/2.3	2d	
EPS	>30	22 ^h	19d02 ^h /19d06 ^h	0.38/0.25	1d	
EPS	>50	23 ^h	19d01 ^h /19d06 ^h	0.13/0.07	1d	
EPS	>60	23 ^h	19d01 ^h /19d06 ^h	0.08/0.04	1d	
EPS	>100	23 ^h	19d01 ^h / -	0.03/ -	1d	
IMP-8						
CPME	>1	22 ^h	19d02 ^h /19d09 ^h	5/77	2d	
CPME	>4	22 ^h	19d02 ^h /19d06 ^h	2.8/18.8	2d	
CPME	>10	22 ^h	19d02 ^h /19d06 ^h	1.7/5.1	2d	
CPME	>30	22 ^h	19d01 ^h /19d06 ^h	0.6/0.8	2d	
CPME	>60	22 ^h	19d01 ^h /19d06 ^h	0.4/0.4	2d	
ACE						
SIS	>10	22 ^h	19d01 ^h /19d06 ^h	1.75/5.3	2d	
SIS	>30	22 ^h	19d01 ^h /19d06 ^h	0.7/1	2d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 1998 October 18

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	19d06 ^h	- /19d06 ^h	- /33	~2d	
CPME	2-4.6	19d03 ^h	- /19d06 ^h	- /12	~2d	
CPME	4.6-15	21 ^h	19d02 ^h /19d06 ^h	0.1/1.2	~2d	
CPME	15-25	22 ^h	19d02 ^h /19d06 ^h	0.07/0.2	~2d	
CPME	25-48	21 ^h	19d01 ^h /19d06 ^h	0.02/0.02	~2d	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	19d05 ^h	- /19d07 ^h	- /6.1	1.5d	
LION	2-6	19d05 ^h	- /19d07 ^h	- /1.4	1.5d	
EPHIN	4-8	23 ^h	19d01 ^h /19d06 ^h	0.2/2.6	2.5d	
EPHIN	8-25	22 ^h	19d01 ^h /19d06 ^h	0.06/0.3	2.5d	
EPHIN	25-41	22 ^h	19d01 ^h /19d06 ^h	0.01/0.017	2.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 October 18

1998 October 15 ○ AR To event 348

DSF	6562 Å	<1005		<1101	N19E10	27°	*
1 – 12	keV						
CME	WL	15 ^d 1004	0199 km/s	3.2 km/s	360°	264°	

The spacecraft detected an interplanetary shock passage at 18/1902UT related to a halo-CME produced by a large filament disappearance on 15 October – PRF1207, 1208.

1998 October 18 ○ AR8358 To event 348

H α	6563 Å	0142	0144	0202	N16 W53	2B	EFH
1 – 12	keV	0138	0145	0152		M2.4	9.9E–3
410	MHz	0158	0318		85	110	
CME	WL						gap

Particle event: To(Ep>10 MeV) – 06d03^h

Tmax(Ep>10 MeV) – 06d12^h, Jmax(Ep>10 MeV) – 4.6 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – E_{qm} = 75 MeV


Sources: • solar flare 05d19^h00^m, M8.4/2B, N22W18, AR8375

Main X-ray burst 1-8 Å: onset – 05d19^h00^m, max – 05d19^h55^m, Φ = 0.11 J/m²



CME: 05d20^h44^m, V = 1118 km/s, Δφ = 360°, dA = 300°

ΔSC 07d08^h15^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 November 06



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 November 06

S/c, instrument	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES-10						
EPS	>5	03 ^h	12 ^h	50.5	1.5d	
EPS	>10	03 ^h	12 ^h	4.6	1.5d	
EPS	>30	03 ^h	12 ^h	0.12	1.5d	
EPS	>50	03 ^h	11 ^h	0.05	1.5d	
EPS	>60	03 ^h	11 ^h	0.02	1.5d	
EPS	>100	-	-	-	-	
POES-15						
MEPED	>0.24	-	12 ^h	4360	1.5d	
MEPED	>0.8	-	12 ^h	2890	1.5d	
MEPED	>2.5	-	12 ^h	760	1.5d	
MEPED	>6.9	-	12 ^h	16	1.5d	
MEPED	>16	-	12 ^h	1	1.5d	
MEPED	>36	-	12 ^h	0.2	1.5d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	01 ^h	12 ^h	1790	1.5d	
CPME	>4	01 ^h	12 ^h	110	1.5d	
CPME	>10	01 ^h	12 ^h	4.5	1.5d	
CPME	>30	01 ^h	12 ^h	0.2	1.5d	
CPME	>60	01 ^h	12 ^h	0.1	-	
ACE						
SIS	>10	07 ^h	10 ^h	6	1.5d	
SIS	>30	07 ^h	08 ^h	0.15	1.5d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 1998 November 06

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	01 ^h	12 ^h	754	1.5d	
CPME	2-4.6	01 ^h	12 ^h	173	1.5d	
CPME	4.6-15	01 ^h	12 ^h	6.7	1.5d	
CPME	15-25	01 ^h	12 ^h	0.05	1.5d	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	02 ^h	17 ^h	200	2d	
LION	2-6	01 ^h	17 ^h	30.2	2d	
EPHIN	4-8	00 ^h	16 ^h	24.8	1.5d	
EPHIN	8-25	00 ^h	12 ^h	0.58	1.5d	
EPHIN	25-41	-	-	-	-	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	



Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 November 06

1998 November 05 • AR8375 To event 349

H α	6563 Å	<1839*	1950	>2334	N22 W18	2B	T
1 – 12	keV	1900	1955	2012		M8.4	1.1E–1
>300 keV	keV	193749	194343	201520		742.61	BATS C
15.4	GHz	1941.0	1944.0	2048.0		2.72	
8.8	GHz	1941.0	1943.0	2006.0		2.94	
6.7	GHz	1937.9	1952.2	2046.9	U0.6 P6.7	3.19	
5	GHz	1941.0	1952.0	2053.0		2.94	
2.7	GHz	1941.0	1952.0	2055.0		2.60	
1.4	GHz	1942.0	1946.0	2055.0		2.18	
610	MHz	1945.0	1954.0	2055.0		1.84	
410	MHz	1944.0	1945.0	2053.0		2.34	
280	MHz	1951.0	2005.2			2.59	
DS II	25-75	1951		1957		2	

8.8	GHz	2026.0	2027.0	2045.0		1.98	
5	GHz	2016.0	2026.0	2047.0	U0.6 P1.4-5	2.11	
2.7	GHz	2016.0	2035.0	2047.0		2.04	
1.4	GHz	2015.0	2026.0	2047.0		2.04	
610	MHz	2025.0	2027.0	2047.0		1.71	
410	MHz	2015.0	2015.0	2047.0		1.87	
280	MHz	1951.0	2005.2			2.59	
DS IV	25-300	<2015		2142		2	
DS III	18-180	2017		2037	N	2	
DS CONT	25-55	2010		0337		1	
CME	WL	2044	1118 km/s	-24.0 km/s ²	360°	300°	

* – No Flare Patrol 5^d 1729 – 1839



Particle event: To($E_p > 10$ MeV) – 07d12^h

$T_{\max 1}(E_p > 10 \text{ MeV})$ – 07d14^h, $J_{\max 1}(E_p > 10 \text{ MeV})$ – $2.8 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV})$ – 08d02^h, $J_{\max 2}(E_p > 10 \text{ MeV})$ – $6 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm1} = 75 \text{ MeV}$

– $E_{qm2} = 80 \text{ MeV}$


Sources: ● solar flare 07d11^h06^m, M2.4/SN, N14W43, AR8375

Main X-ray burst 1-8 Å: onset – 07d11^h02^m, max – 07d11^h06^m, $\Phi = 0.005 \text{ J/m}^2$



CME: 07d11^h54^m, $V = 0632 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 258^\circ$

▲ SC 08d04^h51^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 November 07



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 November 07

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES-10						
EPS	>5	12 ^h	14 ^h /08d02 ^h	15.3/36	3d	
EPS	>10	12 ^h	14 ^h /08d02 ^h	2.8/6	3d	
EPS	>30	12 ^h	14 ^h /08d01 ^h	0.25/0.22	3d	
EPS	>50	12 ^h	14 ^h / -	0.03/0.02	2d	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-15						
MEPED	>0.24	-	- /08d03 ^h	- /5338	3d	
MEPED	>0.8	-	- /08d03 ^h	- /2078	3d	
MEPED	>2.5	-	- /08d03 ^h	- /329	3d	
MEPED	>6.9	-	- /08d03 ^h	- /10.5	2d	
MEPED	>16	-	- /08d03 ^h	- /2.85	2d	
MEPED	>36	-	- /08d03 ^h	- /0.51	2d	
MEPED	>70	-	- /08d03 ^h	- /0.2	2d	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	12 ^h	13 ^h /8d01 ^h	1310/686	3d	
CPME	>4	12 ^h	13 ^h /8d01 ^h	30/45	3d	
CPME	>10	12 ^h	13 ^h /8d01 ^h	4.3/6.1	2d	
CPME	>30	12 ^h	13 ^h /8d01 ^h	0.3/0.45	1.5d	
CPME	>60	12 ^h	13 ^h /8d01 ^h	0.1/0.25	1.5d	

ACE						
SIS	>10	12 ^h	13 ^h /8d01 ^h	2.2/5.4	1.5d	
SIS	>30	12 ^h	13 ^h /8d01 ^h	0.2/0.3	1d	
SOHO						
EPHIN (INT)	>50	-	14 ^h /08d01 ^h	0.02/0.025	-	

Differential fluxes of protons for the event of 1998 November 07

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	-	13 ^h / -	1130/ -	5d	
CPME	2-4.6	-	13 ^h / -	84/ -	5d	
CPME	4.6-15	-	13 ^h / -	1.8/ -	5d	
CPME	15-25	-	13 ^h / -	0.08/ -	5d	
CPME	25-48	-	13 ^h / -	0,008/ -	5d	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	-	- /08d04 ^h	- /99	5d	
LION	2-6	-	- /08d04 ^h	- /11	5d	
EPHIN	4-8	14 ^h	15 ^h /08d01 ^h	4.4/8.5	5d	
EPHIN	8-25	12 ^h -	14 ^h /08d01 ^h	0.13/0.4	5d	
EPHIN	25-41	12 ^h -	14 ^h /08d01 ^h	0.005/0.008	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 November 07

1998 November 07 • AR8375 To event 350

Hα	6563 Å	1104	1108	1116	N14 W43	SN	C
1 – 12.5	keV	1102	1106	1116		M2.4	4.9E-3
53 – 93	keV	<110522	110558	111100		788	HXT Y
>300	MeV	110410	110556	110934			BATS C
3	GHz	1104.5	1106.0	1108.3		1.70	
204	MHz	1104.6	1104.8	1107.8		4.51	
DS DCIM	2000-4375	1104		1106	G	2	
DS DCIM	800-2000	1104		1107	G	2	
CME	WL	1154	0632 km/s	1.3 km/s	360°	258°	

Particle event: To(Ep>10 MeV) – 14d06^h

Tmax₁(Ep>10 MeV) – 14d12^h, Jmax₁ (Ep>10 MeV) – 250 /cm².s.sr

Tmax₂(Ep>10 MeV) – 15d06^h, Jmax₂ (Ep>10 MeV) – 10 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 580 MeV

– Eqm₂ = 190 MeV


Sources: ☐ solar flare 14d05^h00^m, C1.3/, N28W90*, AR8375 ~ 2 days behind W-limb;

Main X-ray burst 1-8 Å: onset – 14d05^h00^m, max – 14d05^h08^m, $\Phi = 0.013 \text{ J/m}^2$



CME: gap

* – localization on BSL

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 November 14



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 November 14

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	06 ^h	12 ^h /15d02 ^h	442/12.5	3.5d	0.35
EPS	>10	06 ^h	12 ^h /15d02 ^h	250/10	3d	0.21
EPS	>30	06 ^h	12 ^h /15d02 ^h	64/3.8	2d	0.12
EPS	>50	06 ^h	12 ^h /15d02 ^h	24/0.8	2d	0.09
EPS	>60	06 ^h	12 ^h / -	15/ -	2d	0.08
EPS	>100	06 ^h	09 ^h / -	5.1/ -	2d	0.05
POES-15						
MEPED	>0.24	-	(11-14) ^h / -	1375/ -	4d	
MEPED	>0.8	-	(11-14) ^h / -	1122/ -	4d	
MEPED	>2.5	-	(11-14) ^h / -	843/ -	3.5d	
MEPED	>6.9	-	(11-14) ^h / -	544/ -	3d	
MEPED	>16	-	(11-14) ^h / -	170/ -	2d	
MEPED	>36	-	(11-14) ^h / -	64.2/ -	2d	
MEPED	>70	-	(11-14) ^h / -	9.67/ -	2d	
MEPED	>140	-	(11-14) ^h / -	1.5/ -	2d	
IMP-8						
CPME	>1	05 ^h	14 ^h /15d03 ^h	550/305	>5d	
CPME	>4	05 ^h	14 ^h /15d03 ^h	305/134	>5d	
CPME	>10	05 ^h	14 ^h /15d03 ^h	173/58	5d	
CPME	>30	05 ^h	13 ^h /15d03 ^h	64/14.6	4d	
CPME	>60	05 ^h	13 ^h /15d03 ^h	37/7.6	4d	
ACE						
SIS	>10	05 ^h	12 ^h /15d02 ^h	192/75	2d	
SIS	>30	05 ^h	12 ^h /15d02 ^h	92/18	2d	

SOHO						
EPHIN (INT)	>50	06 ^h	10 ^h / -	15.5/ -		
BALLOONS						
Mi	>110	-	08 ^h -12 ^h / -	2.5/ -	-	
Mi	>184	-	08 ^h -12 ^h / -	0.5/ -	-	
Mi	>214	-	08 ^h -12 ^h / -	0.1/ -	-	
Mi	>107	-	- /15d(03 ^h -03 ^h)	- /0.08	-	
Mi	>114	-	- /15d(03 ^h -03 ^h)	- /0.07	-	

Differential fluxes of protons for the event of 1998 November 14

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	05 ^h	14 ^h /15d03 ^h	101/78	~2,5d	
CPME	2-4.6	05 ^h	14 ^h /15d03 ^h	65/43	~2,5d	
CPME	4.6-15	05 ^h	14 ^h /15d03 ^h	14.2/7.8	~2,5d	
CPME	15-25	05 ^h	14 ^h /15d03 ^h	6.4/2.4	~2,5d	
CPME	25-48	05 ^h	14 ^h /15d03 ^h	1.3/0.34	~2,5d	
CPME	48-96	05 ^h	14 ^h /15d03 ^h	0.3/0.075	~2,5d	
CPME	96-145	05 ^h	14 ^h /15d03 ^h	0.2/0.08	~2,5d	
CPME	145-440	05 ^h	14 ^h /15d03 ^h	0.05/0.01	~2,5d	
SOHO						
LION	0,75-2	05 ^h	14 ^h / -	46/ -	-	
LION	2-6	05 ^h	14 ^h / -	9.7/ -	-	
EPHIN	4-8	05 ^h	13 ^h / -	30/ -	-	
EPHIN	8-25	05 ^h	13 ^h / -	7.3/ -	-	
EPHIN	25-41	05 ^h	13 ^h / -	1.25/ -	-	

References

Dietrich W. and C. Lopate, 1999.

Leske R.A., R.A. Mewaldt, C. Cummings et al., 2001.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 November 14

1998 November 14 ☐ AR8375 To event 351

Hα	6563 Å	No Flare					
BSL	6563 Å	0518		0543	N28 W90		
1 -12.5	keV	0500	0508	0515		C1.3	6.1E-04
33 – 53	keV	0503:11	0505:13	0506:23		339	HXT Y
5.7	GHz	0500.3	0505.1	0532.3		0.90	
245	MHz	0502.0	0502.0	0503.0		2.04	
DS II	35-160	0506		0524	SH	3	
DS II	18-70	0506		0528	FN	3	
DS IV	30-80	0501		0536		3	
DS III	18-230	0502		0506	G	2	
CME	WL						gap

Particle event: To($E_p > 10$ MeV) – 22d07^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 22\text{d}09^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 1.1 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 22\text{d}14^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 0.6 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 1.5 day

Quasimaximal energy of protons in the event – $E_{\text{qm}1} = 285 \text{ MeV}$


– $E_{\text{qm}2} = 160 \text{ MeV}$

Sources: • solar flare 22d06^h30^m, X3.7/1N, S27W82, AR8384



Main burst X-ray 1-8 Å: onset – 22d06^h30^m, max – 22d06^h42^m, $\Phi = 0.2 \text{ J/m}^2$

CME: gap

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 November 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 November 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	07 ^h	09 ^h /14 ^h	1.33/0.9	1d	
EPS	>10	07 ^h	09 ^h /14 ^h	1.1/0.6	1d	
EPS	>30	07 ^h	08 ^h /13 ^h	0.5/0.2	1d	
EPS	>50	07 ^h	08 ^h /13 ^h	0.25/0.08	-	
EPS	>60	07 ^h	08 ^h /13 ^h	0.15/0.05	-	
EPS	>100	07 ^h	08 ^h /13 ^h	0.1/0.02	-	
POES-15						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	08 ^h / -	2.2/ -	-	
MEPED	>36	-	08 ^h / -	1.6/ -	-	
MEPED	>70	-	08 ^h / -	0.33/ -	-	
MEPED	>140	-	08 ^h / -	0.2/ -	-	
IMP-8						
CPME	>1	07 ^h	12 ^h /16 ^h	10/10.3	2d	
CPME	>4	07 ^h	10 ^h /16 ^h	3.8/1.9	2d	
CPME	>10	07 ^h	09 ^h /16 ^h	1.8/0.65	1,5d	
CPME	>30	07 ^h	09 ^h /16 ^h	0.6/0.3	1d	
CPME	>60	07 ^h	09 ^h /16 ^h	0.25/0.1	1d	
ACE						
SIS	>10	07 ^h	09 ^h /16 ^h	4.3/1.0	2d	
SIS	>30	07 ^h	08 ^h /16 ^h	1.5/0.3	2d	
SOHO						
EPHIN (INT)	>50	07 ^h	08 ^h /17 ^h	0.16/0.08		

Differential fluxes of protons for the event of 1998 November 22

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	08 ^h	12 ^h /18 ^h	4/6	~2d	
CPME	2-4.6	07 ^h	12 ^h /18 ^h	1.4/1.3	~2d	
CPME	4.6-15	07 ^h	10 ^h /18 ^h	0.2/0.1	~2d	
CPME	15-25	07 ^h	10 ^h /18 ^h	0.08/0.025	~2d	
CPME	25-48	07 ^h	09 ^h /19 ^h	0.021/0.009	~2d	
CPME	48-96	07 ^h	09 ^h / -	0.006/ -	~1d	
CPME	96-145	06 ^h	09 ^h / -	0.004/ -	~1d	
CPME	145-440					
SOHO						
LION	0,75-2	07 ^h	10 ^h /19 ^h	6.5/27	~1d	
LION	2-6	07 ^h	11 ^h /19 ^h	1.2/1.8	~1d	
EPHIN	4-8	08 ^h	10 ^h /17 ^h	0.75/0.5	~1d	
EPHIN	8-25	07 ^h	09 ^h /17 ^h	0.13/0.06	~1d	
EPHIN	25-41	07 ^h	08 ^h /17 ^h	0.02/0.005	~1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1998 November 22

1998 November 22 • AR8384 To event 352

H α		0635	0645	0711	S27W82*	1N	
1 -12.5	keV	0630	0642	0649		X3.7	2.0E-1
53 – 93	keV	<063720	064014	081534		6689	HXT Y
>16	MeV		0639				OSSE C
>300	MeV	063336	063917	064837			BATS C

5.7	GHz	0630.8	~0640.0	0742.0	[0.5 / 5.7]	~2.78	5.7
2.7	GHz	0632.0	0636.0	0649.0		2.56	2.7
500	MHz	0636.2	0640.6	0646.2		1.48	
DS II	30-420	0638		0706	SH	3	
DS II	25-170	0639		~0650	FN	3	
CME	WL						gap

* – PRF1212



Particle event: To($E_p > 10$ MeV) – 24d03^h

Tmax($E_p > 10$ MeV) – 24d10^h, Jmax ($E_p > 10$ MeV) – 1.25 /cm².s.sr *)

Duration of the event – 2 days *)

Quasimaximal energy of protons in the event – $E_{qm} = 210$ MeV


*) According to data IMP-8 (CPME)

Sources: • solar flare 24d02^h07^m, X1.0/SF, S30W81, AR8384



Main x-ray burst 1-8 Å: onset – 24d02^h07^m, max – 24d02^h20^m, $\Phi = 0.12$ J/m²

CME: 24d02^h30^m, $V = 1798$ km/s, $\Delta\phi = 360^\circ$, $dA = 225^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1998 November 24



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1998 November 24

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	03 ^h	14 ^h	0.95	7d	
EPS	>10	03 ^h	14 ^h	0.5	6d	
EPS	>30	03 ^h	10 ^h	0.27	4d	
EPS	>50	03 ^h	10 ^h	0.12	4d	
EPS	>60	03 ^h	10 ^h	0.09	3d	
EPS	>100	03 ^h	10 ^h	0.04	2d	
IMP-8						
CPME	>1	03 ^h	10 ^h	2.3	2d	
CPME	>4	03 ^h	10 ^h	1.8	2d	
CPME	>10	03 ^h	10 ^h	1.25	2d	
CPME	>30	03 ^h	10 ^h	0.7	2d	
CPME	>60	03 ^h	10 ^h	0.5	2d	
ACE						
SIS	>10	03 ^h	09 ^h	1.5	2d	
SIS	>30	03 ^h	09 ^h	0.6	2d	
SOHO						
EPHIN (INT)	>50	03 ^h	09 ^h	0.11	1d	

Differential fluxes of protons for the event of 1998 November 24

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	05 ^h	10 ^h	1.2	-	
CPME	2-4.6	05 ^h	10 ^h	0.5	-	
CPME	4.6-15	05 ^h	10 ^h	0.1	-	
CPME	15-25	05 ^h	10 ^h	0.04	2d	
CPME	25-48	05 ^h	10 ^h	0.01	2d	
CPME	48-96	05 ^h	10 ^h	0.0035	2d	
CPME	96-145	05 ^h	10 ^h	0.002	1d	
CPME	145-440	05 ^h	10 ^h	0.001	1d	

SOHO						
LION	0,75-2	02 ^h	10 ^h	0.6	6d	
LION	2-6	02 ^h	10 ^h	0.12	6d	
EPHIN	4-8	03 ^h	09 ^h	0.31	5d	
EPHIN	8-25	03 ^h	09 ^h	0.07	4d	
EPHIN	25-41	03 ^h	09 ^h	0.015	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
1998 November 24**


1998 November 24

•

AR8384

To event 353

H α		0256	0256	0259	S30 W81	SF	
1 -12.5	keV	0207	0220	0237		X1.0	1.2E-1
53 – 93	keV	<0217:19	~0217:19	0407:25		256	HXT Y
>300	keV	0206:04	0214:31	0244:37		0.00	BATS C
15.4	GHz	0209.0	0215.0	0221.0	U0.4-0.6/8.8-15.4	3.18	15.4
8.8	GHz	0208.0	0215.0	0225.0		3.15	8.8
5	GHz	0207.0	0214.0	0225.0		2.78	
2.7	GHz	0208.0	0214.0	0222.0		2.67	
1.4	GHz	0210.0	0214.0	0221.0		2.53	
610	MHz	0210.0	0212.0	0223.0		2.43	
410	MHz	0212.0	0221.0	0222.0		2.45	
245	MHz	0215.0	0215.0	0223.0		2.68	
DS II	60-150	0217		0222	SH,H	2	
DS II	35-60	0218		0220	FN,H	2	
DS III	18-270	0215		0216	G	3	
CME	WL	0230	1798 km/s	-12.5 km/s ²	360°	225°	



Events in 1999

			Page
1.	Event 1999.01.20 – (1999-020)	№ 354	121
2.	Event 1999.01.22 – (1999-022)	№ 355	125
3.	Event 1999.04.24 – (1999-114)	№ 356	129
4.	Event 1999.05.04 – (1999-124)	№ 357	133
5.	Event 1999.05.09 – (1999-129)	№ 358	138
6.	Event 1999.05.27 – (1999-147)	№ 359	142
7.	Event 1999.06.01 – (1999-152)	№ 360	147
8.	Event 1999.06.04 – (1999-155)	№ 361	151
9.	Event 1999.06.11 ? (1999-162)	№ 362	156
10.	Event 1999.06.25 – (1999-176)	№ 363	160
11.	Event 1999.11.17 – (1999-321)	№ 364	165

Particle event: To($E_p > 10$ MeV) – 20d23^h

$T_{\max 1}(E_p > 10 \text{ MeV})$ – 21d11^h, $J_{\max 1}(E_p > 10 \text{ MeV})$ – 1.3 /cm².s.sr

$T_{\max 2}(E_p > 10 \text{ MeV})$ – 22d06^h, $J_{\max 2}(E_p > 10 \text{ MeV})$ – 1 /cm².s.sr

Duration of the event – 1 day

Quasimaximal energy of protons in the event – $E_{qm1} = 270$ MeV

– $E_{qm2} = 250$ MeV


Sources: ☐ solar flare 20d19^h06^m, M5.2/..., n27e90*, AR – unknown

Main X-ray burst 1-8 Å: onset – 20d19^h06^m, max – 20d20^h04^m, $\Phi = 0.25$ J/m²


CME: gap

* – probable localization of the flare event

Particle fluxes and associated phenomena




Temporary profiles of the proton fluxes for the event of 1999 January 20



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 January 20

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	-	21d11 ^h /22d06 ^h	3.6/1.5	1d	
EPS	>10	23 ^h	21d11 ^h /22d06 ^h	1.3/1	1d	
EPS	>30	23 ^h	21d11 ^h /22d06 ^h	0.4/0.4	1d	
EPS	>50	23 ^h	21d11 ^h /22d06 ^h	0.2/0.15	1d	
EPS	>60	23 ^h	21d11 ^h /22d06 ^h	0.14/0.13	1d	
EPS	>100	23 ^h	21d11 ^h /22d06 ^h	0.07/0.05	1d	
METEOR-2						
CBM	>5	22 ^h	21d17 ^h -	11,6/ -	1.5d	
CBM	>15	22 ^h	21d15 ^h -	2,15/ -	1.5d	
CBM	>25	22 ^h	21d14 ^h -	0,35/ -	1.5d	
CBM	>40	22 ^h	21d13 ^h -	0,11/ -	1.5d	
POES-15						
MEPED	>0.24	-	21d18 ^h -	460/ -	-	
MEPED	>0.8	-	21d18 ^h -	270/ -	-	
MEPED	>2.5	-	21d18 ^h -	82/ -	-	
MEPED	>6.9	-	21d18 ^h -	13/ -	-	
MEPED	>16	-	21d18 ^h -	3.16/ -	-	
MEPED	>36	-	21d18 ^h -	1.3/ -	-	
MEPED	>70	-	21d18 ^h -	0.29/ -	-	
MEPED	>140	-	21d18 ^h -	0.24/ -	-	
IMP-8						
CPME	>1	17 ^h	21d15 ^h /22d06 ^h	110/205	1d	
CPME	>4	<21d04 ^h	21d12 ^h /22d06 ^h	12.4/19.6	1d	
CPME	>10	<21d04 ^h	21d15 ^h /22d06 ^h	3/4.7	1d	
CPME	>30	<21d04 ^h	21d15 ^h /22d06 ^h	0.75/0.6	1d	
CPME	>60	<21d04 ^h	21d15 ^h /22d06 ^h	0.45/0.4	1d	
ACE						
SIS	>10	23 ^h	21d14 ^h /22d06 ^h	4.3/4.4	1d	
SIS	>30	23 ^h	21d14 ^h /22d06 ^h	1.4/ 2,1	1d	

Differential fluxes of protons for the event of 1999 January 20

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	07 ^h	21d15 ^h /22d04 ^h	72/124	1d	
CPME	2-4.6	08 ^h	21d15 ^h /22d04 ^h	13/23	1d	
CPME	4.6-15	<21d04 ^h	21d12 ^h /22d04 ^h	0.7/1.2	1d	
CPME	15-25	<21d04 ^h	21d12 ^h /22d04 ^h	0.07/0.12	1d	
CPME	25-48	<21d04 ^h	21d18 ^h /22d04 ^h	0.0180/02	1d	
CPME	48-96	<21d04 ^h	21d15 ^h /22d04 ^h	0.005/0.004	1d	
CPME	96-145	<21d04 ^h	21d18 ^h /22d03 ^h	0.0025/0.004	1d	
CPME	145-440	<21d04 ^h	21d12 ^h /22d03 ^h	0.001/0.0006	1d	
SOHO						
		No	Data			

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
1999 January 20**

1999 January 20




ARXXXX

To event 354

H α	6563 Å	No Flare Patrol			n27e90*		
1 -12.5	keV	1906	2004	2100		M5.2	2.5E-1
33 – 53	keV	190511	191931	>192325		389	HXT Y
8.8	GHz	1911.0	1912.0	1912.0		1.52	
5	GHz	1911.0	1920.0	2130.0		2.11	
2.7	GHz	1909.0	1920.0	2029.0	P2.7	2.36	
1.4	GHz	1905.0	1919.0	2008.0		2.20	
235	MHz	1854.6	1919.0	1940.0		0.60	
DS II	25-55	1914		1923		1	
DS IV	25-45	1926		1944		2	
DS V	30-55	1926		1935		1	
CME	WL						gap

* – probable localization of the flare event



Particle event: To($E_p > 10$ MeV) – 22d02^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 22d16^h$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 3 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 23d14^h$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 5 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 4 days

Quasimaximal energy of protons in the event – $E_{qm1} = 85 \text{ MeV}$

– $E_{qm2} = 85 \text{ MeV}$


Sources: source unknown

Ø solar flare 22d17^h05^m, M1.4/SF, N19W44, AR8440


Main X-ray burst 1-8 Å: onset – 22d17^h05^m, max – 22d17^h24^m, $\Phi = 0.015 \text{ J/m}^2$

CME: gap

Particle fluxes and associated phenomena




Temporary profiles of the proton fluxes for the event of 1999 January 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 January 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	03 ^h	16 ^h /23d14 ^h	7/30	4d	
EPS	>10	03 ^h	16 ^h /23d14 ^h	3/5	4d	
EPS	>30	03 ^h	16 ^h /23d14 ^h	0.7/0.3	4d	
EPS	>50	-	16 ^h /23d12 ^h	0.45/0.05	4d	
EPS	>60	-	16 ^h /23d12 ^h	0.26/0.025	-	
EPS	>100	-	- /23d12 ^h	- /0.004	-	
POES-15						
MEPED	>0.24	03 ^h	23d12 ^h	2820	4d	
MEPED	>0.8	03 ^h	23d12 ^h	1220	4d	
MEPED	>2.5	03 ^h	23d12 ^h	290	4d	
MEPED	>6.9	03 ^h	23d12 ^h	40	4d	
MEPED	>16	03 ^h	23d12 ^h	3.43	4d	
MEPED	>36	03 ^h	23d12 ^h	0.67	4d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	02 ^h	23 ^h /23d12 ^h	270/1030	6d	
CPME	>4	02 ^h	23 ^h /23d12 ^h	33/92	4d	
CPME	>10	02 ^h	23 ^h /23d12 ^h	6.1/9.5	4d	
CPME	>30	02 ^h	23 ^h /23d12 ^h	0.5/ -	-	
CPME	>60	-	- / -	- / -	-	
ACE						
SIS	>10	02 ^h	23 ^h /23d11 ^h	5.7/9	3d	
SIS	>30	-	- / -	- / -	-	

Differential fluxes of protons for the event of 1999 January 22

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion)	Comments
IMP-8						
CPME	1-2	02 ^h	22d23 ^h /23d12 ^h	160/610	10d	
CPME	2-4.6	02 ^h	22d23 ^h /23d12 ^h	32.5/120	9d	
CPME	4.6-15	02 ^h	22d23 ^h /23d12 ^h	2.15/6.35	8d	
CPME	15-25	02 ^h	22d23 ^h /23d12 ^h	0.2/0.23	6d	
CPME	25-48	02 ^h	22d23 ^h /23d12 ^h	0.02/0.913	4d	
CPME	48-96	02 ^h	22d23 ^h /23d12 ^h	0.003//0.002	2d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO		No	Data			

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
1999 January 22**



1999 January 22

Ø

AR8440

To event 355

H α	6563 Å	1708	1708	1715	N20W44	SF	
1 -12.5	keV	1705	1724	1736		M1.4	1.5E-2
22 – 32	keV	170809	170809	170821		36	HXT Y
15.4	GHz	1707.0	1708.0	1709.0		1.57	
9.5	GHz	1706.9	1707.8	1709.3		1.38	
8.8	GHz	1707.0	1707.0	1709.0		1.56	
6.7	GHz	1706.9	1707.8	1709.4		1.30	
15.4	GHz	1717.0	1717.0	1722.0		1.76	
9.5	GHz	1717.0	1717.9	1721.0		1.84	
6.7	GHz	1716.7	1717.8	1723.3		1.93	
5	GHz	1716.0	1717.0	1722.0	P5	2.00	
2.7	GHz	1716.0	1717.0	1722.0		1.85	
1.4	GHz	1717.0	1717.0	1720.0		1.48	
610	MHz	1719.0	1719.0	1722.0		1.08	
CME	WL						gap



Particle event: To(Ep>10 MeV) – 24d15^h

Tmax₁(Ep>10 MeV) – 24d21^h, Jmax₁(Ep>10 MeV) – 3.7 /cm².s.sr

Tmax₂(Ep>10 MeV) – 25d06^h, Jmax₁(Ep>10 MeV) – 4.3 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 85 MeV


– Eqm₂ = 210 MeV

Sources: ☐ solar flare event 24d<13^h31^m, n22w90*, AR8517 ~ 3 days behind W limb



CME: 24d13^h31^m, V = 1495 km/s, Δφ = 360°, dA = 321°

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1999 April 24



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 April 24

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	15 ^h	21 ^h /25d06 ^h	5.1/7.0	2d	
EPS	>10	15 ^h	21 ^h /25d06 ^h	3.7/4.3	2d	
EPS	>30	15 ^h	21 ^h /25d05 ^h	0.77/0.5	2d	
EPS	>50	15 ^h	21 ^h /25d05 ^h	0.19/0.12	1d	
EPS	>60	15 ^h	22 ^h /25d04 ^h	0.08/0.06	1d	
EPS	>100	15 ^h	-/25d00 ^h	-/0.03	1d	
POES-15						
MEPED	>0.24	-	-/25d01 ^h	-/582	-	
MEPED	>0.8	-	-/25d01 ^h	-/480	-	
MEPED	>2.5	-	-/25d01 ^h	-/296	-	
MEPED	>6.9	-	-/25d01 ^h	-/70	-	
MEPED	>16	-	-/25d01 ^h	-/7.83	-	
MEPED	>36	-	-/25d01 ^h	-/1.36	-	
MEPED	>70	-	-/25d01 ^h	-/0.25	-	
MEPED	>140	-	-/25d01 ^h	-/0.16	-	
IMP-8						
CPME	>1	15 ^h	23 ^h /25d04 ^h	250/410	2.5d	
CPME	>4	15 ^h	23 ^h /25d05 ^h	75/85	2.5d	
CPME	>10	15 ^h	22 ^h /25d05 ^h	16.7/13.6	2.5d	
CPME	>30	15 ^h	<22 ^h /25d05 ^h	1.2/0.6	2.5d	
CPME	>60	16 ^h	<22 ^h /25d05 ^h	0.4/0.2	2.5d	

ACE						
SIS	>10	15 ^h	21 ^h /25d02 ^h	13.3/16	1.5d	
SIS	>30	15 ^h	21 ^h /25d02 ^h	1.1/0.9	1d	
SOHO						
EPHIN (INT)	>50	14 ^h	17 ^h /23 ^h	0.3/0.33	~0.5d	

Differential fluxes of protons for the event of 1999 April 24

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	-	- /25d04 ^h	- /179	~2,8d	
CPME	2-4.6	-	- /25d04 ^h	- /64	~2,8d	
CPME	4.6-15	-	- /25d04 ^h	- /6,1	~2,8d	
CPME	15-25	-	- /25d04 ^h	- /0,5	~2,5d	
CPME	25-48	-	- /25d04 ^h	- /0,03	~2d	
CPME	48-96	-	- /25d04 ^h	- /0,01	~2d	
CPME	96-145	-	- / -	- / -	-	
CPME	145-440	-	- / -	- / -	-	
SOHO						
LION	0,75-2	16 ^h	22 ^h /25d02 ^h	17/37	~2d	
LION	2-6	16 ^h	22 ^h /25d02 ^h	5,1/9,3	~2d	
EPHIN	4-8	14 ^h	22 ^h /25d02 ^h	15.5/22	~2.5d	
EPHIN	8-25	14 ^h	21 ^h /25d02 ^h	1.5/1.9	~2.5d	
EPHIN	25-41	14 ^h	21 ^h /25d02 ^h	0.05/0.04	~2.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1999 April 24

1999 April 24 ☐ AR8517 To event 356

CME	WL	1331	1495 km/s	37.1 km/s	360°	321°	
H α		No Flare			n22e90*		
1 - 12.5	keV	No	X-ray	Burst			

* – probable localization of the flare event

Particle event: To($E_p > 10$ MeV) – 04d08^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 05\text{d}21^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 3.7 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 06\text{d}06^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 4 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 4 days

Quasimaximal energy of protons in the event – $E_{qm1} = 75 \text{ MeV}$

– $E_{qm2} = 65 \text{ MeV}$


Sources: ☉ solar flare 03d05^h36^m, M4.4/2N, N15E32, AR8525

Main X-ray burst 1-8 Å: onset – 03d 05^h36^m, max – 03d 06^h 02^m, $\Phi = 0.099 \text{ J/m}^2$



CME: 03d06^h06^m, $V = 1584 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 088^\circ$

▲ SC 05d15^h 43^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1999 May 04



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 May 04

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	08 ^h	05d21 ^h /06d09 ^h	17/36	4d	
EPS	>10	08 ^h	05d19 ^h /06d06 ^h	3.7/4	4d	
EPS	>30	07 ^h	05d18 ^h /06d04 ^h	0.13 /0.16	3d	
EPS	>50	-	- /06d04 ^h	- /0.02	1d	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-15						
MEPED	>0.24	-	05d16 ^h /06d03 ^h	1120/4280	4d	
MEPED	>0.8	-	05d16 ^h /06d03 ^h	430/1940	4d	
MEPED	>2.5	-	05d16 ^h /06d03 ^h	190/570	4d	
MEPED	>6.9	-	05d16 ^h /06d03 ^h	25/48	4d	
MEPED	>16	-	05d16 ^h /06d03 ^h	0.9/1.6	3d	
MEPED	>36	-	05d16 ^h /06d03 ^h	0.08/0.3	1d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	02 ^h	05d19 ^h / -	660/ -	7 d	
CPME	>4	02 ^h	05d19 ^h / -	80/ -	7 d	
CPME	>10	02 ^h	05d19 ^h / -	6.8/ -	6 d	
CPME	>30	01 ^h	05d19 ^h / -	0.13/ -	-	
CPME	>60	-	-	- / -	-	
ACE						
SIS	>10	03d18 ^h	05d19 ^h /06d02 ^h	6.6/6.8	6d	
SIS	>30	03d18 ^h	05d15 ^h / -	0.2/ -	3d	

SOHO						
EPHIN (INT)	>50	03d18	- “ -	- “ -	4d	


Differential fluxes of protons for the event of 1999 May 04

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	03d12 ^h	05d15 ^h -	348/ -	7d	
CPME	2-4.6	03d16 ^h	05d15 ^h -	92/ -	7d	
CPME	4.6-15	03d20 ^h	05d19 ^h -	6,3/ -	5d	
CPME	15-25	01 ^h	05d19 ^h -	0,18/ -	~4d	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	03d22 ^h	05d18 ^h /06d04 ^h	42/106	6d	
LION	2-6	02 ^h	05d18 ^h /06d03 ^h	7.3/18	6d	
EPHIN	4-8	03d14 ^h	05d18 ^h /06d02 ^h	12/19	6d	
EPHIN	8-25	03d14 ^h	05d18 ^h /06d02 ^h	0.7/0.72	6d	
EPHIN	25-41	03d14 ^h	05d18 ^h /06d02 ^h	0.0035/0.0018	5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1999 May 04

1999	May 03	\odot	AR8525	To event 357			
H α	6563Å	0543	0551	0745	N15 E32	2N	U
1 -12.5	keV	0536	0602	0632		M4.4	9.9E-2
23 – 33	keV	0602		0605		290	HXT Y

15.4	GHz	0548.0	0608.0	0651.0		1.86	
8.8	GHz	0544.0	0607.0	0708.0		1.96	
5	GHz	0542.0	0607.0	0655.0		2.11	
2.7	GHz	0540.0	0548.0	0630.0		2.34	
1.4	GHz	0540.0	0548.0	0634.0		2.34	
610	MHz	0543.0	0551.0	0621.0		1.92	
500	MHz	0540.0	0552.0	0630.0		2.32	
410	MHz	0544.0	0551.0	0623.0		2.64	
245	MHz	0547.0	0613.0	0621.0	U0.6 P1.4-2.6	2.54	
DS II	35-85	0520		0606		3	
DS II	25-180	0543		0612	SH,H	2	
DS IV	40-800	<0551		~0710		2	
DS IV	35-85	0606		0709		3	
DS III	40-90	<0551		~1010	N	1	
DS DCIM	800-2000	0542		0641	GG	2	
CME	WL	0606	1584 km/s	15.8 km/s ²	360°	088°	



Particle event: To($E_p > 10$ MeV) – 09d19^h

Tmax($E_p > 10$ MeV) – 09d21^h, Jmax ($E_p > 10$ MeV) – $1.2 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}^*$

Duration of the event – 1 day

Quasimaximal energy of protons in the event = 75 MeV

*) The data from IMP-8


Sources: ■ solar flare 09d17^h53^m, M7.6/ ..., n23w90*, AR8526 one day behind W limb;

Main X-ray burst 1-8 Å: onset – 09d17^h53^m, max – 09d18^h07^m, $\Phi = 0.061 \text{ J/m}^2$



CME: 09d18^h27^m, $V = 0615 \text{ km/s}$, $\Delta\phi = 172^\circ$, $dA = 316^\circ$

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1999 May 09



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 May 09

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	-	22 ^h	0.2	1d	
EPS	>10	-	22 ^h	0.1	1d	
EPS	>30	-	21 ^h	0.05	0.5d	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-15						
MEPED	>0.24	-	23 ^h	96	-	
MEPED	>0.8	-	23 ^h	56	-	
MEPED	>2.5	-	23 ^h	32	-	
MEPED	>6.9	-	23 ^h	8	-	
MEPED	>16	-	23 ^h	0.99	-	
MEPED	>36	-	23 ^h	0.61	-	
MEPED	>70	-	23 ^h	0.2	-	
MEPED	>140	-	23 ^h	0.22	-	
IMP-8						
CPME	>1	21 ^h	23 ^h	27.4	4d	
CPME	> 4	19 ^h	22 ^h	4.8	1d	
CPME	>10	19 ^h	21 ^h	1.2	1d	
CPME	>30	19 ^h	20 ^h	0.3	1d	
CPME	>60	19 ^h	20 ^h	0.1	1d	
ACE						
SIS	>10	18 ^h	21 ^h	2	1d	
SIS	>30	18 ^h	22 ^h	0.4	1d	
SOHO						
EPHIN (INT)	>50	18 ^h	20 ^h	0.03	0.5d	

Differential fluxes of protons for the event of 1999 May 09

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	21 ^h	23 ^h	15.2	3d	
CPME	2-4.6	21 ^h	23 ^h	3.7	2d	
CPME	4.6-15	20 ^h	22 ^h	0,33	2d	
CPME	15-25	<20 ^h	22 ^h	0,04	1d	
CPME	25-48	<20 ^h	21 ^h	0,009	1d	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	22 ^h	10d02 ^h	2.2	3d	
LION	2-6	21 ^h	23 ^h	0.387	3d	
EPHIN	4-8	20 ^h	22 ^h	1.05	5d	
EPHIN	8-25	19 ^h	21 ^h	0.13	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1999 May 09

1999 May 09




AR8526

To event 358

H α	6563Å	No Flare			n23 w90*		
1 -12.5	keV	1753	1807	1815		M7.6	6.1E-2
23 – 33	keV	1804		1821		1740	HXT Y
>300	keV	175542	175751	180725		2204.41	BATS C
>300	keV					2.1+/-0.1	OSSE C

15.4	GHz	1757.0	1757.0	1802.0		2.84	
8.8	GHz	1756.0	1757.0	1802.0		2.69	
5	GHz	1756.0	1758.0	1811.0	U1.4 / 15.4	2.41	
2.7	GHz	1757.0	1759.0	1801.0		2.20	
1.4	GHz	1757.0	1758.0	1800.0		1.88	
410	MHz	1804.0	1805.0	1805.0		2.72	
235	MHz	1800.3	1802.0	1803.0		3.19	
DS III	25-75	1803		1803		1	
CME	WL	1827	0615 km/s	-3.0 km/s ²	172°	316°	

* – probable localization of the flare event



Particle event: To($E_p > 10$ MeV) – 27d12^h

Tmax($E_p > 10$ MeV) – 27d13^h, Jmax ($E_p > 10$ MeV) – 2.75 /cm².s.sr

Duration of the event – 1.5 days


Quasimaximal energy of protons in the event – $E_{qm} = 275$ MeV

Sources: ☐ back side solar flare event < 27d11^h06^m, AR unknown, behind W_L



○ solar flare 26d19^h15^m, M1.2/2N, N17E46, AR8552

CME: 27d11^h06^m, $V = 1691$ km/s, $\Delta\phi = 360^\circ$, $dA = 341^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1999 May 27



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 May 27

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	12 ^h	13 ^h	3.44	1.5d	
EPS	>10	12 ^h	13 ^h	2.75	1.5d	
EPS	>30	12 ^h	13 ^h	0.73	1.5d	
EPS	>50	12 ^h	12 ^h	0.33	1d	
EPS	>60	12 ^h	12 ^h	0.24	1d	
EPS	>100	-	12 ^h	0.09	1d	
IMP-8						
CPME	>1	12 ^h	15 ^h	25.6	2d	
CPME	>4	12 ^h	15 ^h	11	2d	
CPME	>10	12 ^h	14 ^h	3.9	1.5d	
CPME	>30	11 ^h	13 ^h	0.9	1d	
CPME	>60	11 ^h	12 ^h	0.5	1d	
ACE						
SIS	>10	11 ^h	14 ^h	5.7	1d	
SIS	>30	11 ^h	13 ^h	1.5	1d	
SOHO						
EPHIN (INT)	>50	11 ^h	13 ^h	0.41	1d	

Differential fluxes of protons for the event of 1999 May 27

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	14 ^h	19 ^h	14.8	1.5d	
CPME	2-4.6	13 ^h	19 ^h	5.7	1.5d	
CPME	4.6-15	13 ^h	15 ^h	0.7	1.5d	
CPME	15-25	13 ^h	14 ^h	0.17	1.5d	
CPME	25-48	13 ^h	13 ^h	0.025	1d	
CPME	48-96	12 ^h	12 ^h	0.006	1d	
CPME	96-145	12 ^h	12 ^h	-	1d	
CPME	145-440	12 ^h	12 ^h	0.001	1d	
SOHO						
LION	0,75-2	16 ^h	14 ^h	2.1	1.5d	
LION	2-6	16 ^h	15 ^h	0.47	1.5d	
EPHIN	4-8	11 ^h	14 ^h	2.1	1.5d	
EPHIN	8-25	11 ^h	14 ^h	0.47	1.5d	
EPHIN	25-41	11 ^h	14 ^h	0.04	1.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References

Slocum P.L., E.C. Stone, R.A. Leske et al., 2003.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1999 May 27

1999		May 27	☐	AR8552	To event 359	
CME	WL	1106	1691 km/s	-33.5 km/s	360°	341°

1999		May 26	○	AR8552	To event 359		
H α		1917	1921	2026	N17 E46	2N	FU
1 -12.5	keV	1915	1932	1945		M1.2	1.3E-02
6.7	GHz	1911.0	1934.0	2106.0		1.43	
DS III	25-200	1955		1955	B	2	
DS CONT	25-75	1954		2038		1	
CME	WL	2026	0396 km/s	4.4km/s ²	017°	039°	

Particle event: To($E_p > 10$ MeV) – 01d20^h

Tmax₁($E_p > 10$ MeV) – 02d09^h, Jmax₁($E_p > 10$ MeV) – 23 /cm².s.sr

Tmax₂($E_p > 10$ MeV) – 02d21^h, Jmax₂($E_p > 10$ MeV) – 13 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 350 MeV

– Eqm₂ = 240 MeV


Sources: □ back side solar flare 01d18^h53^m, C1.2; n25w90, AR unknown

Main X-ray burst 1-8 Å: onset – 01d18^h53^m, max – 01d19^h04^m, $\Phi = 0.0027$ J/m²



CME: 01d19^h37^m, V = 1772 km/s, $\Delta\phi = 360^\circ$; dA=359°

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1999 June 01



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 June 01

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	20 ^h	02d10 ^h /02d21 ^h	47/20.1	2d	
EPS	>10	20 ^h	02d09 ^h /02d21 ^h	23/13	2d	
EPS	>30	20 ^h	02d09 ^h /02d21 ^h	4.4/1.7	2d	
EPS	>50	20 ^h	02d09 ^h /02d21 ^h	1.42/0.6	2d	
EPS	>60	20 ^h	02d05 ^h /02d20 ^h	0.90/0.24	2d	
EPS	>100	20 ^h	02d03 ^h /02d20 ^h	0.43/0.07	2d	
POES-15						
MEPED	>0.24	-	02d08 ^h / -	755/ -	2d	
MEPED	>0.8	-	02d08 ^h / -	403/ -	2d	
MEPED	>2.5	-	02d08 ^h / -	192/ -	2d	
MEPED	>6.9	-	02d08 ^h / -	106/ -	2d	
MEPED	>16	-	02d08 ^h / -	16/ -	2d	
MEPED	>36	-	02d08 ^h / -	4.65/ -	2d	
MEPED	>70	-	02d08 ^h / -	0.93/ -	2d	
MEPED	>140	-	02d08 ^h / -	0.26/ -	2d	
IMP-8						
CPME	>1	21 ^h	02d09 ^h /02d19 ^h	215/188	2 d	
CPME	>4	21 ^h	02d09 ^h /02d19 ^h	68/65,4	2 d	
CPME	>10	21 ^h	02d09 ^h /02d19 ^h	26/23	2 d	
CPME	>30	21 ^h	02d08 ^h /02d19 ^h	3.9/2.3	2 d	
CPME	>60	21 ^h	02d07 ^h /02d19 ^h	1.5/0.7	2 d	
ACE						
SIS	>10	21 ^h	02d08 ^h /02d18 ^h	29/28.3	2 d	
SIS	>30	21 ^h	02d07 ^h /02d18 ^h	6.5/3.2	2 d	

SOHO						
EPHIN (INT)	>50	20 ^h	02d07 ^h /02d19 ^h	0.85/0.47	-	

Differential fluxes of protons for the event of 1999 June 01

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	02d01 ^h	02d09 ^h /02d19 ^h	80.2/66	2d	
CPME	2-4.6	22 ^h	02d09 ^h /02d19 ^h	30/25.6	2d	
CPME	4.6-15	21 ^h	02d09 ^h /02d19 ^h	4.0/4.1	2d	
CPME	15-25	21 ^h	02d09 ^h /02d19 ^h	1.2/1.05	2d	
CPME	25-48	21 ^h	02d08 ^h /02d19 ^h	0.14/0.095	2d	
CPME	48-96	20 ^h	02d08 ^h /02d19 ^h	0.020/0.01	2d	
CPME	96-145	- / -	- / -	- / -	-	
CPME	145-440	- / -	- / -	- / -	-	
SOHO						
LION	0,75-2	21 ^h	02d07 ^h /02d19 ^h	46.6/21.6	2d	
LION	2-6	21 ^h	02d07 ^h /02d19 ^h	9.6/4.9	2d	
EPHIN	4-8	21 ^h	02d07 ^h /02d19 ^h	20.7/12.1	2d	
EPHIN	8-25	21 ^h	02d07 ^h /02d19 ^h	3.3/1.9	2d	
EPHIN	25-41	20 ^h	02d07 ^h /02d19 ^h	0.17/0.1	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
1999 June 01**

1999 June 01 □ AR To event 360

Hα	6563 Å	No Flare			n25w90*		
1 -12.5	keV	1853	1904	1932		C1.2	2.7E-3
5	GHz	1855.0	1855.0	1856.0		1.75	
2.8	GHz	1852.0	1855.0	1931.0		1.26	2.8
CME	WL	1938	1772 km/s	1.8 km/s ²	360°	359°	

* – probable localization of the flare event

Particle event: To($E_p > 10$ MeV) – 04d08^h

Tmax($E_p > 10$ MeV) – 04d12^h, Jmax ($E_p > 10$ MeV) – 20 /cm².s.sr

Duration of the event – 3 days


Quasimaximal energy of protons in the event – $E_{qm} = 300$ MeV

Sources: • solar flare 04d06^h52^m, M3.9/2N, N18W72, AR8552

Main X-ray burst 1-8 Å: onset – 04d06^h52^m, max – 04d07^h03^m, $\Phi = 0.024$ J/m²



CME: 04d07^h27^m, $V = 2230$ km/s, $\Delta\phi = 150^\circ$, $dA = 287^\circ$

Particle fluxes and associated phenomena




May - June 1999

Time profiles of the proton fluxes for the event of 1999 June 04



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 June 04

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	08 ^h	13 ^h	48	3d	
EPS	>10	08 ^h	12 ^h	20	3d	
EPS	>30	08 ^h	12 ^h	2.2	3d	
EPS	>50	08 ^h	11 ^h	0.55	3d	
EPS	>60	-	11 ^h	0.3	2d	
EPS	>100	-	11 ^h	0.07	2d	
POES-15						
MEPED	>0.24	-	12 ^h	615	3d	
MEPED	>0.8	-	12 ^h	540	3d	
MEPED	>2.5	-	12 ^h	430	3d	
MEPED	>6.9	-	12 ^h	170	3d	
MEPED	>16	-	12 ^h	19.72	3d	
MEPED	>36	-	12 ^h	3.27	3d	
MEPED	>70	-	12 ^h	0.46	2d	
MEPED	>140	-	12 ^h	0.26	2d	
IMP-8						
CPME	>1	09 ^h	20 ^h	1020	5 d	
CPME	>4	09 ^h	17 ^h	220	5 d	
CPME	>10	09 ^h	16 ^h	37	5 d	
CPME	>30	09 ^h	13 ^h	2.9	2 d	
CPME	>60	09 ^h	12 ^h	1.4	2 d	
ACE						
SIS	>10	09 ^h	16 ^h	39	4d	
SIS	>30	09 ^h	13 ^h	2.8	1d	
SOHO						
EPHIN (INT)	>50	09 ^h	12 ^h	0.41	<1d	



Differential fluxes of protons for the event of 1999 June 04

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	10 ^h	23 ^h	560	5d	
CPME	2-4.6	09 ^h	19 ^h	188	5d	
CPME	4.6-15	09 ^h	16 ^h	17	5d	
CPME	15-25	07 ^h	14 ^h	1.2	5d	
CPME	25-48	09 ^h	13 ^h	0,1	4d	
CPME	48-96	08 ^h	12 ^h	0,01	2d	
CPME	96-145	08 ^h	-	-	1d	
CPME	145-440	06 ^h	10 ^h	0,002	1d	
SOHO						
LION	0,75-2	10 ^h	23 ^h	110	4d	
LION	2-6	10 ^h	17 ^h	28	4d	
EPHIN	4-8	09 ^h	15 ^h	49.7	5d	
EPHIN	8-25	09 ^h	14 ^h	40.5	5d	
EPHIN	25-41	09 ^h	13 ^h	0.1	5d	
EPHIN	41-53	- ' -	- ' -	- ' -	- ' -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1999 June 04

1999 June 04		•		AR8552		To event 361	
H α	6563 Å	0655	0700	0716	N18 W72	2N	F
1 -12.5	keV	0652	0703	0711		M3.9	2.4E-2
>300	keV	065329	065958	071125		4242.30	BATS C
15.4	GHz	0656.0	0700.0	0710.0		3.11	
8.8	GHz	0656.0	0700.0	0713.0	U0.9 P8.8	3.38	
5	GHz	0656.0	0700.0	0710.0		3.18	
2.7	GHz	0653.0	0701.0	0708.0		2.56	
1.4	GHz	0654.0	0659.0	0705.0		2.57	
900	MHz	0649.5	0659.1	0708.6		2.24	
600	MHz	0655.0	0700.7	0711.3		2.81	
410	MHz	0654.0	0702.0	0707.0		2.79	
245	MHz	0653.0	0655.0	0716.0		2.82	
DS II	30-90	0704		0710	FN	2	
DS II	50-150	0705		0716		3	
DS IV	35-85	0658		0728		3	
DS III	45-270	0652		~0721	N	2	
DS III	380-550	0654		0658	GG,RS	2	
DS DCIM	800-2000	0652		0709	GG,FS	2	
DS DCIM	270-550	0653		0700	P	2	
DS DCIM	2000-4365	0653		0710	GG	2	

5.7	GHz	0754.7	0756.9	0809.0		1.04	
2.8	GHz	0751.0	0755.0	0820.0		1.21	
1.4	GHz	0756.0	0757.0	0758.0		1.60	
900	MHz	0753.5	0756.1	0804.9		1.83	
600	MHz	0753.5	0756.2	0800.0		2.66	
410	MHz	0754.0	0756.0	0801.0	P0.4	2.83	
245	MHz	0754.0	0757.0	0758.0		2.11	
DS III	45-90	~0753		>1200	N	1	
DS DCIM	800-2000	0754		0801	GG,FS	2	
DS DCIM	290-550	0754		0759	P	1	
DS DCIM	2000-4355	0755		0800		1	
CME	WL	0727	2230 km/s	-158.8 km/s	150°	287°	



Particle event: To($E_p > 10$ MeV) – 11d01^h

Tmax($E_p > 10$ MeV) – 11d03^h, Jmax ($E_p > 10$ MeV) – 2.2 /cm².s.sr *)

Duration of the event – 10 hours

Quasimaximal energy of protons in the event – $E_{qm} = 240$ MeV


*) From CPME s/c IMP-8

Sources: □ solar flare 11d01^h05^m, C1.0/..., AR behind the W- limb



Main X-ray burst 1-8 Å: onset: – 11d01^h05^m, max – 11d01^h10^m, $\Phi = 0.00061$ J/m²

CME: 11d01^h27^m, $V = 0719$ km/s, $\Delta\phi = 101^\circ$, $dA = 288^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1999 June 11



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 June 11

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	01 ^h	03 ^h	0.5	10 h	
EPS	>10	00 ^h	03 ^h	0.4	10 h	
EPS	>30	00 ^h	02 ^h	0.24	10 h	
EPS	>50	00 ^h	02 ^h	0.16	10 h	
EPS	>60	00 ^h	02 ^h	0.13	8 h	
EPS	>100	00 ^h	01 ^h	0.07	6 h	
POES-15						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	02 ^h	2.21	10 h	
MEPED	>36	-	02 ^h	0.88	10 h	
MEPED	>70	-	02 ^h	0.23	8 h	
MEPED	>140	-	02 ^h	0.2	6 h	
IMP-8						
CPME	>1	01 ^h	06 ^h	9	10 h	
CPME	>4	01 ^h	04 ^h	3.6	10 h	
CPME	>10	01 ^h	03 ^h	2.2	10 h	
CPME	>30	01 ^h	02 ^h	0.6	10 h	
CPME	>60	01 ^h	02 ^h	0.28	10 h	
ACE						
SIS	>10	01 ^h	02 ^h	2.1	10 h	
SIS	>30	01 ^h	02 ^h	0.85	8 h	
SOHO						
EPHIN (INT)	>50	01 ^h	02 ^h	0.1	10 h	

Differential fluxes of protons for the event of 1999 June 11

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	04 ^h	06 ^h	4	15h	
CPME	2-4.6	04 ^h	05 ^h	1.5	15h	
CPME	4.6-15	04 ^h	03 ^h	0.22	15h	
CPME	15-25	03 ^h	02 ^h	0.11	12h	
CPME	25-48	-	02 ^h	0.018	10h	
CPME	48-96	-	01 ^h	0.007	10h	
CPME	96-145	-	02 ^h	0.004	8h	
CPME	145-440	-	01 ^h	0.0003	8h	
SOHO						
LION	0,75-2	01 ^h	06 ^h	1	~1d	
LION	2-6	01 ^h	05 ^h	0,3	~1d	
EPHIN	4-8	01 ^h	03 ^h	0.64	1,5d	
EPHIN	8-25	01 ^h	02 ^h	0.1	1d	
EPHIN	25-41	01 ^h	02 ^h	0.015	1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1999 June 11

1999 June 11 □ AR To event 362

H α	6563 Å	No Flare					
1 -12.5	keV	0105	0110	0116		C1.0	6.1E-04
1.4	GHz	0038.0	0038.0	0039.0		1.61	
245	MHz	0039.0	0039.0	~0039.0		1.74	
DS II	50-250	0039		0049	SH,H	3	
DS II	30-130	0039		0049	FN,H	3	
DS IV	25-75	0051		0116		1	
DS III	20-150	0041		0042	B	2	
410	MHz	0109.0	0109.0	0110.0		2.34	
245	MHz	0109.0	0109.0	0110.0		2.23	
DS III	30-55	0103		0104		1	
CME	WL	0127	0719 km/s	-38.2 km/s	101°	262°	

Particle event: To(Ep>10 MeV) – 25d10^h

Tmax(Ep>10 MeV) – 26d12^h, Jmax(Ep>10 MeV) – 1.7 /cm².s.sr *)

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm = 40 MeV

*) The data from IMP-8

Sources: ○ solar flare 24d12^h04^m, C4.1/1F, N29W13, AR8595


Ø solar flare 26d05^h08^m, M2.3/2B, N24E02, AR8598

Main X-ray burst 1-8 Å: onset – 24d12^h04^m, max – 24d14^h12^m, $\Phi = 0.004 \text{ J/m}^2$



CME: 24d13^h31^m, V = 0975 km/s, $\Delta\phi = 360^\circ$, dA = 335°

▲ SC 26d03^h25^m, 26d20^h16^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1999 June 25



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 June 25

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	20 ^h	26d12 ^h	10.3	1d	
EPS	>10	21 ^h	26d11 ^h	0.3	1d	
EPS	>30	-	26d12 ^h	0.003	-	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-15						
MEPED	>0.24	-	26d10 ^h	9310	-	
MEPED	>0.8	-	26d10 ^h	3670	2d	
MEPED	>2.5	-	26d10 ^h	390	2d	
MEPED	>6.9	-	26d10 ^h	14	2d	
MEPED	>16	-	26d10 ^h	0.5	2d	
MEPED	>36	-	26d10 ^h	0.3	2d	
MEPED	>70	-	26d10 ^h	0.22	2d	
MEPED	>140	-	26d10 ^h	0.2	2d	
IMP-8						
CPME	>1	8 ^h	26d12 ^h	2790	2d	
CPME	>4	10 ^h	26d12 ^h	54	2d	
CPME	>10	10 ^h	26d12 ^h	1.7	2d	
CPME	>30	-	-	-	-	
CPME	>60	-	-	-	-	

ACE						
SIS	>10	8 ^h	26d10 ^h	0.6	2d	
SIS	>30	-	26d12 ^h	0.07	-	
SOHO						
EPHIN (INT)	>50	-	-	-	-	Forbush


Differential fluxes of protons for the event of 1999 June 25

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	09 ^h	26d12 ^h	2340	3d	
CPME	2-4.6	08 ^h	26d12 ^h	226	3d	
CPME	4.6-15	08 ^h	26d12 ^h	3.2	3d	
CPME	15-25	-	-	-	-	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	06 ^h	26d13 ^h	222	2d	
LION	2-6	06 ^h	26d13 ^h	18.5	2d	
EPHIN	4-8	06 ^h	26d12 ^h	7.6	2d	
EPHIN	8-25	06 ^h	26d12 ^h	0.09	2d	
EPHIN	25-41	-	-	-	-	
EPHIN	41-53	- “ -	- “ -	- “ -	-	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1999 June 25

1999	June 24	O	AR8595	To event 363			
H α	6563 Å	<1321	1342	1444	N29W13	1F	SU
1 -12.5	keV	1204	1412	1510		C4.1	4.0E-3
6.7	GHz	1330.0	1420.0	1523.0		1.05	
280	MHz	<1300.0					
235	MHz	<1300.0					
33	MHz	1421.0					
CME	WL	1331	0975 km/s	32.4 km/s	360°	335°	

H α	6563 Å	0505	0512	0538	N24E02	2B	EF
1 -12.5	keV	0508	0512	0514		M2.3	3.9E-3
15.4	GHz	0511.0	0511.0	0512.0	0.6/1.4 - 5/15.4	2.48	
8.8	GHz	0511.0	0511.0	0512.0		2.32	
5.7	GHz	0510.4	0511.9	0521.0		2.11	
5	GHz	0511.0	0511.0	0512.0		1.89	
2.8	GHz	0503.0	0512.0	0514.0		1.85	
1.4	GHz	0511.0	0512.0	0512.0		1.83	
900	MHz	0511.5	0512.3	0512.7		1.72	
600	MHz	0511.3	0512.3	0512.9		1.18	
DS III	450-550	0511		0512		1	
DS DCIM	800-2000	0511		0512	GG	2	
CME	WL	0731	0558 km/s	-9.8 km/s ²	360°	021°	



Particle event: To(Ep>10 MeV) – 17d19^h

Tmax₁(Ep>10 MeV) – 19d02^h, Jmax₁(Ep>10 MeV) – 1 /cm².s.sr *)

Tmax₂(Ep>10 MeV) – 19d23^h, Jmax₂(Ep>10 MeV) – 0.4 /cm².s.sr *)

Duration of the event – 2.5 days

Quasimaximal energy of protons in the event – Eqm₁ = 50 MeV


*) The data from IMP-8 – Eqm₂ = 25 MeV

Sources: • solar flare 17d09^h47^m, M7.4/2B, N17E21, AR8766


Main x-ray burst 1-8 Å: onset – 17d09^h47^m, max – 17d09^h57^m, Φ = 0.04 J/m²

CME: gap

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 1999 November 17



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 1999 November 17

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	19 ^h	19d02 ^h /19d23 ^h	3.4/0.7	2.5d	
EPS	>10	19 ^h	19d02 ^h /19d23 ^h	0.48/0.1	2.5d	
EPS	>30	19 ^h	19d02 ^h / -	0.03/ -	2.5d	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
IMP-8						
CPME	>1	19 ^h	19d02 ^h /19d23 ^h	195/202	5 d	
CPME	>4	19 ^h	19d02 ^h /19d23 ^h	8.6/4	4 d	
CPME	>10	19 ^h	19d02 ^h /19d23 ^h	1/0.4	2.5 d	
CPME	>30	19 ^h	19d02 ^h / -	0.08/ -	-	
CPME	>60	-	19d02 ^h / -	0.01/ -	-	
ACE						
SIS	>10	19 ^h	19d02 ^h / -	0.9/ -	1d	
SIS	>30	-	-	-	-	


Differential fluxes of protons for the event of 1999 November 17

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	23 ^h	19d02 ^h /19d20 ^h	160/160	7d	
CPME	2-4.6	20 ^h	19d02 ^h /19d20 ^h	15/11.3	7d	
CPME	4.6-15	18 ^h	19d01 ^h /19d20 ^h	0.6/0.2	4d	
CPME	15-25	18 ^h	19d00 ^h /19d20 ^h	0.018/ -	3d	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO			No	Data		

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 1999 November 17

1999		November 17		•	AR8766	To event 364	
Hα	6563 Å	0951	0955	>1024	N17E21	2B	E
1 -12.5	keV	0947	0957	1002		M7.4	4.0E-2

15.4	GHz	0952.0	0953.0	1002.0		2.72	
8.8	GHz	0951.0	0953.0	1007.0	U0.9 P1.8	3.00	
5	GHz	0950.0	0954.0	0959.0		2.86	
3	GHz	0951.1	0957.2	1006.6		2.65	
900	MHz	0951.2	0952.9	1006.9		2.05	
610	MHz	0951.0	0958.0	1002.0		2.70	
410	MHz	0951.0	0953.0	0959.0		3.00	
204	MHz	0951.9	0952.5	1006.8		4.63	
DS II	30-80	0959		1005		2	
DS IV	35-85	0951		1403		3	
DS III	30-80	0951		0958		3	
DS DCIM	800-2000	0951		1000	GG	2	
CME	WL						gap



Events in 2000

				Page
1.	Event 2000.02.18 – (2000-049)	№ 365	170
2.	Event 2000.04.04 – (2000-095)	№ 366	175
3.	Event 2000.06.07 – (2000-159)	№ 367	180
4.	Event 2000.06.10 – (2000-162)	№ 368	185
5.	Event 2000.06.17 – (2000-169)	№ 369	190
6.	Event 2000.06.25 – (2000-177)	№ 370	196
7.	Event 2000.07.13 – (2000-195)	№ 371	201
8.	Event 2000.07.14 – (2000-196) – GLE-59	№ 372	206
9.	Event 2000.07.16 – (2000-198)	№ 373	212
10.	Event 2000.07.22 – (2000-204)	№ 374	217
11.	Event 2000.07.28 – (2000-210)	№ 375	222
12.	Event 2000.08.11 – (2000-224)	№ 376	226
13.	Event 2000.08.13 – (2000-226)	№ 377	230
14.	Event 2000.09.12 – (2000-256)	№ 378	235
15.	Event 2000.10.16 – (2000-290)	№ 379	240
16.	Event 2000.10.25 – (2000-299)	№ 380	245
17.	Event 2000.10.31 – (2000-305)	№ 381	249
18.	Event 2000.11.08 – (2000-313)	№ 382	253
19.	Event 2000.11.24 – (2000-329)	№ 383	258
20.	Event 2000.11.26 – (2000-331)	№ 384	264

Particle event: To($E_p > 10$ MeV) – 18d06^h

Tmax($E_p > 10$ MeV) – 18d12^h, Jmax ($E_p > 10$ MeV) – 1.7 /cm².s.sr

Duration of the event – 1 day

Quasimaximal energy of protons in the event – $E_{qm} = 290$ MeV

Sources: ☉ solar flare 17d20^h17^m, M1.3/2N, S29E07, AR8872


☾ solar flare 17d18^h41^m, M2.5/1B, S25W16*, AR8869

Main X-ray burst 1-8 Å: onset – 17d20^h17^m, max – 17d20^h45^m, $\Phi = 0.027$ J/m²



CME: 17d21^h30^m, $V = 728$ km/s, $\Delta\phi = 360^\circ$; $dA = 184^\circ$;

* - the contribution to the beginning

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 February 18



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



○

Integral fluxes of protons for the event of 2000 February 18


S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm2.s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	07h	13h	1.9	1d	
EPS	>10	06h	12h	1.7	1d	
EPS	>30	06h	11h	0.8	1d	
EPS	>50	06h	11h	0.42	1d	
EPS	>60	06h	10h	0.32	1d	
EPS	>100	06h	10h	0.14	1d	
POES-15						
MEPED	>0.24	-	10h	250	1d	
MEPED	>0.8	-	10h	228	1d	
MEPED	>2.5	-	10h	136	1d	
MEPED	>6.9	-	10h	80	1d	
MEPED	>16	-	10h	11.5	1d	
MEPED	>36	-	10h	3.27	1d	
MEPED	>70	-	10h	0.33	1d	
MEPED	>140	-	10h	0.18	1d	
IMP-8						
CPME	>1	11h	12h	64	4d	
CPME	>4	11h	12h	14.2	4d	
CPME	>10	11h	12h	4.8	3d	
CPME	>30	11h	12h	1.9	1.5d	
CPME	>60	11h	12h	1.1	1d	
ACE						
SIS	>10	09h	12h	29	3d	
SIS	>30	09h	12h	7.4	2d	
SOHO						
EPHIN (INT)	>50	11h	12h	0.23	1d	

Differential fluxes of protons for the event of 2000 February 18


S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	11 ^h	15 ^h	32	4d	
CPME	2-4.6	10 ^h	14 ^h	9.6	4d	
CPME	4.6-15	10 ^h	14 ^h	0.9	3d	
CPME	15-25	10 ^h	13 ^h	0.17	3d	
CPME	25-48	09 ^h	10 ^h	0.043	2d	
CPME	48-96	09 ^h	10 ^h	0.012	1.5d	
CPME	96-145	09 ^h	10 ^h	0.006	1d	
CPME	145-440	09 ^h	10 ^h	0.002	1d	
SOHO						
LION	0,75-2	12 ^h	15 ^h	29	4d	
LION	2-6	12 ^h	14 ^h	7.3	4d	
EPHIN	4-8	11 ^h	13 ^h	12.5	4d	
EPHIN	8-25	11 ^h	13 ^h	1.2	4d	
EPHIN	25-41	11 ^h	13 ^h	0.07	4d	
EPHIN	41-53	- ' -	- ' -	- ' -	- ' -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 February 18

2000	February 17	☉•			AR8872	To event 365	
H α	6563 Å	2031	2045	2218	S29E07	2N	FSU
1 -12.5	keV	2017	2035	2107		M1.3	2.7E-02
15.4	GHz	2024.0	2030.0	2034.0		1.76	
8.8	GHz	2024.0	2028.0	2039.0		1.72	
5	GHz	2024.0	2028.0	2039.0		1.91	
2.7	GHz	2023.0	2025.0	2039.0	U0.6 P1.4-2.7	2.30	
1.4	GHz	2023.0	2028.0	2039.0		2.28	
610	MHz	2024.0	2029.0	2039.0		1.76	
410	MHz	2024.0	2028.0	2039.0		2.08	
245	MHz	2024.0	2030.0	2039.0		2.45	
DS II	26-180	2025		2040	SH	3	
DS III	25-75	2024		2040		3	
CME		2130	0728 km/s	-22.9 km/s	360°	184	



2000	February 17	Ø	AR8869			To event 365	
Hα	6563 Å	1844	1847	1938	S25W16	1B	FU
1 -12.5	keV	1841	1852	1905		M2.5	2.3E-2
15.4	GHz	1844.0	1849.0	1859.0		2.00	
8.8	GHz	1847.0	1848.0	1855.0		2.04	
5	GHz	1846.0	1848.0	1855.0	U0.6 P5	2.15	
2.8	GHz	1836.0	1848.0	~1932.0		1.91	
1.4	GHz	1847.0	1847.0	1855.0		1.62	
610	MHz	1847.0	1848.0	1855.0		1.59	
410	MHz	1847.0	1847.0	1850.0		1.83	
245	MHz	1849.0	1851.0	1852.0		3.11	
245	MHz	1849.0	1851.0	1852.0		3.08	
DS II	25-75	1852		1905		2	
CME	WL	1931	0543 km/s	-82.5 km/s	064°	184°	



Particle event: To(Ep>10 MeV) – 04d17^h

Tmax₁(Ep>10 MeV) – 5d02^h, Jmax₁ (Ep>10 MeV) – 25 /cm².s.sr

Tmax₂(Ep>10 MeV) – 6d06^h, Jmax₂ (Ep>10 MeV) – 4 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 105 MeV


– Eqm₂ = 75 MeV

Sources: • solar flare 04d15^h11^m, C9.7/2F, N16W66, AR8933



Main X-ray burst 1-8 Å: onset – 04d15^h12^m, max – 04d15^h41^m, $\Phi = 0.023$ J/m²

CME: 04d16^h33^m, V = 1188 km/s, $\Delta\phi = 360^\circ$, dA = 265°

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 April 04



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 April 04

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	17 ^h	5d02 ^h /6d06 ^h	130.9/41.1	4d	
EPS	>10	17 ^h	5d02 ^h /6d06 ^h	25/4	3d	
EPS	>30	17 ^h	23 ^h /6d06 ^h	0.6/0.12	2d	
EPS	>50	17 ^h	22 ^h /6d06 ^h	0.11/0.03	1d	
EPS	>60	18 ^h	21 ^h /6d06 ^h	0.09/0.01	1d	
EPS	>100	18 ^h	20 ^h /6d06 ^h	0.015/0.005	1d	
POES-15						
MEPED	>0.24	18 ^h	5d10 ^h / -	3200/ -	4d	
MEPED	>0.8	18 ^h	5d10 ^h / -	2500/ -	4d	
MEPED	>2.5	18 ^h	5d10 ^h / -	1120/ -	4d	
MEPED	>6.9	18 ^h	5d10 ^h / -	36/ -	4d	
MEPED	>16	18 ^h	5d10 ^h / -	5.2/ -	4d	
MEPED	>36	18 ^h	5d10 ^h / -	0.46/ -	4d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	18 ^h	05d02 ^h /06d06 ^h	1200/1060	7 d	
CPME	>4	18 ^h	05d02 ^h /06d06 ^h	131/94	5 d	
CPME	>10	18 ^h	05d02 ^h /06d06 ^h	24/4	4 d	
CPME	>30	18 ^h	05d02 ^h /06d06 ^h	0.6/0.3	1 d	
CPME	>60	18 ^h	05d02 ^h /06d06 ^h	0.3/0.01	1 d	

ACE						
SIS	>10	18 ^h	05d06 ^h /06d18 ^h	46.5/19.1	2 d	
SIS	>30	18 ^h	21 ^h /06d18 ^h	1.5/ 0.1	1 d	
SOHO						
EPHIN (INT)	>50	18 ^h	21 ^h /06d15 ^h	0.03/0.1	2d	


Differential fluxes of protons for the event of 2000 April 04

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	19 ^h	- /10 ^h	- /950	8d	
CPME	2-4.6	19 ^h	- /10 ^h	- /143	8d	
CPME	4.6-15	19 ^h	- /10 ^h	- /6	5d	
CPME	15-25	19 ^h	- /10 ^h	- /0.07	3d	
CPME	25-48	19 ^h	- / -	- / -	2d	
CPME	48-96	19 ^h	- / -	- / -	1d	
CPME	96-145	19 ^h	- / -	- / -	1d	
CPME	145-440	19 ^h	- / -	- / -	1d	
SOHO						
LION	0,75-2	19 ^h	05d02 ^h /06d15 ^h	59/407	6d	
LION	2-6	19 ^h	05d02 ^h /06d15 ^h	13.4/55.5	6d	
EPHIN	4-8	19 ^h	05d02 ^h /06d15 ^h	20.3/38.7	8d	
EPHIN	8-25	18 ^h	05d03 ^h /06d15 ^h	1.3/0.6	5d	
EPHIN	25-41	17 ^h	05d00 ^h /06d15 ^h	0.028/0.001	4d	
EPHIN	41-53	- ' -	- ' -	- ' -	- ' -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 April 04

2000	April 04	•		AR8933	To event 366		
H α	6563 Å	1512	1533	1724	N16W66	2F	FS
1 -12.5	keV	1512	1541	1605		C9.7	2.3E-02
>300	keV	1523				8.7	O

15.4	GHz	1517.0	1526.0	1552.0		2.11	
8.8	GHz	1517.0	1527.0	1552.0		2.26	
6.7	GHz	1516.3	1526.2	1543.2		2.37	
2.7	GHz	1517.0	1519.0	1552.0	U0.6 P2.7	2.69	
1.4	GHz	1515.0	1524.0	1542.0		2.59	
610	MHz	1516.0	1518.0	1551.0		2.04	
410	MHz	1516.0	1518.0	1552.0		2.43	
280	MHz	1507.9	1522.0	1555.0		2.39	
127	MHz	1516.1	1517.8	1534.1		>2.18	
DS II	110-170	1524		1526	SH	3	
DS II	40-80	1530		1531	F,H	3	
DS IV	35-85	1515		1619		3	
DS IV	40-550	1517		1527	P	3	
DS III	30-80	1516		1539	N	3	
DS DCIM	800-2000	1514		1542	GG	2	
DS DCIM	2000-4500	1517		1543	GG	2	
CME	WL	1633	1188 km/s	12.8 km/s	360°	265°	



Particle event: To($E_p > 10$ MeV) – 07d00^h

Tmax($E_p > 10$ MeV) – 8d10^h, Jmax ($E_p > 10$ MeV) – 54 /cm².s.sr

Duration of the event – 3 days


Quasimaximal energy of protons in the event – $E_{qm} = 100$ MeV

Sources: • solar flare 06d11^h25^m, X2.3/3B, N20E18, AR9026



Main X-ray burst 1-8 Å: onset – 06d14^h28^m, max – 06d15^h25^m, $\Phi = 0.36$ J/m²

CME: 06d15^h54^m, $V = 1119$ km/s, $\Delta\phi = 360^\circ$, $dA = 047^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 June 07



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 June 07

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	01 ^h	08d10 ^h	520	3d	
EPS	>10	01 ^h	08d10 ^h	54	3d	
EPS	>30	00 ^h	08d09 ^h	0.6	3d	
EPS	>50	-	08d08 ^h	0.1	2d	
EPS	>60	-	08d07 ^h	0.07	2d	
EPS	>100	-	08d07 ^h	0.03	2d	
POES-15						
MEPED	>0.24	-	08d10 ^h	25310	3d	
MEPED	>0.8	-	08d10 ^h	13250	3d	
MEPED	>2.5	-	08d10 ^h	3290	3d	
MEPED	>6.9	-	08d10 ^h	320	2d	
MEPED	>16	-	08d10 ^h	6.5	2d	
MEPED	>36	-	08d10 ^h	0.6	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	6d23 ^h	08d11 ^h	13700	3 d	
CPME	>4	6d22 ^h	08d11 ^h	994	3 d	
CPME	>10	02 ^h	08d10 ^h	44	3 d	
CPME	>30	02 ^h	08d09 ^h	0.85	2 d	
CPME	>60	02 ^h	08d09 ^h	0.6	2 d	
ACE						
SIS	>10	01 ^h	08d12 ^h	42	3d	
SIS	>30	06d23 ^h	07d11 ^h	0.7	2d	

SOHO						
EPHIN (INT)	>50	06d22 ^h	08d10 ^h	0.13	2d	

Differential fluxes of protons for the event of 2000 June 07



S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	06d23 ^h	08d10 ^h	980	4d	
CPME	2-4.6	06d23 ^h	08d10 ^h	178	4d	
CPME	4.6-15	06d23 ^h	08d10 ^h	67	4d	
CPME	15-25	06d23 ^h	08d10 ^h	0.9	3d	
CPME	25-48	06d23 ^h	08d10 ^h	0.017	3d	
CPME	48-96	06d23 ^h	08d09 ^h	0.002	2d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	06d23 ^h	08d12 ^h	996	4d	
LION	2-6	07d00 ^h	08d11 ^h	176	4d	
EPHIN	4-8	06d22 ^h	08d12 ^h	205	4d	
EPHIN	8-25	06d21 ^h	08d09 ^h	7.5	3d	
EPHIN	25-41	06d19 ^h	08d09 ^h	0.023	3d	
EPHIN	41-53	- ' -	- ' -	- ' -	- ' -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 June 07

2000 June 06 • AR9026 To event 367

H α	6563 Å	1125	1317	1450	N20E13	2N	EFHKL
H α	6563 Å	1206	1521	1841	N20E14	3B	UZ
1 -12.5	keV	1310	1319	1328		M2.7	1.9E-2
1 -12.5	keV	1330	1339	1346		X1.1	8.1E-2
1 -12.5	keV	1356	1401	1411		M7.1	5.0E-2
1 -12.5	keV	1458	1525	1540		X2.3	3.6E-1
53 - 93	keV	<1543	1548	>1556	199		HXT Y
15.4	GHz	1330.0	1332.0	1349.0		2.62	
8.8	GHz	1330.0	1332.0	1338.0	U1.4 [P2.7 U5] P8.8	2.66	
6.7	GHz	1330.8	1331.2	1333.8		2.41	
5	GHz	1330.0	1333.0	1338.0		2.18	
2.7	GHz	1332.0	1333.0	1338.0		2.75	
1.4	GHz	1333.0	1334.0	1336.0		1.51	
245	MHz	1336.0	1337.0	1338.0		1.88	
DS III	35-70	1322		1449	N	1	
DS DCIM	2000-4500	1331		1334	GG,SP	2	

15.4	GHz	1508.0	1519.0	0000.0		3.28	
8.8	GHz	1508.0	1519.0	0000.0		3.53	
6.7	GHz	1505.0	1516.5	1536.6	P0.6 U2.7 P6.7	3.67	
5	GHz	1506.0	1519.0	0000.0		3.45	
2.7	GHz	1506.0	1519.0	0000.0		3.36	
1.4	GHz	1506.0	1519.0	0000.0		3.81	
610	MHz	1507.0	1523.0	0000.0		4.26	
410	MHz	1507.0	1521.0	0000.0		3.88	
245	MHz	1507.0	1527.0	1607.0		3.63	
DS II	220-550	1516		~1530		3	
DS II	25-119	1528		1539		2	
DS IV	30-80	1457		2121		3	
DS III	220-550	1507		1509	GG,RS	3	
DS III	25-158	1507		1509		1	
DS DCIM	2000-4500	1503		1553	GG	3	
DS DCIM	400-4000	1520		1710	P,Z	3	
CME	WL	1506	0358 km/s	-3.9 km/s	051°	018°	
CME	WL	1530	0929 km/s	111.9 km/s ²	090°	004°	
CME	WL	1554	1119 km/s	1.5 km/s ²	360°	047°	



Particle event: To($E_p > 10$ MeV) – 10d17^h

Tmax($E_p > 10$ MeV) – 10d20^h, Jmax ($E_p > 10$ MeV) – 24 /cm².s.sr

Duration of the event – 4 days

Quasimaximal energy of protons in the event – $E_{qm} = 390$ MeV


Sources: • solar flare 10d16^h22^m, M5.2/3B, N22W39, AR9026

Main X-ray burst 1-8 Å: onset – 10d16^h40^m, max – 10d17^h02^m, $\Phi = 0.073$ J/m²



CME: 10d17^h08^m, $V = 1108$ km/s, $\Delta\phi = 360^\circ$, dA = 307°

Δ SC 11d08^h01^m, Δ SC 12d22^h08^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 June 10



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 June 10

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	17 ^h	20 ^h	34.4	>3d	
EPS	>10	17 ^h	20 ^h	24	>3d	
EPS	>30	17 ^h	19 ^h	6.9	>2d	
EPS	>50	17 ^h	18 ^h	2.8	>1d	
EPS	>60	17 ^h	18 ^h	1.96	>1d	
EPS	>100	17 ^h	18 ^h	0.73	1d	
POES-15						
MEPED	>0.24	17 ^h	19 ^h	137	3d	
MEPED	>0.8	17 ^h	19 ^h	137	3d	
MEPED	>2.5	17 ^h	19 ^h	137	3d	
MEPED	>6.9	17 ^h	19 ^h	119	3d	
MEPED	>16	17 ^h	19 ^h	41.5	3d	
MEPED	>36	17 ^h	19 ^h	9.2	3d	
MEPED	>70	17 ^h	19 ^h	1.2	3d	
MEPED	>140	17 ^h	19 ^h	0.48	3d	
IMP-8						
CPME	>1	14 ^h	23 ^h	303	5 d	
CPME	>4	14 ^h	21 ^h	73.2	4 d	
CPME	>10	15 ^h	19 ^h	31.1	4 d	
CPME	>30	15 ^h	19 ^h	6.8	1.5 d	
CPME	>60	16 ^h	19 ^h	2.1	1.5 d	
ACE						
SIS	>10	17 ^h	18 ^h	92	3d	
SIS	>30	17 ^h	18 ^h	32.5	2d	
SOHO						
EPHIN (INT)	>50	-	22 ^h	0.85	1.5d	



Differential fluxes of protons for the event of 2000 June 10

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	21 ^h	11d01 ^h	197	4d	
CPME	2-4.6	19 ^h	23 ^h	56.4	4d	
CPME	4.6-15	19 ^h	21 ^h	4.9	4d	
CPME	15-25	18 ^h	18 ^h	1.8	3d	
CPME	25-48	18 ^h	19 ^h	0.25	3d	
CPME	48-96	16 ^h	19 ^h	0.052	2d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	19 ^h	21 ^h	46	5d	
LION	2-6	19 ^h	21 ^h	12.7	5d	
EPHIN	4-8	19 ^h	21 ^h	28.4	5d	
EPHIN	8-25	19 ^h	20 ^h	4.9	5d	
EPHIN	25-41	19 ^h	19 ^h	1.1	5d	
EPHIN	41-53	- ' -	- ' -	- ' -	- ' -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 June 10

2000	June 10	•	AR9026	To event 368			
H α	6563 Å	1622	1653	1810	N22W39	3B	FUY
1 -12.5	keV	1640	1702	1719		M5.2	7.3E-2
53 – 93	keV	<1652	1653	1715		18	HXT Y
15.4	GHz	1647.0	1653.0	0000.0		2.36	
8.8	GHz	1648.0	1653.0	0000.0		2.58	
6.7	GHz	1647.0	1653.1	1658.1	[U2.7 P6.7]	3.40	
5	GHz	1648.0	1653.0	0000.0		2.88	
2.7	GHz	1648.0	1653.0	0000.0		2.74	
2.7	GHz	1648.0	1653.0	0000.0		2.74	
1.4	GHz	1647.0	1653.0	1706.0		3.18	
610	MHz	1653.0	1700.0	1708.0		3.73	
410	MHz	1652.0	1700.0	1703.0		1.72	
280	MHz	1653.3	1703.0	1710.1		2.09	
DS II	25-180	1655		1714		2	
DS II	30-80	1655		1718		3	
DS III	25-180	1655		1659		1	
DS DCIM	1000-4000	1647		1716	P,C	3	

1.4	GHz	1756.0	1800.0	1819.0		2.23	
610	MHz	1737.0	1741.0	0000.0	P0.6	3.97	
410	MHz	1738.0	1741.0	0000.0		3.46	
DS DCIM	1000-1500	1757		1806	P,Z	3	
CME	WL	1708	1108 km/s	-21.2km/s^2	360°	307°	



Particle event: To($E_p > 10$ MeV) – 17d07^h

Tmax($E_p > 10$ MeV) – 18d06^h, Jmax ($E_p > 10$ MeV) – $1.7 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}^*$)

Duration of the event – 2 days *)

Quasimaximal energy of protons in the event – $E_{qm} = 110$ MeV

*) According to data IMP-8 (CPME)


Sources: • solar flare 17d02^h25^m, M3.5/2B, N22W72, AR9033

Ø solar flare 18d01^h52^m, X1.0/SF, N23W85, AR9033



Main X-ray burst 1-8 Å: onset – 17d02^h25^m, max – 17d02^h37^m, $\Phi = 0.024 \text{ J/m}^2$

CME: 17d03^h28^m, $V = 857 \text{ km/s}$, $\Delta\varphi = 133^\circ$, $dA = 301^\circ$;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 June 17



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 June 17

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	07 ^h	18d05 ^h	0.6	1d	
EPS	>10	07 ^h	18d04 ^h	0.4	1d	
EPS	>30	07 ^h	18d04 ^h	0.13	1d	
EPS	>50	07 ^h	18d04 ^h	0.07	1d	
EPS	>60	07 ^h	18d04 ^h	0.04	1d	
EPS	>100	07 ^h	18d04 ^h	0.015	1d	
POES-15						
MEPED	>0.24	14 ^h	18d04 ^h	371	3d	
MEPED	>0.8	14 ^h	18d04 ^h	73	3d	
MEPED	>2.5	14 ^h	18d04 ^h	31	3d	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	-	-	-	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	10 ^h	18d03 ^h	356	3 d	
CPME	>4	08 ^h	18d05 ^h	14	2 d	
CPME	>10	07 ^h	18d06 ^h	1.7	2 d	
CPME	>30	-	18d06 ^h	0.13	1.5 d	
CPME	>60	-	18d06 ^h	0.04	0.5 d	
ACE						
SIS	>10	07 ^h	18d06 ^h	1.3	1d	
SIS	>30	07 ^h	18d04 ^h	0.2	1d	
SOHO						
EPHIN (INT)	>50	-	18d05 ^h	0.008	-	

Differential fluxes of protons for the event of 2000 June 17



S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	09 ^h	18d02 ^h	274	3d	
CPME	2-4.6	08 ^h	18d03 ^h	34.6	3d	
CPME	4.6-15	06 ^h	18d03 ^h	0.77	2d	
CPME	15-25	05 ^h	18d01 ^h	0.045	1,5d	
CPME	25-48	05 ^h	18d05 ^h	0.008	0,5d	
CPME	48-96	-	18d05 ^h	0.0025	0,5d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	06 ^h	18d06 ^h	30.3	3d	
LION	2-6	06 ^h	18d06 ^h	3.9	3d	
EPHIN	4-8	06 ^h	18d05 ^h	3.9	3d	
EPHIN	8-25	05 ^h	18d05 ^h	0.81	3d	
EPHIN	25-41	05 ^h	18d04 ^h	0.006	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 June 17

2000 June 17 • AR9033 To event 369

H α	6563 Å	0228	0238	0501	N22W72	2B	EF
1 -12.5	keV	0225	0237	0244		M3.5	2.4E-2
50 – 150	keV	0228	0235	0337		206	HXT Y
15.4	GHz	0231.0	0231.0	0232.0		1.49	
8.8	GHz	0230.0	0231.0	0235.0	U1.4 P5.7-8.8	1.98	
5.7	GHz	0230.3	0232.0	0405.0		1.98	
2.7	GHz	0231.0	0233.0	0235.0		1.79	
1.4	GHz	0231.0	0234.0	0235.0		1.49	
245	MHz	0236.0	0236.0	~0236.0		1.73	
DS III	25-150	0247		0332	N	2	
DS CONT	30-80	0250		0413		1	
5	GHz	0258.0	0305.0	0310.0	U2.7 / 5	1.75	
2.7	GHz	0307.0	0307.0	~0307.0		1.36	
1.4	GHz	0305.0	0307.0	0309.0		1.76	
610	MHz	0306.0	0307.0	0309.0		2.00	
410	MHz	0306.0	0307.0	0308.0		2.04	
245	MHz	0304.0	0304.0	~0304.0		1.69	
DS III	23-600	0301		0309	GG	3	


5	GHz	0353.0	0353.0	0354.0	[0.9 - 5]	1.77	
2.7	GHz	0351.0	0352.0	0356.0		1.94	
1.4	GHz	0346.0	0346.0	0347.0		1.91	
900	MHz	0345.0	0352.2			1.95	
610	MHz	0345.0	0352.0	0357.0		2.26	
410	MHz	0345.0	0347.0	0349.0		2.00	
245	MHz	0340.0	0342.0	0000.0		2.90	
DS III	160-260	0340		0346	G	2	
DS CONT	160-820	0341		0355		1	
CME	WL	0328	0857 km/s	16.4 km/s ²	133°	301°	





2000 June 18 Ø AR9033 To event 369

H α	6563 Å	0156	0157	0213	N23W85	SF	EF
1 -12.5	keV	0152	0159	0213		X1.0	3.4E-2
53 – 93	keV	>020422	~020426	020912		188	HXT Y
15.4	GHz	0156.0	0157.0	0158.0	U0.6 P8.8-15.4	2.75	
8.8	GHz	0156.0	0157.0	0158.0		2.76	
5	GHz	0155.0	0157.0	0159.0		2.53	
2.7	GHz	0155.0	0157.0	0159.0		2.48	
1.4	GHz	0156.0	0157.0	0200.0		2.45	
610	MHz	0156.0	0157.0	0200.0		2.23	
410	MHz	0157.0	0158.0	0200.0		3.62	
245	MHz	0156.0	0158.0	0203.0		4.08	
DS II	30-650	0157		0211	FN	3	
DS III	18-180	0157		0204	GG	3	
BSL	6563 Å	0156		0233	N17W90		
CME	WL	0210	0629 km/s	-1.2 km/s ²	132°	318°	



Particle event: To($E_p > 10$ MeV) – 25d10^h

Tmax($E_p > 10$ MeV) – 26d07^h, Jmax ($E_p > 10$ MeV) – 1.5 /cm².s.sr

Duration of the event – 3 days


Quasimaximal energy of protons in the event – $E_{qm} = 70$ MeV

Sources: ☉ solar flare 25d07^h17^m, M1.9/2N, N16W55, AR9046



Main X-ray burst 1-8 Å : onset – 25d07^h 17^m, max – 25d07^h52^m, $\Phi = 0.043$ J/m²

CME: 25d07^h54^m, $V = 0546$ km/s, $\Delta\phi = 165^\circ$, $dA = 274^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 June 25



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 June 25

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES-10						
EPS	>5	10 ^h	26d07 ^h	14	2d	
EPS	>10	10 ^h	26d07 ^h	1.5	1.5d	
EPS	>30	11 ^h	26d07 ^h	0.06	1d	
EPS	>50	-	-	-		
EPS	>60	-	-	-		
EPS	>100	-	-	-		
POES-15						
MEPED	>0.24	-	26d04 ^h	1320	3 d	
MEPED	>0.8	-	26d04 ^h	785	3 d	
MEPED	>2.5	-	26d04 ^h	118	3 d	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	-	-	-	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	12 ^h	26d01 ^h	655	3 d	
CPME	>4	12 ^h	26d01 ^h	36	3 d	
CPME	>10	12 ^h	26d01 ^h	2.4	2 d	
CPME	>30	-	-	-	-	
CPME	>60	-	-	-	-	
ACE						
SIS	>10	11 ^h	23 ^h	2.3	1.5d	
SIS	>30	12 ^h	22 ^h	0.2	1d	

SOHO						
EPHIN (INT)	>50	-	-	-	-	


Differential fluxes of protons for the event of 2000 June 25

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	12 ^h	26d08 ^h	313	~3d	
CPME	2-4.6	12 ^h	26d08 ^h	38	~3d	
CPME	4.6-15	11 ^h	26d08 ^h	1.2	~3d	
CPME	15-25	11 ^h	26d08 ^h	0.02	~1,5d	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	12 ^h	26d05 ^h	102	3.5d	
LION	2-6	12 ^h	26d05 ^h	12.1	3.5d	
EPHIN	4-8	12 ^h	26d01 ^h	9.6	3.5d	
EPHIN	8-25	11 ^h	26d01 ^h	0.3	3d	
EPHIN	25-41	10 ^h	26d10 ^h	0,0014	2.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 June 25

2000 June 25		☉		AR9046		To event 370	
H α	6563 Å	0720	0741	0853	N16W55	2N	FH
1 -12.5	keV	0717	0752	0821		M1.9	4.3E-02
3	GHz	0716.2	0720.8	0726.8		1.49	
245	MHz	0719.0	0720.0	0721.0		1.82	
204	MHz	0720.2	0720.6	0721.5		2.16	
DS III	220-450	0720		0721	GG,S	3	
DS III	50-270	0720		0721	GG	2	
DS DCIM	2088-4500	0720		0721	G	1	

8.8	GHz	0741.0	0741.0	0000.0		1.48	
5	GHz	0740.0	0741.0	0744.0		1.52	
3	GHz	0733.9	0741.6	0801.4		1.60	
1.4	GHz	0741.0	0741.0	0742.0		1.49	
900	MHz	0715.0	0735.6			2.06	
600	MHz	0719.5	0738.6		[P0.6]	2.15	
410	MHz	0740.0	0741.0	0742.0		1.91	
204	MHz	0736.8	0745.8	0752.0		1.88	
DS II	40-90	0751		0759		2	
DS I	50-160	0738		~0804	N,C	2	
DS III	45-95	0742		~0808	N	2	
DS CONT	25-180	0739		0805		1	
DS DCIM	800-2000	0730		0746	GG,SP	2	
DS DCIM	220-550	0734		0747	C	2	
CME	WL	0754	1617 km/s	-17.5 km/s ²	165°	274°	



Particle event: To($E_p > 10$ MeV) – 13d06^h

Tmax($E_p > 10$ MeV) – 13d10^h, Jmax ($E_p > 10$ MeV) – 5 /cm².s.sr

Duration of the event – 1.5 days

Quasimaximal energy of protons in the event – $E_{qm} = 40$ MeV


Sources: ☉ solar flare 12d18^h41^m, M5.7/2F, N16W64, AR9070

Main X-ray burst 1-8 Å : onset – 12d18^h41^m, max – 12d18^h47^m, $\Phi = 0.063$ J/m²



CME: 12d20^h30^m, $V = 820$ km/s, $\Delta\phi = 101^\circ$, $dA = 281^\circ$

▲ SC 13d09^h42^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 July 13



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 July 13

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES 10						
EPS	>5	06 ^h	15 ^h	45	1.5d	
EPS	>10	06 ^h	15 ^h	5	1.5d	
EPS	>30	06 ^h	14 ^h	0.08	1d	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-15						
MEPED	>0.24	-	09 ^h	11490	1.5 d	
MEPED	>0.8	-	09 ^h	3480	1.5 d	
MEPED	>2.5	-	09 ^h	362	1.5 d	
MEPED	>6.9	-	09 ^h	7	1.5 d	
MEPED	>16	-	09 ^h	0.6	1.5 d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	02 ^h	10 ^h	2920	1.5 d	
CPME	>4	02 ^h	10 ^h	62	1.5 d	
CPME	>10	02 ^h	10 ^h	2.3	1.5 d	
CPME	>30	-	-	-	-	
CPME	>60	-	-	-	-	
ACE						
SIS	>10	12d20 ^h	14 ^h	2.5	1.5 d	
SIS	>30	-	-	-	1.5 d	
SOHO						
EPHIN (INT)	>50	-	14 ^h	0.008	-	


Differential fluxes of protons for the event of 2000 July 13

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	12d17 ^h	10 ^h	2630	2d	
CPME	2-4.6	12d17 ^h	10 ^h	200	2d	
CPME	4.6-15	12d17 ^h	10 ^h	3.85	2d	
CPME	15-25	-	-	-	-	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	12d19 ^h	12 ^h	111	1.5d	
LION	2-6	12d19 ^h	12 ^h	10.8	1.5d	
EPHIN	4-8	12d20 ^h	12 ^h	5.5	1.5d	
EPHIN	8-25	12d20 ^h	11 ^h	0.16	1.5d	
EPHIN	25-41	12d20 ^h	10 ^h	0.0014	1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 July 13

2000 July 12		☉		AR9070		To event 371	
H α	6563 Å	<1849	1848	>1957	N16W64	2F	
1 -12.5	keV	1841	1849	1907		M5.7	6.3E-2

15.4	GHz	1841.0	1848.0	1858.0	1.4 / 15.4	2.98	
8.8	GHz	1843.0	1848.0	1853.0		2.81	
5	GHz	1843.0	1848.0	1855.0		2.62	
2.7	GHz	1843.0	1848.0	1853.0		2.38	
1.4	GHz	1844.0	1848.0	1852.0		1.77	
DS III	25-86	1848		1849		1	
CME	WL	2030	0820 km/s	-3.2 km/s ²	101°	281°	



Particle event: To($E_p > 10$ MeV) – 14d10^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 14\text{d}18^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 7.2 \cdot 10^3 \text{ cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 15\text{d}13^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 1.8 \cdot 10^4 \text{ cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm1} = 2160 \text{ MeV}$

– $E_{qm2} = 630 \text{ MeV}$

Sources: ● solar flare 14d10^h03^m, X5.7/3B, N22W07, AR9077


⊙ solar flare 15d08^h20^m, M1.3/SF, N16W12, AR9077

Main X-ray burst 1-8 Å: onset – 14d10^h03^m, max – 14d10^h24^m, $\Phi = 0.75 \text{ J/m}^2$



CME: 14d10^h54^m, $V = 1674 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 273^\circ$

▲ SC14d15^h32^m, ▲ SC 15d14^h37^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 July 14



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 July 14

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	10 ^h	18 ^h /15d13 ^h	8600/30475	2d	
EPS	>10	10 ^h	18 ^h /15d13 ^h	7200/18000	2d	
EPS	>30	10 ^h	18 ^h /15d12 ^h	2860/4070	2d	
EPS	>50	10 ^h	17 ^h /15d12 ^h	1300/1160	2d	
EPS	>60	10 ^h	17 ^h /15d12 ^h	805/514	2d	
EPS	>100	10 ^h	16 ^h /15d12 ^h	330/91.3	2d	
POES-16						
MEPED	>0.24	-	13 ^h /15d15 ^h	26430/192050	2d	
MEPED	>0.8	-	13 ^h /15d15 ^h	16250/83230	2d	
MEPED	>2.5	-	13 ^h /15d15 ^h	11690/47710	2d	
MEPED	>6.9	-	13 ^h /15d15 ^h	10120/26470	2d	
MEPED	>16	-	13 ^h /15d15 ^h	3250/11080	2d	
MEPED	>36	-	13 ^h /15d15 ^h	2280/2900	2d	
MEPED	>70	-	13 ^h /15d15 ^h	570/170	2d	
MEPED	>140	-	13 ^h /15d15 ^h	180/12	2d	
IMP-8						
CPME	>1	11 ^h	16 ^h /15d17 ^h	6700/32400	2d	
CPME	>4	10 ^h	16 ^h /15d17 ^h	6000/8440	2d	
CPME	>10	10 ^h	16 ^h /15d17 ^h	4960/3960	2d	
CPME	>30	10 ^h	17 ^h /15d16 ^h	2270/675	2d	
CPME	>60	10 ^h	16 ^h /15d16 ^h	1200/250	2d	
ACE						
SIS	>10	10 ^h	16 ^h / -	4040/ -	2d	
SIS	>30	10 ^h	16 ^h / -	1770/ -	2d	

SOHO						
EPHIN (INT)	>50	10 ^h	17 ^h /15d13 ^h	1760/1510	2d	
BALLOONS						
Mu-1	>173	-	14 ^h -16 ^h / -	29/ -	-	
Mu-1	>402	-	14 ^h -16 ^h / -	2.3/ -	-	
Mu-1	>609	-	14 ^h -16 ^h / -	0.28/ -	-	
NM						
Network	>433	-	11 ^h 10 ^m / -	34	-	
Network	>1000	-	11 ^h 10 ^m / -	0.37	-	
Network	>2000	-	11 ^h 10 ^m / -	0.0088	-	
Network	>3700	-	11 ^h 10 ^m / -	0.00032	-	

Differential fluxes of protons for the event of 2000 July 14

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	10 ^h	19 ^h /15d13 ^h	167/15900	2d	
CPME	2-4.6	10 ^h	19 ^h /15d13 ^h	255/3850	2d	
CPME	4.6-15	10 ^h	19 ^h /15d13 ^h	183/435	2d	
CPME	15-25	09 ^h	18 ^h /15d13 ^h	234/208	2d	
CPME	25-48	09 ^h	18 ^h /15d13 ^h	61/23.6	2d	
CPME	48-96	09 ^h	17 ^h /15d13 ^h	24.5/4.6	2d	
CPME	96-145	09 ^h	17 ^h /15d13 ^h	8.3/6.5	2d	
CPME	145-440	09 ^h	17 ^h /15d13 ^h	1/0.17	2d	
SOHO						
LION	0,75-2	10 ^h	17 ^h /15d13 ^h	1315/8600	2d	
LION	2-6	10 ^h	17 ^h /15d13 ^h	201/1330	2d	
EPHIN	4-8	10 ^h	18 ^h /15d14 ^h	52/1001	2d	
EPHIN	8-25	10 ^h	18 ^h /15d15 ^h	41/766	2d	
EPHIN	25-41	10 ^h	18 ^h /15d11 ^h	24.5/36.7	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References


- Belov A.V., J.W. Bieber, E.A. Eroshenko et al., 2001.
Share G.H., R.J. Murphy, A.J. Tylka et al., 2001.
Smith C.W., N.F. Ness, L.F. Burlaga et al., 2001.
Caroubalos C. C.E. Alissandrakis, A. Hillaris et al., 2001.
Ding L., 2001.
Tonwar S., 2001.
Kallenrode M.-B. and E.W. Cliver, 2001.
Quack M., M.-B. Kallenrode, M. von König et al., 2001.
Vashenyuk E.V., B.B. Gvozdevsky, V.V. Pchelkin et al., 2001.
Lepping R.P., D.B. Berdichevsky, L.F. Burlaga et al., 2001.
Watari S., M. Kunitake, and T. Watanabe, 2001.
Leske R.A., R.A. Mewaldt, A.C. Cummings et al., 2001.
Bieber J.W., W. Dröge, P.A. Evenson et al., 2002.
Zhao H., G. Zhu, S. Wang et al., 2002.
Struminsky A.B., 2003.
Duldig M.L., D.J. Bombardieri, and J.E. Humble, 2003.
Perez-Peraza J., A. Gallegos-Cruz, E.V. Vashenyuk et al., 2003.

Ma Yu.Q., 2003.
 Nitta N.V., E.W. Cliver, A.J. Tylka et al., 2003.
 Wan W., L. Liu, H. Yuan et al., 2005.
 Le G.M., 2005.
 Tang Y.H., Y. Dai, 2003.
 Iles R.H., J.B.L. Jones, G.C. Taylor et al., 2004.
 Al-Thoyaib S.S., 2005.
 Achard P., O. Adriani, M. Aguilar-Benitez et al., 2006.
 Bombardieri D.J., M.L. Duldig, K.J. Michael et al., 2006.
 Wang R.G. and J.X. Wang, 2006.
 Rawat R., S. Alex, and G.S. Lakhina, 2006.
 Dmitriev A.V., H.-C. Yeh, J.-K. Chao et al., 2006.
 El-Borie M.A., 2003a.
 El-Borie M.A., 2003b.
 Li C., Y.H. Tang, Y. Dai et al., 2007.
 Miroshnichenko L.I. and J. Perez-Peraza, 2008.
 Lario D., R.B. Decker, and A. Aran, 2008.
 Lario D, A. Aran, R.B. Decker, 2009
 Perez-Peraza J., A. Gallegos-Cruz, E.V. Vashenyuk et al., 2009.
 Wang R.G., 2009.
 Miroshnichenko L.I., J.A. Pérez-Peraza, V.M. Velasco-Herrera et al., 2012
 Somov B.V., 2012.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2000 July 14**

2000 July 14		• AR9077		To event 372			
H α	6563 Å	<1012	~1021	1330	N22W07	3B	U
1 -12.5	keV	1003	1024	1043		X5.7	7.5E-1
50 – 150	keV	101113	102011	120115		8244	HXT Y

15.4	GHz	1012.0	1027.0	1132.0		3.83	
8.8	GHz	1007.0	1027.0	1157.0	P0.4 U3 P8.8	3.95	
5	GHz	1005.0	1027.0	1159.0		3.89	
3	GHz	1001.3	1028.6	1240.7		3.51	
1.4	GHz	1006.0	1030.0	0000.0		3.61	
610	MHz	1009.0	1032.0	1132.0		4.52	
410	MHz	1010.0	1031.0	1132.0		4.61	
245	MHz	1011.0	1036.0	1132.0		4.49	
204	MHz	1013.9	1028.2	1043.8		4.24	
DS II	220-440	1017		1027	H	2	
DS II	35-85	1020		1026		3	
DS IV	30-80	1026		1330		3	
DS III	25-270	1015		~1039	S,C	2	
DS CONT	25-270	1026		~1041		3	
DS DCIM	800-2000	1003		1151	GG,FS	3	
DS DCIM	2000-4500	1007		1135	GG	3	
CME	WL	1054	1674 km/s	-96.1 km/s ²	360°	273°	



2000	July 15	☉		AR9087		To event 372	
H α	6563 Å	0822	0826	1146	N16W12	SF	U
1 -12.5	keV	0820	0833	0848		M1.3	1.8E-02
3	GHz	0831.6	0832.2	0836.7		1.18	
DS III	25-160	0832		0833	G	2	
DS III	30-210	0839		0839	G	2	
DS III	25-160	0843		0845	GG FS	2	
DS III	200-270	0903		0904	GG	1	

Particle event: To(Ep>10 MeV) – 16d11^h

Tmax₁(Ep>10 MeV) – 16d12^h, Jmax₁(Ep>10 MeV) – 100 /cm².s.sr

Tmax₂(Ep>10 MeV) – 17d02^h, Jmax₂(Ep>10 MeV) – 37 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 370 MeV

– Eqm₂ = 320 MeV


Sources: ☉ solar flare 15d08^h20^m, M1.3/SF, N16W12 AR9077

☽ solar flare 16d23^h37^m, M1.4/2F, N17W40 AR9077



Main X-ray burst 1-8 Å: onset – 15d08^h20^m, max – 15d08^h33^m, Φ = 0.018 J/m²

▲ SC 15d14^h37^m, ▲ SC 19d15^h27^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 July 16



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 July 16

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	11 ^h	12 ^h /17d03 ^h	236/75.8	3d	
EPS	>10	11 ^h	12 ^h /17d02 ^h	100/37	3d	
EPS	>30	11 ^h	12 ^h /17d02 ^h	15.7/7.6	3d	
EPS	>50	11 ^h	12 ^h /17d02 ^h	5.4/2.6	3d	
EPS	>60	11 ^h	12 ^h /17d02 ^h	2.1/1.0	3d	
EPS	>100	11 ^h	12 ^h /17d01 ^h	0.6/0.26	3d	
POES-15						
MEPED	>0.24	11 ^h	12 ^h /17d02 ^h	4660/1825	3d	
MEPED	>0.8	11 ^h	12 ^h /17d02 ^h	3254/1290	3d	
MEPED	>2.5	11 ^h	12 ^h /17d02 ^h	1690/712	3d	
MEPED	>6.9	11 ^h	12 ^h /17d02 ^h	620/294	3d	
MEPED	>16	11 ^h	12 ^h /17d02 ^h	184/84	3d	
MEPED	>36	11 ^h	12 ^h /17d02 ^h	32/14	3d	
MEPED	>70	11 ^h	12 ^h /17d02 ^h	5.2/2.1	3d	
MEPED	>140	11 ^h	12 ^h /17d02 ^h	1./0.9	3d	

IMP-8						
CPME	>1	11 ^h	13 ^h /17d03 ^h	3210/1060	3d	
CPME	>4	11 ^h	12 ^h /17d02 ^h	810/323	3d	
CPME	>10	10 ^h	12 ^h /17d02 ^h	258/110	3d	
CPME	>30	10 ^h	11 ^h /17d02 ^h	35.2/13.6	3d	
CPME	>60	10 ^h	11 ^h /17d01 ^h	13.4/5.6	3d	
ACE						
SIS	>10	10 ^h	-	-	-	
SIS	>30	10 ^h	-	-	-	
SOHO						
EPHIN (INT)	>50	11 ^h	15 ^h /17d03 ^h	4.3/1.7	3d	


Differential fluxes of protons for the event of 2000 July 16

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	10 ^h	12 ^h /17d02 ^h	1610/508	3d	
CPME	2-4.6	10 ^h	12 ^h /17d03 ^h	466/145	3d	
CPME	4.6-15	10 ^h	12 ^h /17d02 ^h	50.6/19.8	3d	
CPME	15-25	10 ^h	12 ^h /17d03 ^h	12.3/6.1	3d	
CPME	25-48	10 ^h	11 ^h /17d02 ^h	1.35/0.5	3d	
CPME	48-96	10 ^h	11 ^h /17d02 ^h	0.15/0.05	3d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	11 ^h	16 ^h /17d03 ^h	332/115	3d	
LION	2-6	11 ^h	16 ^h /17d03 ^h	67.4/23.5	3d	
EPHIN	4-8	11 ^h	15 ^h /17d03 ^h	158/63.2	3d	
EPHIN	8-25	11 ^h	16 ^h /17d03 ^h	20.8/9.4	3d	
EPHIN	25-41	11 ^h	15 ^h /17d03 ^h	1.2/0.56	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 July 16

2000	July 15	☉			AR9090	To event 373	
Hα		0159	0203	0214	N09E81	1N	U
1 -12.5	keV	0159	0203	0214		M5.5	1.6E-02

15.4	GHz	0202.0	0202.0	0204.0		2.63	
8.8	GHz	0202.0	0202.0	0204.0	U2.8 P8.8	2.86	
5	GHz	0202.0	0202.0	0205.0		2.59	
2.8	GHz	0200.0	0204.0	0210.0		1.40	
610	MHz	0147.0	0156.0	0000.0		2.68	
DS III	35-90	0203		0203	B	1	
DS III	30-75	0225		0227	G	1	



2000		July 16		Ø	AR9077	To event 373	
H α		2341	2352	0051	N17W40	2F	EFU
1 -12.5	keV	2337	0004	0015		M1.4	1.7E-02
2.8	GHz	2333.0	2342.0	0143.0		1.40	
610	MHz	2354.0	2355.0	2356.0		1.91	
DS III	28-180	2310		2339	N	1	
DS III	57-180	2317		2324	N	1	

Particle event: To(Ep>10 MeV) – 22d12^h

Tmax₁(Ep>10 MeV) – 22d14^h, Jmax₁ (Ep>10 MeV) – 13 /cm².s.sr

Tmax₂(Ep>10 MeV) – 22d20^h, Jmax₂ (Ep>10 MeV) – 6 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 340 MeV

– Eqm₂ = 80 MeV


Sources: ● solar flare 22d11^h17^m, M3.7/2N, N14W56, AR9085

Main X-ray burst 1-8 Å : onset – 22d11^h17^m, max – 22d11^h34^m, Φ = 0.07 J/m²



CME: 22d11^h54^m, V = 1230 km/s, Δφ = 229°, dA = 275°;

▲ SC 23d10^h41^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 July 22



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 July 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES-10						
EPS	>5	12 ^h	17 ^h /20 ^h	27/20.4	2d	
EPS	>10	12 ^h	14 ^h /20 ^h	13/6	2d	
EPS	>30	12 ^h	13 ^h /20 ^h	3.1/0.55	1d	
EPS	>50	12 ^h	13 ^h /20 ^h	1.1/0.12	1d	
EPS	>60	12 ^h	13 ^h /20 ^h	0.65/0.07	1d	
EPS	>100	12 ^h	12 ^h / -	0.24/ -	1d	
POES-15						
MEPED	>0.24	-	- /24 ^h	- /940	2d	
MEPED	>0.8	-	- /24 ^h	- /650	2d	
MEPED	>2.5	-	- /24 ^h	- /200	2d	
MEPED	>6.9	-	- /24 ^h	- /30	2d	
MEPED	>16	-	- /24 ^h	- /2.5	1d	
MEPED	>36	-	- /24 ^h	- /0.14	1d	
MEPED	>70	-	- /24 ^h	- /0.1	1d	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	13 ^h	16 ^h /22 ^h	42/344	1.5 d	
CPME	>4	12 ^h	16 ^h /22 ^h	22/38	1.5 d	
CPME	>10	12 ^h	15 ^h /20 ^h	8/4.7	1.5d	
CPME	>30	12 ^h	13 ^h /18 ^h	1.8/0.3	1 d	
CPME	>60	12 ^h	13 ^h /18 ^h	0.75/0.05	1 d	

ACE						
SIS	>10	12 ^h	15 ^h /19 ^h	11/6.5	1.5d	
SIS	>30	12 ^h	13 ^h /18 ^h	3.2/0.1	1d	
SOHO						
EPHIN (INT)	>50	12 ^h	14 ^h /20 ^h	0.28/0.04	1d	


Differential fluxes of protons for the event of 2000 July 22

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	15 ^h	-	- /189	2d	
CPME	2-4.6	14 ^h	16 ^h /14 ^h	8.7/61	2d	
CPME	4.6-15	11 ^h	15 ^h	1.5/3.7	2d	
CPME	15-25	11 ^h	13 ^h	0.37/0.22	1.5d	
CPME	25-48	11 ^h	13 ^h	0.06/0.018	1.5d	
CPME	48-96	11 ^h	13 ^h	0.013/0.0015	1d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	14 ^h	16 ^h /23 ^h	2.5/43	2d	
LION	2-6	14 ^h	17 ^h /23 ^h	1/7.3	2d	
EPHIN	4-8	14 ^h	16 ^h /22 ^h	3.1/12.8	2d	
EPHIN	8-25	14 ^h	16 ^h /22 ^h	0.6/0.75	2d	
EPHIN	25-41	13 ^h	15 ^h /21 ^h	0.06/0.022	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 July 22

2000	July 22	•	AR9085	To event 374			
Hα	6563 Å	1117	1125	1246	N14W56	2N	FU
1 -12.5	keV	1117	1134	1202		M3.7	7.0E-2
53-93		<112059	~112335	>112339		253	HXT Y
5	GHz	1125.0	1125.0	~1125.0		1.41	
2.7	GHz	1119.0	1124.0	1133.0		1.91	
610	MHz	1118.0	1119.0	1142.0	[P0.6]	2.30	
410	MHz	1119.0	1120.0	1122.0		1.87	
8.8	GHz	1124.0	1134.0	1142.0		1.63	
5	GHz	1122.0	1134.0	1142.0		1.72	
1.4	GHz	1119.0	1134.0	1141.0		1.88	
DS II	25-160	1125		1144	FS	2	
DS DCIM	800-2000	1117		1204	GG	2	
DS DCIM	2000-4500	1118		1153	GG	1	

15.4	GHz	1133.0	1135.0	1158.0	[U8.8 / 15.4]	1.82	
8.8	GHz	1133.0	1146.0	1158.0		1.91	
5	GHz	1133.0	1146.0	1158.0		1.81	
2.7	GHz	1133.0	1134.0	1158.0		1.73	
610	MHz	1133.0	1134.0	1158.0		1.94	
410	MHz	1133.0	1134.0	1158.0		1.85	
245	MHz	1133.0	1134.0	1158.0		2.15	
DS IV	35-85	1137		1751		1	
DS III	200-260	1136		1136	G	2	
DS CONT	135-270	~1133		~1138		2	
CME	WL	1154	1230 km/s	-12.4 km/s ²	229°	275°	



Particle event: To($E_p > 10$ MeV) – 28d02^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 28d06^h$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 5 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 28d12^h$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 13 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm1} = 140 \text{ MeV}$

– $E_{qm2} = 105 \text{ MeV}$


Sources: ☉ solar flare 27d23^h37^m, M1.2/SF, N11W78, AR9090

Main X-ray burst 1-8 Å: onset – 27d23^h37^m, max – 27d23^h42^m, $\Phi = 0.0044 \text{ J/m}^2$



CME: 28d00^h54^m, $V = 0447 \text{ km/s}$, $\Delta\phi = 057^\circ$; $dA = 296^\circ$

▲ SC 28d06^h34^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 July 28



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 July 28

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES-10						
EPS	>5	02 ^h	06 ^h /12 ^h	10.7/36	3d	
EPS	>10	02 ^h	06 ^h /12 ^h	5/13	3d	
EPS	>30	02 ^h	04 ^h /11 ^h	0.5/0.6	2d	
EPS	>50	02 ^h	04 ^h /11 ^h	0.14/0.12	1d	
EPS	>60	02 ^h	04 ^h /11 ^h -	0.1/0.08	1d	
EPS	>100	02 ^h	04 ^h /11 ^h -	0.04/0.03	1d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	04 ^h /12 ^h	4.2 / 8.2	2d	
MEPED	>36	-	04 ^h /12 ^h	0.5 / 0.4	1d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	<3 ^h	04 ^h /11 ^h	14/48	2d	
CPME	>4	<3 ^h	04 ^h /11 ^h	9/21.7	2d	
CPME	>10	<3 ^h	04 ^h /11 ^h	4/8.2	2d	
CPME	>30	<3 ^h	04 ^h /11 ^h	0.35/0.3	1d	
CPME	>60	<3 ^h	04 ^h /11 ^h	0.16/0.05	1d	
ACE						
SIS	>10	2 ^h	03 ^h /11 ^h	2.1/7.5	1d	
SIS	>30	2 ^h	03 ^h /10 ^h	0.55/0.35	1d	

SOHO						
EPHIN (INT)	>50	00 ^h	04 ^h /12 ^h	0.03/0.02	1d	

Differential fluxes of protons for the event of 2000 July 28

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	01 ^h	06 ^h /12 ^h	3.8/15	1d	
CPME	2-4.6	01 ^h	04 ^h /12 ^h	1.96/8.5	1d	
CPME	4.6-15	01 ^h	04 ^h /11 ^h	0.54/1.56	1d	
CPME	15-25	01 ^h	04 ^h /11 ^h	0.2/0.39	1d	
CPME	25-48	01 ^h	04 ^h /11 ^h	0.016/0.02	1d	
CPME	48-96	00 ^h	04 ^h /11 ^h	0.002/0.0016	1d	
CPME	96-145	00 ^h	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	01 ^h	04 ^h /12 ^h	0.38/13.2	4d	
LION	2-6	01 ^h	04 ^h /12 ^h	0.16/3.1	4d	
EPHIN	4-8	01 ^h	04 ^h /12 ^h	0.4/10	4d	
EPHIN	8-25	01 ^h	04 ^h /12 ^h	0.11/1.5	4d	
EPHIN	25-41	01 ^h	04 ^h /12 ^h	0.011/0.04	4d	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 July 28

2000 July 27

☉

AR9090

To event 375

H α	6563 Å	<2340	~2341	>0046	N11W78	SF	FU
1 -12.5	keV	2337	2342	2347		M1.2	4.4E-3
53-93	keV	<233922	234114	234406		237	HXT Y
5.7	GHz	2340.8	2342.3	2344.4		1.18	
2.8	GHz	2335.0	2341.0	2347.0		0.85	
CME	WL	0054	0447 km/s	-1.8 km/s ²	057°	296°	

Particle event: To($E_p > 10$ MeV) – 11d15^h

Tmax($E_p > 10$ MeV) – 11d17^h, Jmax ($E_p > 10$ MeV) – $3.2 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 1 day


Quasimaximal energy of protons in the event – $E_{qm} = 75$ MeV

Sources: \square back side solar flare event < 11d16^h54^m, AR unknown, behind E_L ;



CME: 11d16^h54^m, $V = 0300$ km/s, $\Delta\phi = 271^\circ$, $dA = 035^\circ$

\blacktriangle SC 11d18^h46^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 August 11



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 August 11

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	15 ^h	17 ^h	19.6	1d	
EPS	>10	15 ^h	17 ^h	3.2	1d	
EPS	>30	15 ^h	17 ^h	0.07	1d	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	15 ^h	16 ^h	7340	1d	
MEPED	>0.8	15 ^h	16 ^h	5740	1d	
MEPED	>2.5	15 ^h	16 ^h	820	1d	
MEPED	>6.9	15 ^h	16 ^h	20	1d	
MEPED	>16	15 ^h	16 ^h	1.5	1d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	15 ^h	17 ^h	2100	1 d	
CPME	>4	15 ^h	17 ^h	170	1 d	
CPME	>10	15 ^h	17 ^h	6.6	1 d	
CPME	>30	15 ^h	16 ^h	0.17	1 d	
CPME	>60	15 ^h	16 ^h	0.08	1 d	

ACE						
SIS	>10	14 ^h	16 ^h	3.1	1 d	
SIS	>30	14 ^h	17 ^h	0.07	1 d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 2000 August 11

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	10 ^h	17 ^h	1390	1d	
CPME	2-4.6	10 ^h	17 ^h	240	1d	
CPME	4.6-15	10 ^h	17 ^h	11.8	1d	
CPME	15-25	10 ^h	16 ^h	0.12	1d	
CPME	25-48	10 ^h	16 ^h	0.0015	0,5d	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	10d23 ^h	16 ^h	193	1d	
LION	2-6	10d23 ^h	16 ^h	24.8	1d	
EPHIN	4-8	10d22 ^h	16 ^h	24,7	~1,5d	
EPHIN	8-25	10d22 ^h	16 ^h	0,75	~1,5d	
EPHIN	25-41	10 ^h	15 ^h	0,0025	~1,5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Wang R., 2009

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 August 11

2000	August 11	<input type="checkbox"/>	AR	To event 376			
CME	WL	1654	0300 km/s	6.6 km/s ²	035°	273°	

Particle event: To($E_p > 10$ MeV) – 13d01^h

Tmax($E_p > 10$ MeV) – 13d06^h, Jmax ($E_p > 10$ MeV) – $1.2 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 1 day


Quasimaximal energy of protons in the event – $E_{qm} = 70$ MeV

Sources: • solar flare 12d09^h56^m, M1.1/SN, S16W79, AR9119



Main X-ray burst 1-8 Å : onset – 12d09^h45^m, max – 12d09^h56^m, $\Phi = 0.011 \text{ J/m}^2$

CME: 12d10^h35^m, $V = 0662 \text{ km/s}$, $\Delta\phi = 168^\circ$; $dA = 262^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 August 13



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 August 13

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	02 ^h	06 ^h	25	1d	
EPS	>10	01 ^h	06 ^h	1.2	1d	
EPS	>30	-	04 ^h	0.04	-	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100				-	
POES-16						
MEPED	>0.24	01 ^h	06 ^h	1514	1d	
MEPED	>0.8	01 ^h	06 ^h	1154	1d	
MEPED	>2.5	01 ^h	06 ^h	512	1d	
MEPED	>6.9	01 ^h	06 ^h	44	1d	
MEPED	>16	-	-	-	-	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	01 ^h	06 ^h	634	1d	
CPME	>4	01 ^h	06 ^h	78	1d	
CPME	>10	01 ^h	06 ^h	2.2	1d	
CPME	>30	01 ^h	05 ^h	0.04	1d	
CPME	>60	-	-	-	-	
ACE						
SIS	>10	01 ^h	04 ^h	1.8	<1d	
SIS	>30	-	05 ^h	0.12	-	
SOHO						
EPHIN (INT)	>50	-	-	-	-	


Differential fluxes of protons for the event of 2000 August 13

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	01 ^h	06 ^h	327	7d	
CPME	2-4.6	01 ^h	06 ^h	96.4	6d	
CPME	4.6-15	01 ^h	06 ^h	5.5	5d	
CPME	15-25	01 ^h	06 ^h	0.012	1d	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	01 ^h	05 ^h	25.9	6d	
LION	2-6	01 ^h	05 ^h	5.8	5d	
EPHIN	4-8	01 ^h	05 ^h	7	5d	
EPHIN	8-25	01 ^h	05 ^h	0.2	4d	
EPHIN	25-41	-	-	-	-	
EPHIN	41-53	-	-	-	-	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 August 13

2000		August 12		•	AR9119		To event 377
H α	6563 Å	0951	0954	1015	S16W79	SN	AL
1 -12.5	keV	0945	0956	1009		M1.1	1.1E-2
50 -150	keV	0948	0949	0957		351	HXT Y

8.8	GHz	0948.0	0951.0	0955.0		1.72	
5	GHz	0947.0	0949.0	0955.0	[U0.4 P5]	1.81	
2.8	GHz	0947.0	1002.7	1016.0		1.75	
610	MHz	0947.0	0951.0	0955.0		1.70	
410	MHz	0955.0	0956.0	0956.0		1.65	
245	MHz	0954.0	1003.0	1004.0		1.77	
DS III	220-500	0956		0956	GG	2	
DS DCIM	2000-4500	0948		0955	GW	1	
DS DCIM	800-1672	0949		0953		2	
DS DCIM	2000-4500	1001		1006	G	1	
DS DCIM	800-2000	1001		1006		1	
CME	WL	1035	0662 km/s	-6.7 km/s ²	168°	262°	



Particle event: To($E_p > 10$ MeV) – 12d14^h

Tmax($E_p > 10$ MeV) – 13d02^h, Jmax ($E_p > 10$ MeV) – 180 /cm².s.sr

Duration of the event – 6 days

Quasimaximal energy of protons in the event – $E_{qm} = 350$ MeV


Sources: • solar flare 12d11^h22^m, 2F/M1.0, S19W08, AR9163

Main X-ray burst 1-8 Å: onset – 12d11^h31^m, max – 12d12^h00^m, $\Phi = 0.045$ J/m²



CME: 12d11^h54^m, $V = 1550$ km/s, $\Delta\phi = 360^\circ$, $dA = 220^\circ$

▲ SC15d04^h50^m, ▲ SC15d19^h12^m, ▲ SC18d14^h44^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 September 12



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 September 12

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES-10						
EPS	>5	14 ^h	13d03 ^h	670	6d	
EPS	>10	14 ^h	13d02 ^h	180	6d	
EPS	>30	14 ^h	13d02 ^h	8.4	5d	
EPS	>50	14 ^h	13d02 ^h	1.6	5d	
EPS	>60	14 ^h	13d00 ^h	0.97	5d	
EPS	>100	14 ^h	21 ^h	0.27	3d	
POES-16						
MEPED	>0.24	-	13d00 ^h	2961	6d	
MEPED	>0.8	-	13d00 ^h	2501	6d	
MEPED	>2.5	-	13d00 ^h	1424	6d	
MEPED	>6.9	-	13d00 ^h	381	6d	
MEPED	>16	-	13d00 ^h	31.5	5d	
MEPED	>36	-	13d00 ^h	4.54	5d	
MEPED	>70	-	13d00 ^h	0.63	5d	
MEPED	>140	-	13d00 ^h	0.11	3d	
IMP-8						
CPME	>1	14 ^h	13d03 ^h	2610	7d	
CPME	>4	14 ^h	13d03 ^h	920	7d	
CPME	>10	14 ^h	13d03 ^h	128	6d	
CPME	>30	14 ^h	13d03 ^h	6.6	6d	
CPME	>60	14 ^h	13d03 ^h	2.7	5d	
ACE						
SIS	>10	14 ^h	13d03 ^h	188	3d	
SIS	>30	14 ^h	13d03 ^h	10	2d	
SOHO						
EPHIN (INT)	>50	14 ^h	13d01 ^h	1.2	3d	

BALLOONS						
Mu	> 94		13d(08 ^h -08 ^h)	1.01		
Mu	>116		13d(08 ^h -08 ^h)	0.36		
Mu	>140		13d(08 ^h -08 ^h)	0.11		

Differential fluxes of protons for the event of 2000 September 12

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	15 ^h	13d04 ^h	810	7d	
CPME	2-4.6	15 ^h	13d03 ^h	445	7d	
CPME	4.6-15	14 ^h	13d03 ^h	57	6d	
CPME	15-25	14 ^h	13d03 ^h	4.4	6d	
CPME	25-48	14 ^h	13d02 ^h	0.22	5d	
CPME	48-96	13 ^h	13d02 ^h	0.028	5d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	15 ^h	13d04 ^h	412	7d	
LION	2-6	15 ^h	13d04 ^h	109	7d	
EPHIN	4-8	14 ^h	13d04 ^h	296	7d	
EPHIN	8-25	14 ^h	13d04 ^h	26.8	6d	
EPHIN	25-41	14 ^h	13d04 ^h	0.35	6d	

References:

Nitta N.V., E.W. Cliver, A.J. Tylka et al., 2003.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 September 12

2000 September 12




•

AR9163

To event 378

H α	6563 Å	1122	1200	1458	S19W08	2F	U
DSF	6563 Å	~1050		1230	S12W18	23°	
1 -12.5	keV	1131	1213	1313		M1.0	4.5E-2
3	GHz	1122.8	1129.2	>1255.8		1.75	
410	MHz	1130.0	1133.0	1137.0		1.79	
245	MHz	1128.0	1129.0	1129.0		1.92	
DS II	25-180	1133		1149	HARM,FS	2	
DS I	80-270	1151		~1231	N	1	
DS III	160-270	~1137		1144	S	1	
DS III	25-250	1152		1211	GG,DS	2	
15.4	GHz	1152.0	1215.0	1309.0		1.66	
8.8	GHz	1154.0	1222.0	1327.0		1.65	
5	GHz	1154.0	1222.0	1341.0		1.68	
2.7	GHz	1210.0	1231.0	0000.0		1.67	
410	MHz	1215.0	1215.0	1220.0		2.46	
DS III		1216		1224	25-40	1	

5	GHz	1210.0	1257.0	1345.0		1.80	
2.7	GHz	1210.0	1257.0	0000.0		1.72	
410	MHz	1300.0	1300.0	1302.0		1.98	
CME		1154	1550 km/s	58.2 km/s ²	360°	220°	



Particle event: To($E_p > 10$ MeV) – 16d08^h

$T_{\max 1}(E_p > 10 \text{ MeV})$ – 16d11^h, $J_{\max 1}(E_p > 10 \text{ MeV})$ – $3.5 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV})$ – 16d17^h, $J_{\max 2}(E_p > 10 \text{ MeV})$ – $9.8 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$ *)

Duration of the event – 4 days

Quasimaximal energy of protons in the event – $E_{qm1} = 310 \text{ MeV}$

– $E_{qm2} = 140 \text{ MeV}$

*) The data from IMP-8


Sources: □ solar flare 16d06^h40^m, M2.5/ -, n05w90*, AR9182

Main X-ray burst 1-8 Å onset – 16d06^h40^m, max – 16d07^h28^m, $\Phi = 0.16 \text{ J/m}^2$



CME: 16d07^h27^m, $V = 1336 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 270^\circ$

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 October 16



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 October 16

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	08 ^h	11 ^h / -	4.8/ -	4d	
EPS	>10	08 ^h	11 ^h / -	3.5/ -	4d	
EPS	>30	08 ^h	11 ^h / -	1.8/ -	3d	
EPS	>50	08 ^h	11 ^h / -	0.8/ -	3d	
EPS	>60	08 ^h	11 ^h / -	0.5/ -	2d	
EPS	>100	08 ^h	11 ^h / -	0.16/ -	1.5d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	12 ^h / -	56/ -	4d	
MEPED	>6.9	-	12 ^h / -	25/ -	4d	
MEPED	>16	-	12 ^h / -	9.1/ -	3d	
MEPED	>36	-	12 ^h / -	2.5/ -	3d	
MEPED	>70	-	12 ^h / -	0.42/ -	2d	
MEPED	>140	-	-	-	-	

IMP-8						
CPME	>1	08 ^h	11 ^h /17 ^h	23.2/61	6 d	
CPME	>4	08 ^h	11 ^h /18 ^h	15.2/25	5 d	
CPME	>10	08 ^h	11 ^h /17 ^h	8.5/9.8	4 d	
CPME	>30	08 ^h	12 ^h /17 ^h	2.2/1.4	2 d	
CPME	>60	08 ^h	12 ^h /17 ^h	0.75/0.4	1 d	
ACE						
SIS	>10	08 ^h	10 ^h /16 ^h	13.7/14.1	3 d	
SIS	>30	08 ^h	10 ^h / -	4.9/ -	2 d	
SOHO						
EPHIN (INT)	>50	08 ^h	10 ^h / -	0.34/ -	1d	

Differential fluxes of protons for the event of 2000 October 16

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	08 ^h	13 ^h /16 ^h	3.3/16.2	6d	
CPME	2-4.6	08 ^h	11 ^h /16 ^h	2.6/9.4	6d	
CPME	4.6-15	08 ^h	11 ^h /16 ^h	0.66/1.6	6d	
CPME	15-25	08 ^h	11 ^h /16 ^h	0.4/0.43	4d	
CPME	25-48	08 ^h	11 ^h /16 ^h	0.08/0.06	3d	
CPME	48-96	08 ^h	11 ^h /16 ^h	0.014/0.01	2d	
CPME	96-145	08 ^h	-	-	-	
CPME	145-440	08 ^h	-	-	-	
SOHO						
LION	0,75-2	08 ^h	11 ^h /16 ^h	1.3/4.8	7d	
LION	2-6	08 ^h	11 ^h /16 ^h	0.3/1.1	7d	
EPHIN	4-8	08 ^h	11 ^h /16 ^h	0.78/3.2	7d	
EPHIN	8-25	08 ^h	11 ^h /16 ^h	0.22/0.7	6d	
EPHIN	25-41	08 ^h	11 ^h /16 ^h	0.053/0.06	6d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 October 16

2000 October 16 ☐ AR9182 To event 379

Hα		No Data			n05w90*		
1 -12.5	keV	0640	0728	0911		M2.5	1.6E-1
53 – 93	keV	0640	0709	0730		8	HXT Y

9.1	GHz	0658.5	0739.0	0839.0		1.20	
5	GHz	0728.0	0728.0	0729.0		1.41	
3	GHz	0722.9	0728.8	0732.0	U0.9 P3	1.64	
1.4	GHz	0728.0	0730.0	0732.0		1.32	
900	MHz	0718.0	0728.2	0909.0		1.23	
600	MHz	0722.5	0728.0	0744.3		1.94	
DS II	20-60	0705		0718	UE	2	
DS IV	25-146	0704		0820		1	
DS III	45-95	0718		~0916	N	1	
DS CONT	800-2000	0722		0741		1	
CME	WL	0727	1366 km/s	9.9 km/s ²	360°	2270°	

* – probable localization of the flare event



Particle event: To($E_p > 10$ MeV) – 25d13^h

Tmax($E_p > 10$ MeV) – 25d23^h, Jmax ($E_p > 10$ MeV) – 4.1 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm} = 95$ MeV


Sources: ☐ solar flare 25d08^h45^m, C4.0/..., ... W90, AR– unknown

Main X-ray burst 1-8 Å : onset – 25d08^h45^m, max – 25d11^h25^m, $\Phi = 0.065$ J/m²



CME: 25d08^h26^m, $V = 0770$ km/s, $\Delta\phi = 360^\circ$, $dA = 275^\circ$

▲ SC 28d09^h54^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 October 25



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 October 25

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Duration	Comments
GOES-10						
EPS	>5	13 ^h	23 ^h	6.5	3d	
EPS	>10	13 ^h	23 ^h	4.1	3d	
EPS	>30	13 ^h	23 ^h	0.42	2d	
EPS	>50	13 ^h	23 ^h	0.12	2d	
EPS	>60	-	24 ^h	0.09	1.5d	
EPS	>100	-	24 ^h	0.01	-	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	19 ^h	2.52	3d	
MEPED	>36	-	19 ^h	0.6	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	13 ^h	23 ^h	30	3d	
CPME	>4	13 ^h	23 ^h	12	3d	
CPME	>10	13 ^h	22 ^h	5	3d	
CPME	>30	13 ^h	21 ^h	0.8	1d	
CPME	>60	13 ^h	21 ^h	0.14	1d	
ACE						
SIS	>10	12 ^h	20 ^h	6.8	2 d	
SIS	>30	12 ^h	19 ^h	0.8	5 h	

SOHO						
EPHIN (INT)	>50	12 ^h	20 ^h	0.025	0.5d	

Differential fluxes of protons for the event of 2000 October 25

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	13 ^h	26d02 ^h	19	3d	
CPME	2-4.6	13 ^h	26d02 ^h	8.5	3d	
CPME	4.6-15	13 ^h	22 ^h	0.8	3d	
CPME	15-25	13 ^h	22 ^h	0.24	3d	
CPME	25-48	13 ^h	21 ^h	0.026	1.5d	
CPME	48-96	13 ^h	19 ^h	0.002	1.5d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	12 ^h	20 ^h	3.6	3.5d	
LION	2-6	12 ^h	19 ^h	1.1	3.5d	
EPHIN	4-8	12 ^h	20 ^h	2.3	3.5d	
EPHIN	8-25	12 ^h	20 ^h	0.47	3.5d	
EPHIN	25-41	12 ^h	20 ^h	0.031	3.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2000 October 25**

2000	October 25	☐			AR	To event 380	
Hα	6563 Å	No Flare					
1 -12.5	keV	0845	1125	1521		C4.0	6.5E-2
CME	WL	0826	0770 km/s	17.4km/s ²	360°	275°	

Particle event: To(Ep>10 MeV) – 31d07^h

Tmax(Ep>10 MeV) – 01d03^h, Jmax (Ep>10 MeV) – 2.1/cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – E_{qm} = 70 MeV


Sources: o solar flare 31d02^h51^m, C6.0/1N, S20E08, AR9209

Main X-ray burst 1-8 Å: onset – 31d02^h51^m, max – 31d03^h00^m, $\Phi = 0.0036$ J/m²



CME: 31d03^h50^m, V = 0074 km/s, $\Delta\phi = 189^\circ$, dA = 113°

▲ SC 31d17^h14^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 October 31



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 October 31


S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES 10						
EPS	>5	07 ^h	01d03 ^h	11.5	3d	
EPS	>10	07 ^h	01d03 ^h	2.1	3d	
EPS	>30	07 ^h	01d01 ^h	0.07	2d	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
IMP-8						
CPME	>1	03 ^h	01d02 ^h	420	3d	
CPME	>4	03 ^h	01d02 ^h	28	3d	
CPME	>10	03 ^h	01d02 ^h	2.6	2d	
CPME	>30	03 ^h	01d02 ^h	0.02	-	
CPME	>60	-	-	-	-	
ACE						
SIS	>10	03 ^h	01d02 ^h	2.8	2d	
SIS	>30	-	--	-	-	
SOHO						
EPHIN (INT)	>50	-	21 ^h	0.006	-	

Differential fluxes of protons for the event of 2000 October 31

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	06 ^h	01d02 ^h	315	3d	
CPME	2-4.6	06 ^h	01d02 ^h	43.8	3d	
CPME	4.6-15	06 ^h	01d02 ^h	1.9	3d	
CPME	15-25	06 ^h	01d02 ^h	0.06	3d	
CPME	25-48	06 ^h	01d02 ^h	0.0018	2d	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	05 ^h	01d02 ^h	28	3d	
LION	2-6	05 ^h	01d02 ^h	3.7	3d	
EPHIN	4-8	05 ^h	01d02 ^h	4.7	3d	
EPHIN	8-25	05 ^h	01d02 ^h	0.22	3d	
EPHIN	25-41	05 ^h	01d02 ^h	0.002	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 October 31

2000 October 31		o			AR9209	To event 381	
H α		0253	0300	0320	S20E08	1N	EF
1 -12.5	keV	0251	0300	0307		C6.0	3.6E-03
5.7	GHz	0252.7	0300.0	0321.5		1.45	
2.8	GHz	0251.0	0254.8	0306.0		1.28	
610	MHz	0259.0	0259.0	~0259.0		2.43	
CME	WL	0350	0074km/s	2.5km/s ²	189°	113°	



Particle event: To($E_p > 10$ MeV) – 08d23^h

Tmax($E_p > 10$ MeV) – 09d15^h, Jmax ($E_p > 10$ MeV) – $9.7 \cdot 10^3 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 8 days

Quasimaximal energy of protons in the event – $E_{qm} = 650$ MeV


Sources: ● solar flare 08d22^h40^m, 3F/M7.4, N20W66, AR9213

Main X-ray burst 1-8 Å: onset – 08d22^h42^m, max – 08d23^h28^m, $\Phi = 0.21 \text{ J/m}^2$



CME: 08d23^h06^m, $V = 1738 \text{ km/s}$, $\Delta\varphi = 170^\circ$, $dA = 299^\circ$

▲ SC 10d06^h28^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 November 08



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 November 08

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	23 ^h	09d15 ^h	12524	8d	
EPS	>10	23 ^h	09d15 ^h	9740	6d	
EPS	>30	23 ^h	09d11 ^h	3113	6d	
EPS	>50	23 ^h	09d04 ^h	1239	6d	
EPS	>60	23 ^h	09d04 ^h	877	5d	
EPS	>100	23 ^h	09d04 ^h	306	5d	
POES-16						
MEPED	>0.24	-	09d04 ^h	23230	8d	
MEPED	>0.8	-	09d04 ^h	19670	8d	
MEPED	>2.5	-	09d04 ^h	15740	6d	
MEPED	>6.9	-	09d04 ^h	13670	6d	
MEPED	>16	-	09d04 ^h	7000	8d	
MEPED	>36	-	09d04 ^h	3884	6d	
MEPED	>70	-	09d04 ^h	625	5d	
MEPED	>140	-	09d04 ^h	50	5d	
IMP-8						
CPME	>1	23 ^h	09d09 ^h	10500	8 d	
CPME	>4	23 ^h	09d08 ^h	9320	8 d	
CPME	>10	23 ^h	09d08 ^h	7310	8 d	
CPME	>30	23 ^h	09d04 ^h	3220	6 d	
CPME	>60	23 ^h	09d04 ^h	1200	6 d	
ACE						
SIS	>10	23 ^h	09d06 ^h	4570	6d	
SIS	>30	23 ^h	09d05 ^h	1980	6d	

SOHO						
EPHIN (INT)	>50	23 ^h	09d04 ^h	2080	5d	
BALLOONS						
Mi-1	>167		09d09 ^h	43.2		
Mi-1	>259		09d09 ^h	4.0		
Mi-1	>356		09d09 ^h	0.21		

Differential fluxes of protons for the event of 2000 November 08

S/c, instruments	ΔE , MeV	To	Tmax	J_{\max} /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	23 ^h	09d03 ^h	200	8d	
CPME	2-4.6	23 ^h	09d03 ^h	260	8d	
CPME	4.6-15	23 ^h	09d03 ^h	167	8d	
CPME	15-25	23 ^h	09d03 ^h	200	8d	
CPME	25-48	23 ^h	09d03 ^h	79	8d	
CPME	48-96	23 ^h	09d03 ^h	32	7d	
CPME	96-145	23 ^h	09d03 ^h	10	6d	
CPME	145-440	23 ^h	09d00 ^h	1.1	5d	
SOHO						
LION	0,75-2	23 ^h	09d07 ^h	2320	8d	
LION	2-6	23 ^h	09d07 ^h	320	8d	
EPHIN	4-8	23 ^h	09d06 ^h	85	8d	
EPHIN	8-25	23 ^h	09d06 ^h	55	8d	
EPHIN	25-41	23 ^h	09d05 ^h	35	6d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

El-Borie M.A., 2003a.
El-Borie M.A., 2003b.
Struminsky A.B., 2003.
Nitta N.V., E.W. Cliver, A.J. Tylka, and P. Smit, 2003.
Al-Thoyaib S.S., 2005.
Rawat R., S. Alex, and G.S. Lakhina., 2006.
Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Lario D., R.B. Decker, and A. Aran., 2008.
Lario D., A. Aran, R.B. Decker, 2009.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 November 08

2000	November 08	•	AR9213	To event 382			
H α	6563 Å	2240	2323	0030	N20W66	3F	F
1 -12.5	keV	2242	2328	0005		M7.4	2.1E-1
50 – 150	keV	<2316	~2316	2355		236	HXT Y

15.4	GHz	2306.0	2343.0	0012.0		2.20	
8.8	GHz	2305.0	2318.0	0041.0		2.38	
5	GHz	2300.0	2318.0	0041.0		2.64	
2.7	GHz	2258.0	2318.0	0052.0	U0.6 P2.7	2.87	
1.4	GHz	2258.0	2318.0	0027.0		2.62	
610	MHz	2326.0	2326.0	~2326.0		1.32	
245	MHz	2304.0	2305.0	2305.0		1.60	
DS IV	25-180	2251		2330		1	
DS IV	500-2000	2306		2345		1	
DS III	25-400	2249		2324	G	2	
DS III	25-106	2330		0002	N	1	
DS UNCLF	100-200	2316		2322		2	
CME	WL	2306	1738 km/s	69.9km/s ²	170°	299°	



Particle event: To($E_p > 10$ MeV) – 24d07^h

Tmax($E_p > 10$ MeV) – 24d21^h, Jmax ($E_p > 10$ MeV) – 65 /cm².s.sr

Duration of the event – 1.5 days

Quasimaximal energy of protons in the event – $E_{qm} = 460$ MeV


Sources: • solar flare 24d04^h55^m, X2.0/3B, N20W05, AR9236

Ø solar flare 24d14^h51^m, X2.3/2B, N20W08, AR9236



Main X-ray burst 1-8 Å: onset – 24d04^h55^m, max – 24d05^h02^m, $\Phi = 0.083$ J/m²

CME: 24d05^h30^m, $V = 1298$ km/s, $\Delta\phi = 360^\circ$, $dA = 313^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 November 24



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 November 24

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	07 ^h	21 ^h	153	1.5d	
EPS	>10	07 ^h	21 ^h	65	1.5d	
EPS	>30	07 ^h	18 ^h	11	1.5d	
EPS	>50	07 ^h	18 ^h	4.2	1.5d	
EPS	>60	07 ^h	17 ^h	3	1.5d	
EPS	>100	07 ^h	17 ^h	1.1	1.5d	
POES-16						
MEPED	>0.24	-	24 ^h	1360	1.5d	
MEPED	>0.8	-	24 ^h	1100	1.5d	
MEPED	>2.5	-	24 ^h	660	1.5d	
MEPED	>6.9	-	24 ^h	170	1.5d	
MEPED	>16	-	18 ^h	46	1.5d	
MEPED	>36	-	18 ^h	14.3	1.5d	
MEPED	>70	-	18 ^h	2.05	1.5d	
MEPED	>140	-	18 ^h	0.56	1.5d	
IMP-8						
CPME	>1	06 ^h	21 ^h	748	1.5 d	
CPME	>4	06 ^h	21 ^h	255	1.5 d	
CPME	>10	06 ^h	21 ^h	66	1.5 d	
CPME	>30	06 ^h	20 ^h	8.7	1.5 d	
CPME	>60	06 ^h	18 ^h	4.2	1.5 d	
ACE						
SIS	>10	06 ^h	20 ^h	83	1.5 d	
SIS	>30	06 ^h	18 ^h	18	1.5 d	

SOHO						
EPHIN (INT)	>50	07 ^h	18 ^h	3	1.5d	

Differential fluxes of protons for the event of 2000 November 24

S/c, instruments	ΔE , MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	07 ^h	21 ^h	243	1.5d	
CPME	2-4.6	07 ^h	21 ^h	115	1.5d	
CPME	4.6-15	07 ^h	21 ^h	17.9	1.5d	
CPME	15-25	07 ^h	19 ^h	2.6	1,5d	
CPME	25-48	07 ^h	19 ^h	0.33	1.5d	
CPME	48-96	07 ^h	18 ^h	0.08	1.5d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	07 ^h	21 ^h	78	1.5d	
LION	2-6	07 ^h	21 ^h	20.6	1.5d	
EPHIN	4-8	07 ^h	21 ^h	41.3	1.5d	
EPHIN	8-25	07 ^h	21 ^h	4.1	1.5d	
EPHIN	25-41	07 ^h	18 ^h	0.34	1.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 November 24

2000 November 24			•		AR9236	To event 383	
H α	6563 Å	0457	0501	0535	N20W05	3B	EF
1 -12.5	keV	0455	0502	0508		X2.0	8.3E-2
15.4	GHz	0458.0	0500.0	0516.0	U5 / 15.4	3.74	
8.8	GHz	0458.0	0500.0	0520.0		3.54	
5	GHz	0458.0	0501.0	0512.0		3.11	
2.7	GHz	0458.0	0459.0	0507.0		3.34	
1.4	GHz	0458.0	0459.0	0507.0	[U0.6 P1.4]	3.96	


610	MHz	0458.0	0501.0	0510.0		3.18	
410	MHz	0458.0	0459.0	0506.0		3.71	
245	MHz	0458.0	0501.0	0506.0		3.86	
200	MHz	0459.0	0459.0	0514.0		3.46	
DS II	25-180	0502		0529		3	
DS III	18-900	0459		0502	G	3	
DS III	18-80	0510		0516	G	3	
CME	WL	0530	1298 km/s	2.1km/s ²	360°	313°	



2000 November 24 Ø AR9213 To event 383

H α	6563 Å	1454	1510	1601	N20W08	2B	FH
1 -12.5	keV	1451	1513	1521		X2.3	1.6E-1
53 – 93	keV	1454	1511	1536		1498	HXT Y
1.2 – 5.6	MeV	1454	1508	1536		195	GRS Y
15.4	GHz	1500.0	1508.0	1612.0	U2.7 / 15.4	4.15	
8.8	GHz	1454.0	1509.0	1612.0		3.86	
5	GHz	1453.0	1508.0	1609.0		3.43	
2.7	GHz	1455.0	1508.0	1612.0		3.15	
1.4	GHz	1454.0	1508.0	1525.0		3.56	
610	MHz	1454.0	1508.0	1612.0		4.54	
410	MHz	1453.0	1509.0	1612.0		3.79	
245	MHz	1453.0	1507.0	1612.0		3.58	
DS II	30-80	1512		1523		3	
DS III	30-80	1457		1523	N	3	

^o n						Bolivia	
CME	WL	1530	1454 km/s	-3.3km/s ²	360°	324°	



Particle event: To($E_p > 10$ MeV) – 26d03^h

Tmax($E_p > 10$ MeV) – 26d20^h, Jmax ($E_p > 10$ MeV) – 670 /cm².s.sr

Duration of the event – 4 days

Quasimaximal energy of protons in the event – $E_{qm} = 400$ MeV

Sources: ● solar flare 25d00^h59^m, M8.2/2N, N07E50, AR9240


○ solar flare 25d18^h33^m, X1.9/2B, N19W24, AR9236

Main X-ray burst 1-8 Å: onset – 25d00^h59^m, max – 25d01^h31^m, $\Phi = 0.21$ J/m²



CME: 26d01^h31^m, $V = 2519$ km/s, $\Delta\phi = 360^\circ$, $dA = 082^\circ$

▲ SC 26d07^h58^m, ▲ SC 26d11^h58^m, ▲ SC 28d05^h31^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2000 November 26



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2000 November 26

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES 10						
EPS	>5	03 ^h	20 ^h	1705	5d	
EPS	>10	03 ^h	20 ^h	670	4d	
EPS	>30	03 ^h	20 ^h	64.5	3d	
EPS	>50	03 ^h	20 ^h	12.6	2d	
EPS	>60	03 ^h	20 ^h	5.6	2d	
EPS	>100	03 ^h	20 ^h	0.95	1.5d	
POES 16						
MEPED	>0.24	-	17 ^h	5910	5d	
MEPED	>0.8	-	17 ^h	3820	5d	
MEPED	>2.5	-	17 ^h	2145	5d	
MEPED	>6.9	-	17 ^h	900	4d	
MEPED	>16	-	17 ^h	179.5	3d	
MEPED	>36	-	17 ^h	38	2d	
MEPED	>70	-	17 ^h	2.25	2d	
MEPED	>140	-	17 ^h	0.23	1.5d	
IMP-8						
CPME	>1	02 ^h	20 ^h	7600	5d	
CPME	>4	02 ^h	20 ^h	1900	5d	
CPME	>10	02 ^h	20 ^h	568	4d	
CPME	>30	02 ^h	20 ^h	70	4d	
CPME	>60	02 ^h	20 ^h	34	3d	
ACE						
SIS	>10	02 ^h	20 ^h	532	4d	
SIS	>30	02 ^h	20 ^h	68.5	3d	

SOHO						
EPHIN (INT)	>50	04 ^h	21 ^h	6.5	3d	


Differential fluxes of protons for the event of 2000 November 26

S/c, instruments	ΔE, MeV	To	Tmax	Jmax /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	01 ^h	20 ^h	3350	7d	
CPME	2-4.6	01 ^h	20 ^h	1110	7d	
CPME	4.6-15	01 ^h	20 ^h	124	6d	
CPME	15-25	01 ^h	20 ^h	24	5d	
CPME	25-48	01 ^h	20 ^h	2.2	4d	
CPME	48-96	01 ^h	20 ^h	0.24	4d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0,75-2	04 ^h	21 ^h	345	7d	
LION	2-6	04 ^h	21 ^h	70	7d	
EPHIN	4-8	04 ^h	21 ^h	193	7d	
EPHIN	8-25	04 ^h	21 ^h	23.7	6d	
EPHIN	25-41	04 ^h	21 ^h	1.1	6d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2000 November 26

2000	November 25	•	AR9240	To event 384			
Hα	6563 Å	0100	0106	0324	N07E50	2N	FU
1 -12.5	keV	0059	0131	0201		M8.2	2.1E-1
53 – 93	keV	0056	0122	0133		81	HXT Y
1.2 – 5.6	MeV	005535	012226	013239		1911	GRS Y
410	MHz	0113.0	0113.0	0125.0		2.73	
245	MHz	0107.0	0108.0	0225.0		2.94	
200	MHz	0108.0	0114.0	0130.0		2.56	
DS II	25-146	0107		0137		3	
DS IV	25-180	0107		1031		2	
DS IV	400-2000	0118		0259		2	
DS III	25-180	0104		0215	N	3	
15.4	GHz	0104.0	0122.0	0201.0		3.23	
8.8	GHz	0100.0	0122.0	0235.0		3.45	
5	GHz	0058.0	0122.0	0249.0		3.70	
2.8	GHz	0057.0	0131.0	0247.0	P2.8	4.05	
1.4	GHz	0055.0	0142.0	0245.0		3.90	



610	MHz	0103.0	0145.0	0239.0		3.30	
500	MHz	0059.0	0146.0	0221.0		2.92	
410	MHz	0113.0	0146.0	0223.0		2.81	
CME	WL	0131	2519 km/s	-5.0km/s ²	360°	082°	



2000 November 25 Ø AR9236 To event 384

H α	6563 Å	1836	1841	2016	N19W24	2B	EFU
1 -12.5	keV	1833	1844	1855		X1.9	1.5E-1
53 – 93	keV	1836	1842	1948		170	HXT Y
1.2 – 5.6	MeV	1836	1838	1948		137	GRS Y
15.4	GHz	1833.0	1838.0	1841.0	U1.4 / 15.4	3.73	
8.8	GHz	1835.0	1838.0	1907.0		3.58	
5	GHz	1833.0	1842.0	1843.0		3.30	
2.7	GHz	1834.0	1838.0	1841.0		2.88	
1.4	GHz	1833.0	1838.0	1841.0		2.81	
610	MHz	1836.0	1838.0	1907.0		3.26	
410	MHz	1835.0	1838.0	1841.0		4.23	
245	MHz	1833.0	1838.0	1841.0		4.23	
DS II	25-180	1839		1849		2	
DS II	30-80	1845		1857		3	
DS III	25-180	1838		1842		2	
2.7	GHz	1836.0	1904.0	1907.0		3.04	
1.4	GHz	1902.0	1903.0	1904.0		3.11	
610	MHz	1902.0	1904.0	1904.0		2.82	
410	MHz	1902.0	1903.0	1904.0		3.46	
245	MHz	1902.0	1902.0	1905.0		3.00	

15.4	GHz	1939.0	1941.0	1943.0		2.00	
8.8	GHz	1938.0	1941.0	1947.0		2.36	
5	GHz	1938.0	1941.0	1945.0		2.61	
2.7	GHz	1937.0	1941.0	1944.0		3.36	
1.4	GHz	1927.0	1939.0	1947.0	P0.4 - 1.4	4.00	
610	MHz	1932.0	1941.0	1947.0		3.98	
410	MHz	1938.0	1941.0	1945.0		3.98	
245	MHz	1938.0	1939.0	1942.0		3.11	
DS III	20-160	<2000		2210	S	1	
CME	WL	1932	1932 km/s	-10.8 km/s ²	360°	348°	



Events in 2001

			Page
1. Event 2001.01.28 – (2001-028)	№ 385	271
2. Event 2001.02.26 – (2001-057)	№386	276
3. Event 2001.03.26 – (2001- 085)	№ 387	280
4. Event 2001.03.29 – (2001-088)	№ 388	285
5. Event 2001.04.02 – (2001-092)	№ 389	290
6. Event 2001.04.09 – (2001-099)	№ 390	295
7. Event 2001.04.10 – (2001-100)	№ 391	300
8. Event 2001.04.12– (2001-102)	№ 392	307
9. Event 2001.04.15 – (2001-105) – GLE-60	№ 393	313
10. Event 2001.04.18 – (2001-108) ? GLE-61	№ 394	318
11. Event 2001.04.27 – (2001-117)	№ 395	323
12. Event 2001.05.07 – (2001-127)	№ 396	328
13. Event 2001.05.20 – (2001-140)	№ 397	332
14. Event 2001.06.15 – (2001-166)	№ 398	337
15. Event 2001.08.09 – (2001-221)	№ 399	342
16. Event 2001.08.16 – (2001-228)	№ 400	346
17. Event 2001.09.15 – (2001-258)	№ 401	350
18. Event 2001.09.24 – (2001-267)	№ 402	355
19. Event 2001.10.01 – (2001-274)	№ 403	361
20. Event 2001.10.19 – (2001-292)	№ 404	366
21. Event 2001.10.22 – (2001-295)	№ 405	371
22. Event 2001.10.28 – (2001-301)	№ 406	377
23. Event 2001.11.04 – (2001-308) – GLE-62	№ 407	381
24. Event 2001.11.17 – (2001-321)	№ 408	387
25. Event 2001.11.22 – (2001-326)	№ 409	391
26. Event 2001.12.26 – (2001-360) – GLE-63	№ 410	397
27. Event 2001.12.29 – (2001-363)	№ 411	402
28. Event 2001.12.30 – (2001-364)	№ 412	408

Particle event: To($E_p > 10$ MeV) – 28d18^h

Tmax($E_p > 10$ MeV) – 29d01^h, Jmax ($E_p > 10$ MeV) – 29 /cm².s.sr

Duration of the event – 3 days


Quasimaximal energy of protons in the event – $E_{qm} = 325$ MeV

Sources: • solar flare 28d15^h08^m, 1N/ M1.5, S04W59, AR9313



Main X-ray burst 1-8 Å: 28d15^h40^m, max – 28d16^h40^m, $\Phi = 0.03$ J/m²

CME: 28d15^h54^m, $V = 0916$ km/s, $\Delta\phi = 360^\circ$; $dA = 254^\circ$;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 January 28



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 January 28

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	18 ^h	29d02 ^h	45.3	3d	
EPS	>10	18 ^h	29d01 ^h	29.0	3d	
EPS	>30	18 ^h	29d01 ^h	4.2	2d	
EPS	>50	18 ^h	21 ^h	1.3	2d	
EPS	>60	18 ^h	21 ^h	0.84	2d	
EPS	>100	18 ^h	20 ^h	0.27	2d	
POES-16						
MEPED	>0.24	-	29d11 ^h	3805	3d	
MEPED	>0.8	-	29d11 ^h	3090	3d	
MEPED	>2.5	-	29d11 ^h	970	3d	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	20 ^h	8.3	2d	
MEPED	>36	-	20 ^h	4.8	2d	
MEPED	>70	-	20 ^h	0.7	2d	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	23 ^h	29d16 ^h	3610	4 d	
CPME	>4	17 ^h	29d07 ^h	295	3 d	
CPME	>10	17 ^h	29d06 ^h	34.3	3 d	
CPME	>30	17 ^h	29d02 ^h	3.1	2 d	
CPME	>60	17 ^h	29d02 ^h	1	2 d	

ACE						
SIS	>10	16 ^h	29d06 ^h	43	3 d	
SIS	>30	16 ^h	29d02 ^h	5.6	2 d	
SOHO						
EPHIN (INT)	>50	17 ^h	29d02 ^h	0,3	2d	


Differential fluxes of protons for the event of 2001 January 28

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	29d03 ^h	29d21 ^h	2520	6d	
CPME	2-4.6	29d00 ^h	29d15 ^h	550	6d	
CPME	4.6-15	19 ^h	29d13 ^h	23,3	5d	
CPME	15-25	17 ^h	29d07 ^h	1,2	4d	
CPME	25-48	17 ^h	29d06 ^h	0,11	4d	
CPME	48-96	17 ^h	29d02 ^h	0,02	3d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	19 ^h	29d16 ^h	440	5d	
LION	2-6	19 ^h	29d13 ^h	72	5d	
EPHIN	4-8	19 ^h	29d12 ^h	118	5d	
EPHIN	8-25	18 ^h	29d10 ^h	3.6	5d	
EPHIN	25-41	17 ^h	29d02 ^h	0.145	5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 January 28

2001	January 28	•	AR9313	To event 385			
H α	6563 Å	1508	1546	1730	S04 W59	1N	FH
1 – 12	keV	1540	1600	1624		M1.5	3.0E-2

15.4	GHz	1551.0	1604.0	1607.0	P2.8 [U1.4 P2.7]	1.62	
8.8	GHz	1548.0	1602.0	1607.0	U6.7/15.4	1.59	
6.7	GHz	1527.0	1553.0	2112.0		1.46	
5	GHz	1548.0	1552.0	1607.0		1.57	
2.7	GHz	1542.0	1552.0	1607.0		2.11	
1.4	GHz	1542.0	1553.0	1607.0		1.98	
610	MHz	1534.0	1537.0	1607.0		3.38	
410	MHz	1534.0	1539.0	1607.0		3.61	
280	MHz	1536.2	1543.2	1615.5		3.97	
235	MHz	1536.2	1541.5	1615.5		3.07	
DS DCIM	2000-4000	1522		1527	C	1	
CME	WL	1554	0916km/s	3.5 km/s ²	360°	254°	



Particle event: To($E_p > 10$ MeV) – 26d09^h

Tmax($E_p > 10$ MeV) – 26d20^h, Jmax ($E_p > 10$ MeV) – 1 /cm².s.sr *)

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm} = 65$ MeV

*) The data from all s/c


Sources: ☐ solar flare 26d05^h14^m, C1.6/..., s04w90*, AR9354

Main X-ray burst 1-8 Å: onset – 26d05^h14^m, max – 26d07^h41^m, $\Phi = 0.0042$ J/m²



CME: 26d05^h30^m, $V = 0851$ km/s, $\Delta\phi = 152^\circ$, $dA = 263^\circ$;

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 February 26



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 February 26

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	10 ^h	21 ^h	3.8	1.5d	
EPS	>10	10 ^h	21 ^h	0.6	1.5d	
EPS	>30	10 ^h	21 ^h	0.03	1d	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	-	22 ^h	930	2.5 d	
MEPED	>0.8	-	22 ^h	334	2.5 d	
MEPED	>2.5	-	22 ^h	45	2.5 d	
MEPED	>6.9	-	22 ^h	5	2 d	
MEPED	>16	-	22 ^h	0.4	2 d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	9 ^h	20 ^h	170	2.5 d	
CPME	>4	9 ^h	20 ^h	11.5	2 d	
CPME	>10	9 ^h	20 ^h	1.05	2 d	
CPME	>30	-	15 ^h	0.05	-	
CPME	>60	-	-	-	-	

ACE						
SIS	>10	8 ^h	17 ^h	1.45	1.5 d	
SIS	>30	8 ^h	17 ^h	0.15	1 d	
SOHO						
EPHIN (INT)	>50	09 ^h	15 ^h	0.01	0.3d	

Differential fluxes of protons for the event of 2001 February 26

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	10 ^h	20 ^h	118	5d	
CPME	2-4.6	10 ^h	20 ^h	18.7	5d	
CPME	4.6-15	10 ^h	19 ^h	0.77	2d	
CPME	15-25	10 ^h	18 ^h	0.018	1d	
CPME	25-48	-	21 ^h	0.003	-	
CPME	48-96	-	21 ^h	0.0004	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	09 ^h	20 ^h	24.2	2.5d	
LION	2-6	09 ^h	20 ^h	2.75	2.5d	
EPHIN	4-8	09 ^h	19 ^h	3.2	7d	
EPHIN	8-25	09 ^h	17 ^h	0.13	5d	
EPHIN	25-41	09 ^h	16 ^h	0.0008	2.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 February 26

2001 February 26 ☐ AR9354 To event 386

H α		No Flare			s04w90*		
1 – 12	keV	0514	0741	0749		C1.6	4.2E-03
CME	WL	0530	0851 km/s	8.1km/s ²	152°	263°	

* – probable localization of the flare event

Particle event: To($E_p > 10$ MeV) – 26d20h

Tmax($E_p > 10$ MeV) – 27d08h, Jmax ($E_p > 10$ MeV) – 1.8 /cm².s.sr

Duration of the event – 1.5 days

Quasimaximal energy of protons in the event – $E_{qm} = 55$ MeV


Sources: ☉ solar flare 26d13h03m, M2.2/1F, N15E27, AR9393

Main X-ray burst 1-8 Å: onset – 26d13h03m, max – 26d13h26m, $\Phi = 0.037$ J/m²;



CME: 26d13h50m, $V = 0541$ km/s, $\Delta\phi = 055^\circ$, $dA = 104^\circ$

▲ SC 27d17^h47^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 March 26



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 March 26

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	20 ^h	08 ^h	22.2	2.5d	
EPS	>10	20 ^h	08 ^h	1.8	2.5d	
EPS	>30	21 ^h	09 ^h	0.07	2d	
EPS	>50	-	09 ^h	0.02	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	-	18 ^h	$2.5 \cdot 10^4$	3 d	
MEPED	>0.8	-	18 ^h	$6.2 \cdot 10^3$	3 d	
MEPED	>2.5	-	18 ^h	355	3 d	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	-	-	-	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	01 ^h	12 ^h	1160	3 d	
CPME	>4	01 ^h	12 ^h	66	3 d	
CPME	>10	01 ^h	11 ^h	2.2	3 d	
CPME	>30	-	11 ^h	0.03	-	
CPME	>60	-	-	-	-	

ACE						
SIS	>10	16 ^h	10 ^h -	1.4	1.5d	
SIS	>30		10 ^h	0.1	1d	
SOHO						
EPHIN (INT)	>50	23 ^h	17 ^h	0.04 *)	-	*) The increase on Forbush


Differential fluxes of protons for the event of 2001 March 26

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	13 ^h	12 ^h	867	3,5d	
CPME	2-4.6	12 ^h	12 ^h	112	3d	
CPME	4.6-15	16 ^h	12 ^h	3.2	3d	
CPME	15-25	08 ^h	11 ^h	0.029	2,5d	
CPME	25-48	-	-	-	-	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	17 ^h	17 ^h	60	2d	
LION	2-6	17 ^h	17 ^h	5.6	2d	
EPHIN	4-8	17 ^h	17 ^h	10.3	2d	
EPHIN	8-25	17 ^h	17 ^h	0.21	2d	
EPHIN	25-41	17 ^h	16 ^h	3.6·10 ⁻⁴	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 March 26

2001	March 26	☉	AR9393	To event 387			
Hα	6563Å	1306	1321	1423	N15E27	1F	F
1 – 12	keV	1303	1326	1344		M2.2	3.7E-2
53 – 93	keV	<131310	~131622	>132018		8	HXT Y

15.4	GHz	1309.0	1327.0	1408.0		1.74	
8.8	GHz	1309.0	1326.0	1408.0		1.80	
5	GHz	1309.0	1314.0	1359.0		1.76	
2.7	GHz	1309.0	1313.0	1351.0		1.62	
1.4	GHz	1320.0	1320.0	1322.0		1.97	
610	MHz	1312.0	1323.0	1328.0		1.88	
410	MHz	1312.0	1314.0	1328.0		1.57	
245	MHz	1307.0	1307.0	1308.0		2.28	
DS DCIM	800-2000	1310		1326	G	2	
DS DCIM	1415-4000	1312		1322	C	3	
CME	WL	1350	0541km/s	4.8 km/s ²	055°		



Particle event: To($E_p > 10$ MeV) – 29d13^h

$T_{\max 1}(E_p > 10 \text{ MeV})$ – 29d19^h, $J_{\max 1}(E_p > 10 \text{ MeV})$ – 7 /cm².s.sr

$T_{\max 2}(E_p > 10 \text{ MeV})$ – 31d00^h, $J_{\max 2}(E_p > 10 \text{ MeV})$ – 22 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm1} = 215$ MeV

– $E_{qm2} = 115$ MeV


Sources: • solar flare 29d09^h55^m, 1N/X1.7, N16W12, AR9393

Main X-ray burst 1-8 Å: onset – 29d09^h55^m, max – 29d10^h15^m, $\Phi = 0.22$ J/m²



CME: 29d10^h26^m, $V = 0942$ km/s, $\Delta\phi = 360^\circ$, $dA = 071^\circ$

▲ SC 31d00^h52^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 March 29



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 March 29

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	13 ^h	19 ^h /31d00 ^h	9.4/106.5	3d	
EPS	>10	13 ^h	19 ^h /31d00 ^h	7/22	3d	
EPS	>30	13 ^h	19 ^h /31d00 ^h	1.9 /0.85	2d	
EPS	>50	13 ^h	18 ^h /31d00 ^h	0.59/0.45	2d	
EPS	>60	13 ^h	18 ^h /31d00 ^h	0.31/0.08	2d	
EPS	>100	13 ^h	17 ^h /31d00 ^h	0.06/0.01	2d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	18 ^h /31d00 ^h	10.2/6.5	2d	
MEPED	>36	-	18 ^h /31d00 ^h	3.5/0.4	2d	
MEPED	>70	-	18 ^h / -	0.44/ -	2d	
MEPED	>140	-	18 ^h / -	0.06/ -	2d	

IMP-8						
CPME	>1	12 ^h	23 ^h /31d01 ^h	91/3500	2.5 d	
CPME	>4	11 ^h	23 ^h /31d00 ^h	38.7/173	2.5 d	
CPME	>10	11 ^h	21 ^h /30d23 ^h	14.5/20.5	2.5 d	
CPME	>30	11 ^h	17 ^h /30d23 ^h	3.2/0.85	2.5 d	
CPME	>60	11 ^h	17 ^h /30d23 ^h	1.9/0.37	2.5 d	
ACE						
SIS	>10	11 ^h	19 ^h /30d23 ^h	15.5/22.2	2.5 d	
SIS	>30	11 ^h	19 ^h /30d23 ^h	3.5/0.8	2.5 d	
SOHO						
EPHIN (INT)	>50	13 ^h	17 ^h /31d00 ^h	0.26/0.29	1.5d	


Differential fluxes of protons for the event of 2001 March 29

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	17 ^h	30d01 ^h /31d00 ^h	44/2720	3d	
CPME	2-4.6	15 ^h	30d01 ^h /31d00 ^h	18.4/308	3d	
CPME	4.6-15	13 ^h	30d01 ^h /31d00 ^h	3.35/13.5	3d	
CPME	15-25	12 ^h	23 ^h /31d00 ^h	0.88/0.61	2d	
CPME	25-48	11 ^h	20 ^h /31d00 ^h	0.09/0.033	2d	
CPME	48-96	11 ^h	17 ^h /31d00 ^h	0.013/0.003	2d	
CPME	96-145	11 ^h	-	-	-	
CPME	145-440	11 ^h	-	-	-	
SOHO						
LION	0.75-2	16 ^h	30d01 ^h /31d00 ^h	4.8/46.2	3d	
LION	2-6	16 ^h	30d01 ^h /31d00 ^h	1.3/14.4	3d	
EPHIN	4-8	13 ^h	30d01 ^h /31d00 ^h	6/46.4	3d	
EPHIN	8-25	13 ^h	21 ^h /31d00 ^h	0.76/2.4	3d	
EPHIN	25-41	11 ^h	21 ^h /31d00 ^h	0.072/0.025	1.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 March 29

2001	March 29	•	AR9393	To event 388			
H α	6563 Å	0955	1005	1108	N16 W12	2N	EFZ
1 – 12	keV	0957	1015	1032		X1.7	2.2E-1

15.4	GHz	0959.0	1010.0	1019.0		3.88	
8.8	GHz	0958.0	1011.0	1019.0	P0.6 U1.4 P 8.8	4.11	
5	GHz	0958.0	1011.0	1020.0		3.98	
2.7	GHz	0958.0	1011.0	1019.0		3.67	
1.4	GHz	1004.0	~1009.0	1029.0		3.59	
900	MHz	0957.0	1010.4	1021.0		3.81	
610	MHz	1004.0	~1011.0	1033.0		4.95	
245	MHz	1000.0	1011.0	1020.0		4.71	
204	MHz	1003.0	1010.8	1013.6		4.37	
DS II	HARM	1003		~1007	HARM	2	
DS II	UE	~1004		~1010	UE	3	
DS IV		0958		~1038		3	
DS I	GG,DC	1000		1001	GG,DC	2	
DS III	G	1006		1012	G	2	
DS III	GG,C	1008		1012	GG,C	2	
DS III	GG,RS	1012		1014	GG,RS	2	
DS III	G,C	1025		1025	G,C	2	
DS CONT		~1005		>1158		2	
DS DCIM	GG,FS	0956		1034	GG,FS	3	
DS DCIM	GG,FS	0956		1023	GG,FS	3	
n°							Swiss(A,T)
CME	WL	1026	0942 km/s	3.5 km/s ²	360°	071°	



Particle event: To(Ep>10 MeV) – 02d23^h

Tmax(Ep>10 MeV) – 03d07^h, Jmax (Ep>10 MeV) – 112 /cm².s.sr

Duration of the event – 5 days

Quasimaximal energy of protons in the event – E_{qm} = 575 MeV

Sources: ■ solar flare 02d21^h32^m, X>17.5/..., n19w90*, AR9393


Main X-ray burst 1-8 Å: onset – 02^d 21^h 32^m, max – 02d21^h51^m, Φ = 1.5 J/m²

CME: 02d22^h06^m, V = 2505 km/s, Δφ = 244°, dA = 293°



▲ SC 04d14^h55^m

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 April 02



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 April 02

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	23 ^h	03d07 ^h	140	5d	
EPS	>10	23 ^h	03d07 ^h	112	5d	
EPS	>30	23 ^h	03d07 ^h	51	4d	
EPS	>50	23 ^h	03d07 ^h	11.5	3d	
EPS	>60	23 ^h	03d06 ^h	13	2d	
EPS	>100	23 ^h	03d06 ^h	4.4	2d	
POES-16						
MEPED	>0.24	-	03d03 ^h	2035	5d	
MEPED	>0.8	-	03d03 ^h	1670	5d	
MEPED	>2.5	-	03d03 ^h	1330	5d	
MEPED	>6.9	-	03d03 ^h	560	5d	
MEPED	>16	-	03d03 ^h	282	4d	
MEPED	>36	-	03d03 ^h	200	3d	
MEPED	>70	-	03d03 ^h	12.4	2d	
MEPED	>140	-	03d03 ^h	1.4	2d	
IMP-8						
CPME	>1	23 ^h	03d07 ^h	1370	5d	
CPME	>4	23 ^h	03d07 ^h	815	5d	
CPME	>10	23 ^h	03d07 ^h	452	5d	
CPME	>30	23 ^h	03d07 ^h	95	5d	
CPME	>60	23 ^h	03d07 ^h	34.6	5d	

ACE						
SIS	>10	22 ^h	03d06 ^h	845	5d	
SIS	>30	22 ^h	03d06 ^h	237	5d	
SOHO						
EPHIN (INT)	>50	23 ^h	03d06 ^h	29	3d	
BALLOONS						
Mi	>125		03d(09 ^h 30 ^m -10 ^h 12 ^m)	2.5		
Mi	>168		03d(09 ^h 30 ^m -10 ^h 12 ^m)	0.8		
Mi	>320			0.1		

Differential fluxes of protons for the event of 2001 April 02

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	19 ^h	03d06 ^h	236	□4d	
CPME	2-4.6	18 ^h	03d07 ^h	144	□4d	
CPME	4.6-15	23 ^h	03d07 ^h	40.3	□4d	
CPME	15-25	23 ^h	03d07 ^h	21.9	□4d	
CPME	25-48	23 ^h	03d07 ^h	3.4	□4d	
CPME	48-96	23 ^h	03d07 ^h	0.59	□4d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	23 ^h	03d07 ^h	43.3	7d	
LION	2-6	23 ^h	03d07 ^h	30.5	7d	
EPHIN	4-8	23 ^h	03d07 ^h	120	7d	
EPHIN	8-25	23 ^h	03d07 ^h	34	7d	
EPHIN	25-41	23 ^h	03d07 ^h	3.9	7d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 April 02

2001 April 02




AR9393

To event 389

Hα	6563 Å	No Flare Patrol			n19w90*		
1 – 12	keV	2132	2151	2203		X>17.5	1.5E00
53 – 93	keV	212902	213634	>213750		55	HXT Y
53 – 93	keV	<221334	~221334	>005640		11	HXT Y

15.4	GHz	2132.0	2149.0	2316.0	P0.2 U0.5/15.4	4.15	15.4
8.8	GHz	2135.0	2147.0	2321.0		3.88	8.8
5	GHz	2135.0	2147.0	2210.0		3.74	5
2.7	GHz	2143.0	2148.0	2212.0		3.56	2.7
1.4	GHz	2144.0	2148.0	2323.0		3.64	1.4
610	MHz	2146.0	2150.0	2205.0		2.97	610
500	MHz	2146.0	2150.0	2203.0		2.71	500
410	MHz	2148.0	2149.0	2201.0		2.85	410
245	MHz	2146.0	2152.0	2214.0		4.65	245
200	MHz	2149.0	2152.0	2207.0		3.65	200
DS II	25-280	2152		2157		3	DS II
DS III	25-180	2135		0156	N	1	DS III
DS III	70-120	2149		2150	B	2	DS III
n°							Armenia(S)
CME	WL	2206	2505 km/s	108.5 km/s ²	244°	293°	

* – probable localization of the flare event



Particle event: To($E_p > 10$ MeV) – 09d17^h

Tmax($E_p > 10$ MeV) – 09d20^h, Jmax ($E_p > 10$ MeV) – 2.2 /cm².s.sr

Duration of the event – 0.6 days


Quasimaximal energy of protons in the event – $E_{qm} = 390$ MeV

Sources: • solar flare 09d15^h20^m, M7.9/1B, S21W04, AR9415



Main X-ray burst 1-8 Å: onset – 09d15^h20^m, max – 09d15^h34^m, $\Phi = 0.13$ J/m²

CME: 09d15^h54^m, $V = 1192$ km/s, $\Delta\phi = 360^\circ$, $dA = 211^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 April 09



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 April 09

S/c, instruments	Ep. MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura-tion	Comments
GOES-10						
EPS	>5	17 ^h	20 ^h	2.9	0.6d	
EPS	>10	17 ^h	20 ^h	2.2	0.6d	
EPS	>30	17 ^h	20 ^h	1.6	0.6d	
EPS	>50	17 ^h	20 ^h	1.0	0.6d	
EPS	>60	17 ^h	20 ^h	0.8	0.6d	
EPS	>100	17 ^h	20 ^h	0.4	0.6d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	20 ^h	4.1	0.6d	
MEPED	>36	-	20 ^h	2.8	0.6d	
MEPED	>70	-	20 ^h	0.7	0.6d	
MEPED	>140	-	20 ^h	0.4	0.6d	
IMP-8						
CPME	>1	17 ^h	20 ^h	139	0.6d	
CPME	>4	17 ^h	20 ^h	8.8	0.6d	
CPME	>10	17 ^h	20 ^h	4.3	0.6d	
CPME	>30	17 ^h	20 ^h	1.6	0.6d	
CPME	>60	17 ^h	20 ^h	1.2	0.6d	

ACE						
SIS	>10	17 ^h	23 ^h	6.8	0.5d	
SIS	>30	17 ^h	22 ^h	4.3	0.5d	
SOHO						
EPHIN (INT)	>50	16 ^h	20 ^h	0.45	0.5d	

Differential fluxes of protons for the event of 2001 April 09

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura-tion	Comments
IMP-8						
CPME	1-2	-	10d01 ^h	148	0.5d	
CPME	2-4.6	-	10d01 ^h	11.5	0.5d	
CPME	4.6-15	-	10d01 ^h	0.42	0.5d	
CPME	15-25	-	23 ^h	0.155	0.5d	
CPME	25-48	-	23 ^h	0.037	0.5d	
CPME	48-96	-	23 ^h	0.0127	0.5d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	-	21 ^h	4.2	0.5d	
LION	2-6	-	22 ^h	0.47	0.5d	
EPHIN	4-8	19 ^h	23 ^h	1	0.5d	
EPHIN	8-25	18 ^h	23 ^h	0.18	0.5d	
EPHIN	25-41	17 ^h	23 ^h	0.034	0.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 April 09

2001	April 09	•	AR9415	To event 390			
H α	6563Å	1524	1534	1712	S21W04	1B	EFH
1 – 12	keV	1520	1534	1600		M7.9	1.3E-1
53 – 93	keV	152259	~152459	>152513		25	HXT Y

15.4	GHz	1522.0	1526.0	1614.0	P0.6 U2.7/15.4	3.77	
8.8	GHz	1522.0	1530.0	1609.0		3.73	
6.7	GHz	1522.8	1536.2	1603.8		3.75	
5	GHz	1522.0	1535.0	1614.0		3.58	
2.7	GHz	1523.0	1537.0	1614.0		3.20	
1.4	GHz	1524.0	1526.0	1614.0		3.65	
610	MHz	1524.0	1530.0	1614.0		4.59	
245	MHz	1524.0	1529.0	1614.0		3.83	
DS II	25-180	1528		1549		3	
DS IV	40-800	1523		~1606	P	3	
DS III	25-180	1527		1550	N	2	
DS DCIM	2000-4500	1522		1605	GG	3	
DS DCIM	800-2000	1524		1609	GG,SP,FS	3	
CME	WL	1554	1192km/s	108.5km/s ²	360°	211°	



Particle event: To($E_p > 10$ MeV) – 10d08^h

$T_{\max 1}(E_p > 10 \text{ MeV})$ – 11d01^h, $J_{\max 1}(E_p > 10 \text{ MeV})$ – 50 /cm².s.sr

$T_{\max 2}(E_p > 10 \text{ MeV})$ – 11d20^h, $J_{\max 2}(E_p > 10 \text{ MeV})$ – 280 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm1} = 350$ MeV

– $E_{qm2} = 260$ MeV


Sources: • solar flare 10d04^h59^m, 3N/X2.3, S23W09, AR9415

Ø solar flare 11d12^h56^m, M2.3/1F, S20W28 AR9415



Main X-ray burst 1-8 Å: onset – 10d05^h06^m, max – 10d05^h26^m, $\Phi = 0.3 \text{ J/m}^2$

CME 10d05^h30^m, $V = 2411 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 166^\circ$

ΔSC 11d13^h43^m, ΔSC 11d15^h19^m




Time profiles of the proton fluxes for the event of 2001 April 10



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 April 10

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	08 ^h	11d01 ^h /11d20 ^h	109/1256	2d	
EPS	>10	08 ^h	11d01 ^h /11d20 ^h	50/280	2d	
EPS	>30	08 ^h	11d01 ^h /11d20 ^h	9.6/11.6	2d	
EPS	>50	08 ^h	21 ^h /11d20 ^h	2.5/ 1.9	2d	
EPS	>60	08 ^h	19 ^h /11d18 ^h	1.1/0.47	2d	
EPS	>100	08 ^h	17 ^h /11d17 ^h	0.28/0.08	2d	
POES-16						
MEPED	>0.24	08 ^h	11d01 ^h /11d20 ^h	5500/20230	2d	
MEPED	>0.8	08 ^h	11d01 ^h /11d20 ^h	3085/10330	2d	
MEPED	>2.5	08 ^h	11d01 ^h /11d20 ^h	1485/3340	2d	
MEPED	>6.9	09 ^h	11d01 ^h /11d20 ^h	285/540	2d	
MEPED	>16	09 ^h	11d01 ^h /11d20 ^h	55/50	2d	
MEPED	>36	08 ^h	11d01 ^h /11d20 ^h	11/8.05	2d	
MEPED	>70	08 ^h	11d01 ^h /11d20 ^h	0.6/1.1	2d	
MEPED	>140	08 ^h	11d01 ^h /11d20 ^h	0.09/0.12	2d	

IMP-8						
CPME	>1	09 ^h	11d01 ^h /11d20 ^h	3230/9760	2d	
CPME	>4	08 ^h	11d00 ^h /11d20 ^h	450/1060	2d	
CPME	>10	08 ^h	11d00 ^h /11d20 ^h	86/172	2d	
CPME	>30	08 ^h	20 ^h /11d20 ^h	8.4/7.5	2d	
CPME	>60	08 ^h	16 ^h /11d20 ^h	2.7/2.5	2d	
ACE						
SIS	>10	08 ^h	11d01 ^h /11d20 ^h	92/195	2d	
SIS	>30	08 ^h	20 ^h /11d20 ^h	15/9.2	2d	
SOHO						
EPHIN (INT)	>50	08 ^h	16 ^h /11d18 ^h	0.7/0.8	2d	

Differential fluxes of protons for the event of 2001 April 10




S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	11 ^h	11d02 ^h /11d18 ^h	1680/8070	2d	
CPME	2-4.6	09 ^h	11d01 ^h /11d20 ^h	536/1200	2d	
CPME	4.6-15	08 ^h	11d01 ^h /11d20 ^h	30.8/80	2d	
CPME	15-25	08 ^h	11d01 ^h /11d20 ^h	3.3/6.1	2d	
CPME	25-48	08 ^h	11d01 ^h /11d20 ^h	0.4/0.33	2d	
CPME	48-96	08 ^h	11d01 ^h /11d20 ^h	0.05/0.025	2d	
CPME	96-145	08 ^h	-	-	-	
CPME	145-440	08 ^h	-	-	-	
SOHO						
LION	0.75-2	10 ^h	21 ^h /11d21 ^h	69.6/285	2d	
LION	2-6	10 ^h	19 ^h /11d21 ^h	11.5/59.6	2d	
EPHIN	4-8	08 ^h	21 ^h /11d20 ^h	42.7/265	2d	
EPHIN	8-25	08 ^h	18 ^h /11d20 ^h	2.6/18.9	2d	
EPHIN	25-41	08 ^h	18 ^h /11d20 ^h	0.29/0.36	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2001 April 10**

2001	April 10	•	AR9415	To event 391			
H α	6563Å	0459	0520	0810	S23W09	3N	UZ
1 – 12	keV	0506	0526	0542		X2.3	3.0E-1
53 – 93	keV	<051343	~051907	055729		89	HXT Y
15.4	GHz	0510.0	0523.0	0651.0		3.70	
8.8	GHz	0509.0	0522.0	0707.0	P8.8	3.84	
5	GHz	0509.0	0523.0	0706.0		3.73	
3	GHz	0504.4	0525.8	0541.1		3.73	
500	MHz	0504.0	0539.0	0700.0		3.45	
200	MHz	0504.0	0526.0			2.66	
DS II	25-180	0518		0536		3	
DS IV	25-600	0515		0535		3	
DS IV	40-800	<0518		~1010		3	
DS III	25-180	0509		0535	N	3	
DS CONT	23-130	0515		0534		3	
9.1	GHz	0607.2	0608.5	0612.5		1.26	
3	GHz	0606.4	0608.8	0622.8		1.52	
900	MHz	<0504.2	0613.0	>0617.9		3.71	
610	MHz	0524.0	0612.0	0706.0		4.04	
410	MHz	0524.0	0612.0	0743.0	P0.4	4.74	
204	MHz	0606.0	0612.3	0614.2		3.62	
DS I	45-270	<0559		>1200	S,C	2	
DS III	45-270	<0559		0659	S	2	
DS III	25-180	0656		0658		3	
DS CONT	220-270	0605		0724		2	
DS DCIM	800-2000	0556		0619	GG,SP	3	
9.1	GHz	0656.9	0657.4	0657.9		1.18	
3	GHz	0634.5	0638.3	0642.0		1.83	
900	MHz	0645.1	0648.8			2.00	
245	MHz	0451.0	0648.0	0743.0	P0.2	4.00	
204	MHz	0629.4	0657.9	0707.1		3.63	
DS III		0656		0658	25-180	3	
DS DCIM	GG,SP	0623		0706	800-2000	1	
n°						Arm, Tib, Nor	
CME	WL	0530	2411km/s	211.6*km/s ²	360°	166°	



2001


April 11

Ø

AR9415

To event 391

H α	6563Å	1309	1321	1424	S20W28	1F	FU
1 – 12	keV	1256	1326	1349		M2.3	4.8E-2
53 – 93	keV	<132529	~132531	134203		9	HXT Y
15.4	GHz	1303.0	1317.0	1354.0		3.11	
9.5	GHz	1257.0	1317.4	1324.0	P0.4 U2.7 P9.5	3.43	
6.7	GHz	1310.0	1318.0	1330.8		3.32	
5	GHz	1259.0	1317.0	1340.0		3.20	
2.7	GHz	1300.0	1317.0	1404.0		2.73	
1.4	GHz	1300.0	1316.0	1402.0		2.87	
610	MHz	1305.0	1323.0	1354.0		3.89	
410	MHz	1259.0	1318.0	1411.0		4.08	
245	MHz	1300.0	1318.0	1354.0		3.49	
DS II	40-70	1303		1304	UE	3	
DS IV	40-800	1304		~1420	P	3	
DS III	25-180	1300		1321	N	3	
DS III	1415-3150	1303		1316	GG	3	
DS DCIM	2000-3768	1343		1401	GG	1	
CME	WL	1332	1103km/s	-13.0 km/s ²	360°	224°	



Particle event: To($E_p > 10$ MeV) – 12d12^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 12\text{d}17^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 4.3 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 13\text{d}10^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 8.7 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm1} = 410 \text{ MeV}$


– $E_{qm2} = 275 \text{ MeV}$

Sources: ● solar flare 12d09^h39^m, X2.0/2B, S20W42, AR9415



Main X-ray burst 1-8 Å: onset – 12d09^h39^m, max – 12d10^h28^m, $\Phi = 0.3 \text{ J/m}^2$

CME: 12d10^h31^m, $V = 1184 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 221^\circ$

▲ SC 13d07^h34^m




Time profiles of the proton fluxes for the event of 2001 April 12



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 April 12

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	-	17 ^h /13d10 ^h	5.8/26.2	2d	
EPS	>10	12 ^h	17 ^h /13d10 ^h	4.3/8.7	2d	
EPS	>30	12 ^h	17 ^h /13d10 ^h	4/1.1	2d	
EPS	>50	12 ^h	17 ^h /13d10 ^h	2.6/0.5	2d	
EPS	>60	12 ^h	17 ^h /13d08 ^h	2.1/0.3	2d	
EPS	>100	11 ^h	17 ^h /13d08 ^h	1.1/0.11	2d	
POES-16						
MEPED	>0.24	14 ^h	18 ^h / -	1470/-	1d	
MEPED	>0.8	14 ^h	18 ^h / -	850/-	1d	
MEPED	>2.5	14 ^h	18 ^h / -	345/-	1d	
MEPED	>6.9	14 ^h	18 ^h / -	90/-	-	
MEPED	>16	14 ^h	18 ^h / -	17.5/-	-	
MEPED	>36	14 ^h	18 ^h / -	7.8/-	-	
MEPED	>70	14 ^h	18 ^h / -	1.8/-	-	
MEPED	>140	14 ^h	18 ^h / -	0.83/-	-	
IMP-8						
CPME	>1	12 ^h	19 ^h /13d10 ^h	840/240	2d	
CPME	>4	12 ^h	17 ^h /13d10 ^h	110/30	2d	
CPME	>10	12 ^h	17 ^h /13d14 ^h	30/6.6	2d	
CPME	>30	12 ^h	17 ^h /13d14 ^h	7/1	2d	
CPME	>60	12 ^h	17 ^h /13d14 ^h	3.2/0.5	2d	

ACE						
SIS	>10	12 ^h	19 ^h / -	49/ -	2d	
SIS	>30	12 ^h	19 ^h / -	15/ -	2d	
SOHO						
EPHIN (INT)	>50	10 ^h	18 ^h / -	2.7/ -	1.5d	

Differential fluxes of protons for the event of 2001 April 12

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	10 ^h	19 ^h /13d14 ^h	610/156	3d	
CPME	2-4.6	10 ^h	19 ^h /13d14 ^h	94.5/26.8	3d	
CPME	4.6-15	10 ^h	19 ^h /13d14 ^h	6.5/2	3d	
CPME	15-25	10 ^h	19 ^h /13d14 ^h	1.33/0.22	3d	
CPME	25-48	10 ^h	19 ^h / -	0.24/ -	3d	
CPME	48-96	10 ^h	19 ^h / -	0.057/ -	2d	
CPME	96-145	10 ^h	19 ^h / -	0.052/ -	2d	
CPME	145-440	10 ^h	19 ^h / -	0.0036/ -	1d	
SOHO						
LION	0.75-2	10 ^h	20 ^h /13d12 ^h	28.7/10.8	3d	
LION	2-6	10 ^h	19 ^h /13d12 ^h	5/1.7	3d	
EPHIN	4-8	10 ^h	19 ^h /13d14 ^h	22.8/4.8	3d	
EPHIN	8-25	10 ^h	19 ^h /13d14 ^h	2.8/0.44	3d	
EPHIN	25-41	10 ^h	19 ^h / -	0.29/ -	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:



Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 April 12

2001	April 12	•	AR9415	To event 392			
H α	6563 Å	<1018	1040	1130	S23W42	2B	E
1 – 12	keV	0939	1028	1049		X2.0	3.0E-1
53 – 93	keV	101058	102042	115042		79	HXT Y
15.4	GHz	1014.0	1017.0	1050.0	U2.7/15.4	3.79	
9.1	GHz	1009.0	1020.5			3.68	
5	GHz	1014.0	1020.0	1052.0		3.41	
2.7	GHz	1014.0	1017.0	1052.0		3.08	
900	MHz	1013.6	1018.2	1110.0		3.38	
610	MHz	1014.0	1020.0	1100.0		4.30	
204	MHz	1014.8	1018.5	1022.1		4.80	

DS II	40-250	~1017		~1038	SH,H	3	
DS IV	40-800	1014		~1116		3	
DS I	25-270	1019		>1230	N,C	2	
DS III	1415-3200	1014		1038	GG,RS	2	
DS III	25-180	1017		1047	N	3	
DS DCIM	2000-4500	1014		1055	GG	3	
DS DCIM	800-2000	1014		1059	GG,FS	3	
3	GHz	1013.8	1032.3			2.60	
1.4	GHz	1014.0	1036.0	1052.0		3.15	
610	MHz	<1029.0	~1036.0	>1100.0		3.52	
410	MHz	1015.0	1036.0	1101.0	P0.4	3.91	
245	MHz	<1029.0	~1033.0	>1100.0		2.78	
245	MHz	1028.0	1033.0	0000.0		2.78	
DS II	135-210	1031		1034	HARM	2	
DS IV	25-180	1032		1341		1	
DS III	130-270	1027		1038	S	2	
15.4	GHz	<1046.0	~1101.0	>1102.0	0.9 \ 15.4	1.88	
8.8	GHz	<1029.0	~1047.0	>1102.0		2.38	
5	GHz	<1029.0	~1047.0	>1056.0		2.43	
3	GHz	1045.8	1047.3	1158.6		2.46	
900	MHz	1013.6	1046.5			3.99	
DS III	40-155	1046		1047	G	3	
DS CONT	30-80	1046		2233		2	
DS DCIM	800-2000	1102		1129	GG	2	
CME	WL	1031	1184 km/s	-20.0 km/s ²	360°	221°	





Particle event: To(Ep>10 MeV) – 15d14^h

Tmax(Ep>10 MeV) – 15d16^h, JmaxEp>10 MeV) – 270 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – E_{qm} = 3480 MeV


Sources: • solar flare 15d13^h19^m, X14.4/2B, S20W84, AR9415

Main X-ray burst 1-8 Å onset – 15d13^h19^m, max – 15d13^h50^m, Φ = 0.3 J/m²



CME: 15d14^h06^m, V=1199 km/s, Δφ=167°, dA= 268°

Δ SC 18d00^h46^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 April 15



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 April 15

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	14 ^h	16 ^h	275	2.5d	
EPS	>10	14 ^h	16 ^h	273	2.5d	
EPS	>30	14 ^h	16 ^h	271	2.5d	
EPS	>50	14 ^h	15 ^h	231	2.5d	
EPS	>60	14 ^h	15 ^h	210	2.5d	
EPS	>100	14 ^h	15 ^h	138	2.5d	
POES-16						
MEPED	>0.24	15 ^h	15 ^h	7750	> 2d	
MEPED	>0.8	15 ^h	15 ^h	3940	> 2d	
MEPED	>2.5	15 ^h	15 ^h	2410	> 2d	
MEPED	>6.9	15 ^h	15 ^h	1900	> 2d	
MEPED	>16	15 ^h	15 ^h	885	> 2d	
MEPED	>36	15 ^h	15 ^h	675	> 2d	
MEPED	>70	15 ^h	15 ^h	180	> 2d	
MEPED	>140	15 ^h	15 ^h	102	> 2d	

IMP-8						
CPME	>1	14 ^h	22 ^h	1650	2.5d	
CPME	>4	14 ^h	19 ^h	1030	2.5d	
CPME	>10	14 ^h	19 ^h	699	2.5d	
CPME	>30	14 ^h	17 ^h	334	2.5d	
CPME	>60	14 ^h	16 ^h	253	2.5d	
ACE						
SIS	>10	14 ^h	16 ^h	1680	2.5d	
SIS	>30	14 ^h	16 ^h	980	2.5d	
SOHO						
EPHIN (INT)	>50	13 ^h	16 ^h	247	2,5d	
BALLOONS						
Mi	>241		16d(07 ^h -08 ^h)	6.4		After
Mi	>279		16d(07 ^h -07 ^h)	3.5		maximum
Mi	>341		16d(07 ^h -07 ^h)	0.64		- “ -
NM						
Network	>433		15 ^h	29.8		
Network	>1000		15 ^h	0.814		
Network	>2000		15 ^h	0.0413		
Network	>3700		15 ^h	0.0029		
Network	>5800		15 ^h	0.00038		

Differential fluxes of protons for the event of 2001 April 15

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	16 ^h	23 ^h	262	2.5d	
CPME	2-4.6	14 ^h	23 ^h	214	2.5d	
CPME	4.6-15	14 ^h	23 ^h	47.2	2.5d	
CPME	15-25	14 ^h	20 ^h	25.4	2.5d	
CPME	25-48	14 ^h	19 ^h	6	2.5d	
CPME	48-96	14 ^h	17 ^h	2.4	2.5d	
CPME	96-145	14 ^h	17 ^h	1.3	2.5d	
CPME	145-440	14 ^h	16 ^h	0.4	2.5d	
SOHO						
LION	0.75-2	14 ^h	21 ^h	28.2	2.5d	
LION	2-6	14 ^h	20 ^h	16.5	2.5d	
EPHIN	4-8	14 ^h	20 ^h	90.8	2.5d	
EPHIN	8-25	14 ^h	20 ^h	27.2	2.5d	
EPHIN	25-41	14 ^h	19 ^h	4.1	2.5d	


References:

Quack M., M.-B. Kallenrode, M. von König et al., 2001
Perez-Peraza J., A. Gallegos-Cruz, E.V. Vashenyuk et al., 2003.
Struminsky A.B., 2003.
Bieber J.W., P. Evenson, W. Dröge et al., 2004.
Kuwabara T., Bieber J.W., Clem J., et.al., 2006.
Bombardieri D.J., M.L. Duldig, K.J. Michael et al., 2007.
Miroshnichenko L.I. and J. Perez-Peraza, 2008.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2001 April 15

2001 April 15 • AR9415 To event 393

H α	6563Å	1336	1349	1529	S20W84	2B	FHY
1 – 12	keV	1319	1350	1355		X14.4	3.0E-1
53 – 93	keV	133534	134902	144407		376	HXT Y
1.2 – 5.6	MeV	133534	134634	144407		116	GRS Y
15.4	GHz	1334.0	1349.0	1521.0		3.71	
8.8	GHz	1334.0	1349.0	1521.0		3.18	
5	GHz	1333.0	1346.0	1521.0		3.38	
2.7	GHz	1333.0	1417.0	1521.0		4.68	
1.4	GHz	1345.0	1418.0	1521.0	[U0.4 P1.4 U8.8/15.4]	4.73	
410	MHz	1346.0	1347.0	1514.0		3.18	
245	MHz	1346.0	1347.0	1424.0		4.79	
127	MHz	1346.5	1349.4	1406.5		>4.41	
DS II	25-180	1347		1350		3	
DS IV	30-80	1352		1505		3	
DS IV	40-800	~1406		~1519	RS	3	
DS III	25-180	1344		1522	N	2	
DS DCIM	2000-4500	1331		1552	GG	3	
DS DCIM	800-2000	1332		1615	GG	2	
°n						Bolivia	
CME	WL	1031	1184 km/s	-35.9 km/s ²	360°	221°	



Particle event: To($E_p > 10$ MeV) – 18d03^h

Tmax($E_p > 10$ MeV) – 18d10^h, Jmax ($E_p > 10$ MeV) – 190 /cm².s.sr

Duration of the event – 4 days

Quasimaximal energy of protons in the event – E_{qm} = 2100 MeV

Sources: ☐ solar flare 18d02^h11^m, C2.2, s20w90*, AR9415


Main X-ray burst 1-8 Å: onset – 18d02^h11^m, max – 18d02^h14^m, $\Phi = 0.0004$ J/m²

CME: 18d02^h30^m, V=2465 km/s, $\Delta\phi = 360^\circ$, dA = 263°



Δ SC 21d16^h01^m

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 April 18



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 April 18

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	03 ^h	10 ^h	342	3d	
EPS	>10	03 ^h	10 ^h	190	3d	
EPS	>30	03 ^h	06 ^h	64.5	3d	
EPS	>50	03 ^h	06 ^h	33.9	3d	
EPS	>60	03 ^h	06 ^h	26.4	3d	
EPS	>100	03 ^h	06 ^h	12.4	3d	
POES-16						
MEPED	>0.24	03 ^h	05 ^h	1572	3d	
MEPED	>0.8	03 ^h	05 ^h	1004	3d	
MEPED	>2.5	03 ^h	05 ^h	830	3d	
MEPED	>6.9	03 ^h	05 ^h	620	3d	
MEPED	>16	03 ^h	05 ^h	197	3d	
MEPED	>36	03 ^h	05 ^h	90	3d	
MEPED	>70	03 ^h	05 ^h	15.4	3d	
MEPED	>140	03 ^h	05 ^h	6.5	3d	
IMP-8						
CPME	>1	-	07 ^h	1470	-	
CPME	>4	-	07 ^h	296	-	
CPME	>10	-	07 ^h	156	-	
CPME	>30	03 ^h	06 ^h	54.2	4 d	
CPME	>60	03 ^h	06 ^h	27.4	4 d	

ACE						
SIS	>10	03 ^h	09 ^h	320	4 d	
SIS	>30	03 ^h	09 ^h	126	4 d	
SOHO						
EPHIN (INT)	>50	03 ^h	09 ^h	16.9	3d	
BALLOONS						
Mi	>153		06 ^h 45 ^m -07 ^h 39 ^m	9.7		
Mi	>287		06 ^h 45 ^m -07 ^h 39 ^m	2.07		
Mi	>707		06 ^h 45 ^m -07 ^h 39 ^m	0.15		
NM						
Network	>433	-	04 ^h 20 ^m	0.41	-	
Network	>1000	-	04 ^h 20 ^m	0.046	-	
Network	>2000	-	04 ^h 20 ^m	0.0075	-	
Network	>3700	-	04 ^h 20 ^m	0.0014	-	
Network	>5800	-	04 ^h 20 ^m	0.00036	-	

Differential fluxes of protons for the event of 2001 April 18

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	04 ^h	10 ^h	771	7d	
CPME	2-4.6	06 ^h	10 ^h	269	7d	
CPME	4.6-15	03 ^h	10 ^h	30.3	6d	
CPME	15-25	03 ^h	12 ^h	8.2	5d	
CPME	25-48	03 ^h	10 ^h	1.6	5d	
CPME	48-96	03 ^h	10 ^h	0.4	4d	
CPME	96-145	03 ^h	10 ^h	0.34	4d	
CPME	145-440	02 ^h	10 ^h	0.026	3d	
SOHO						
LION	0.75-2	05 ^h	10 ^h	41	6d	
LION	2-6	04 ^h	10 ^h	13	6d	
EPHIN	4-8	03 ^h	09 ^h	70	6d	
EPHIN	8-25	02 ^h	09 ^h	11.9	6d	
EPHIN	25-41	03 ^h	09 ^h	1.6	6d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Struminsky A.B., 2003.
Matsubara Y., Y. Muraki, T. Sako et al., 2005.
Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Miroshnichenko L.I. and J. Perez-Peraza, 2008.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2001 April 18

2001	April 18	☐			AR9415	To event 394	
H α	6563 Å	No data			s20w90*		
1 – 12	keV	0211	0214	0216		C2.2	4.0E-4
53 – 93	keV	021343	021443	021755		20	HXT Y
15.4	GHz	0213.0	0214.0	0219.0		1.90	
8.8	GHz	0213.0	0214.0	0219.0		2.20	
5	GHz	0213.0	0214.0	0221.0		2.67	
2.7	GHz	0213.0	0215.0	0217.0	U0.6 P2.7	2.80	
1.4	GHz	0213.0	0215.0	0218.0		2.62	
610	MHz	0215.0	0216.0	0217.0		1.88	
410	MHz	0214.0	0214.0	0221.0		3.65	
245	MHz	0214.0	0214.0	0215.0		3.26	
DS II	25-500	0217		0246		3	
DS III	23-480	0214		0216	G	3	
DS CONT	300-470	0216		0221		1	
CME	WL	0230	2465 km/s	-9.5 km/s ²	360°	263°	

* – probable localization of the flare event



Particle event: To($E_p > 10$ MeV) – 27d03^h

Tmax($E_p > 10$ MeV) – 28d05^h, Jmax ($E_p > 10$ MeV) – 15 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm} = 80$ MeV


Sources: ●◇ solar flare 26d11^h26^m, M7.8/2B, N17W31, AR9433

Main X-ray burst 1-8 Å: onset – 26d11^h26^m, max – 26d13^h20^m, $\Phi = 0.092$ J/m²



CME 26d12^h30^m, V=1006 km/s, $\Delta\phi = 360^\circ$; dA = 037°;

▲ SC 28d05^h00^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 April 27



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 April 27

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	03 ^h	28d05 ^h	388	2d	
EPS	>10	03 ^h	28d05 ^h	15	2d	
EPS	>30	03 ^h	28d05 ^h	0.2	2d	
EPS	>50	03 ^h	28d05 ^h	0.07	-	
EPS	>60	03 ^h	28d05 ^h	0.04	-	
EPS	>100	03 ^h	28d05 ^h	0.02	-	
POES-16						
MEPED	>0.24	03 ^h	28d04 ^h	45520	4d	
MEPED	>0.8	03 ^h	28d04 ^h	19820	4d	
MEPED	>2.5	03 ^h	28d04 ^h	3170	4d	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	-	-	-	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	26d17 ^h	28d04 ^h	16700	3d	
CPME	>4	26d23 ^h	28d04 ^h	715	3d	
CPME	>10	26d23 ^h	28d04 ^h	9.5	2d	
CPME	>30	-	28d03 ^h	0.16	-	
CPME	>60	-	28d03 ^h	0.05	-	

ACE						
SIS	>10	03 ^h	28d04 ^h	4.5	2d	
SIS	>30	-	-	-	-	
SOHO						
EPHIN (INT)	>50	-	28d04 ^h	0.17	0.4d	


Differential fluxes of protons for the event of 2001 April 27

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	26d15 ^h	28d06 ^h	6560	6d	
CPME	2-4.6	26d17 ^h	28d04 ^h	2230	6d	
CPME	4.6-15	26d21 ^h	28d04 ^h	47.4	6d	
CPME	15-25	01 ^h	28d04 ^h	0.09	6d	
CPME	25-48	01 ^h	28d02 ^h	0.0015	2d	
CPME	48-96	-	-	-	-	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	26d21 ^h	28d04 ^h	444	6d	
LION	2-6	26d21 ^h	28d03 ^h	58	6d	
EPHIN	4-8	26d21 ^h	28d04 ^h	114	6d	
EPHIN	8-25	26d23 ^h	28d03 ^h	1.5	6d	
EPHIN	25-41	08 ^h	28d03 ^h	0.0008	4d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 April 27

2001	April 26	•	AR9433	To event 395			
H α	6563 Å	1211	1311	1618	N17W31	2B	FZT
1 – 12	keV	1126	1312	1319		M7.8	9.2E-2
53 – 93	keV	<115658	~115658	122528		8	HXT Y
53 – 93	keV	<130816	131007	131904		26	HXT Y

15.4	GHz	1307.0	1310.0	1333.0		2.79	
8.8	GHz	1307.0	1310.0	1333.0		2.96	
6.7	GHz	1305.2	1310.2	1312.0	[U0.6 P1.4 U2.7] P6.7	3.12	
5	GHz	1307.0	1309.0	1333.0		2.73	
2.7	GHz	1307.0	1307.0	1317.0		2.56	
1.4	GHz	1307.0	1308.0	1333.0		3.43	
610	MHz	1307.0	1310.0	1333.0		2.62	
410	MHz	1307.0	1308.0	1312.0		3.11	
245	MHz	1307.0	1311.0	1323.0		3.23	
DS II	38-50	1335		1339		2	
DS IV	30-80	1341		2314		2	
DS III	30-80	1309		1310		2	
DS CONT	25-180	1241		1341		1	
DS DCIM	800-2000	1222		1312	GG	3	
DS DCIM	2000-4500	1307		1311	GG	3	
CME	WL	1230	1006km/s	21.1 km/s ²	360°	037°	



Particle event: To(Ep>10 MeV) – 07d14^h

Tmax₁(Ep>10 MeV) – 07d18^h, Jmax₁(Ep>10 MeV) – 7.7 /cm².s.sr

Tmax₂(Ep>10 MeV) – 08d12^h, Jmax₂(Ep>10 MeV) – 11.5 /cm².s.sr

Duration of the event – 3 days


Quasimaximal energy of protons in the event – Eqm₁ = 80 MeV

– Eqm₂ = 85 MeV



Sources: ☐ back side solar flare event < 07d12^h06^m, AR unknown, behind W_L

CME: 07d12^h06^m, V=1223 km/s, Δφ=205°, dA= 267°

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 May 07



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 May 07


S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	14 ^h	19 ^h /08d12 ^h	14.2/46.6	3d	
EPS	>10	14 ^h	18 ^h /08d12 ^h	7.7/11.5	3d	
EPS	>30	14 ^h	18 ^h /08d12 ^h	0.9/0.26	2d	
EPS	>50	14 ^h	14 ^h /08d12 ^h	0.2/0.07	1.5d	
EPS	>60	14 ^h	14 ^h /08d12 ^h	0.08/ 0.02	1d	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	-	18 ^h /08d13 ^h	90/570	-	
MEPED	>0.8	-	18 ^h /08d13 ^h	80/440	-	
MEPED	>2.5	-	18 ^h /08d13 ^h	50/215	-	
MEPED	>6.9	-	18 ^h /08d13 ^h	24/16	-	
MEPED	>16	-	18 ^h /08d13 ^h	4.4 /4	-	
MEPED	>36	-	18 ^h /08d13 ^h	1.5/0.6	-	
MEPED	>70	-	18 ^h /08d13 ^h	0.2/0.03	-	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	14 ^h	19 ^h /08d12 ^h	50.3/268	2.5d	
CPME	>4	14 ^h	19 ^h /08d11 ^h	18.7/79	2.5d	
CPME	>10	14 ^h	18 ^h /08d09 ^h	6.8/14	2d	
CPME	>30	14 ^h	18 ^h /08d08 ^h	0.7/0.6	1d	
CPME	>60	14 ^h	18 ^h /08d08 ^h	0.2/0.2	1d	

ACE						
SIS	>10	14 ^h	18 ^h /08d13 ^h	6.4/9	2d	
SIS	>30	14 ^h	18 ^h /08d13 ^h	1.4/0.3	1d	
SOHO						
EPHIN (INT)	>50	14 ^h	16 ^h / -	0.07/ -	0.4d	

Differential fluxes of protons for the event of 2001 May 07

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	16 ^h	18 ^h /08d13 ^h	15.3/107	5d	
CPME	2-4.6	15 ^h	18 ^h /08d13 ^h	7.5/43.7	5d	
CPME	4.6-15	13 ^h	18 ^h /08d09 ^h	1.2/5.7	5d	
CPME	15-25	13 ^h	18 ^h /08d08 ^h	0.33/0.67	3d	
CPME	25-48	12 ^h	17 ^h /08d08 ^h	0.03/0.026	2d	
CPME	48-96	12 ^h	16 ^h /16 ^h	0.0037/0.001	1d	
CPME	96-145	13 ^h	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	14 ^h	19 ^h /08d09 ^h	2.1/12.8	5d	
LION	2-6	14 ^h	19 ^h /08d09 ^h	0.87/4.4	5d	
EPHIN	4-8	14 ^h	17 ^h /08d08 ^h	5.34/24.1	5d	
EPHIN	8-25	14 ^h	17 ^h /08d08 ^h	0.96/2.24	5d	
EPHIN	25-41	14 ^h	17 ^h /08d08 ^h	0.5/0.024	5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 May.07

2001	May 7		AR9433	To event 396			
CME	WL	1206	1223 km/s	19.2 km/s ²	205°	267°	

Particle event: To($E_p > 10$ MeV) – 20d07^h

Tmax($E_p > 10$ MeV) – 20d10^h, Jmax ($E_p > 10$ MeV) – 1.8 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm} = 410$ MeV


Sources: • solar flare 20d06^h00^m, M6.4/..., s18w90*, AR9455

Main X-ray burst 1-8 Å: onset – 20d06^h00^m, max – 20d06^h03^m, $\Phi = 0.012$ J/m²



CME: 20d06^h26^m, $V = 546$ km/s, $\Delta\phi = 179^\circ$; dA = 231°

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 May 20



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 May 20

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	07 ^h	10 ^h	1.9	3d	
EPS	>10	07 ^h	10 ^h	1.8	3d	
EPS	>30	07 ^h	10 ^h	1.5	2d	
EPS	>50	07 ^h	10 ^h	1	2d	
EPS	>60	07 ^h	10 ^h	0.8	2d	
EPS	>100	07 ^h	09 ^h	0.46	2d	
POES-16						
MEPED	>0.24	07 ^h	-	-	-	
MEPED	>0.8	07 ^h	-	-	-	
MEPED	>2.5	07 ^h	-	-	-	
MEPED	>6.9	07 ^h	-	-	-	
MEPED	>16	07 ^h	09 ^h	5.2	2d	
MEPED	>36	07 ^h	09 ^h	3.2	2d	
MEPED	>70	07 ^h	09 ^h	1.0	2d	
MEPED	>140	07 ^h	09 ^h	0.6	2d	
IMP-8						
CPME	>1	07 ^h	11 ^h	8.7	-	
CPME	>4	07 ^h	11 ^h	6.7	-	
CPME	>10	07 ^h	11 ^h	4.4	-	
CPME	>30	07 ^h	11 ^h	1.6	-	
CPME	>60	07 ^h	10 ^h	0.85	-	

ACE						
SIS	>10	07 ^h	11 ^h	11.2	2d	
SIS	>30	07 ^h	10 ^h	5.1	1.5 d	
SOHO						
EPHIN (INT)	>50	07 ^h	11 ^h	0.62	2d	

Differential fluxes of protons for the event of 2001 May 20

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	11 ^h	<18 ^h	>4.5	8d	
CPME	2-4.6	08 ^h	<18 ^h	>1.6	8d	
CPME	4.6-15	07 ^h	>11 ^h	>0.26	5d	
CPME	15-25	07 ^h	>11 ^h	>0.17	4d	
CPME	25-48	07 ^h	>11 ^h	>0.045	4d	
CPME	48-96	07 ^h	10 ^h	0.011	3d	
CPME	96-145	07 ^h	10 ^h	0.008	2d	
CPME	145-440	07 ^h	10 ^h	0.002	0,5d	
SOHO						
LION	0.75-2	08 ^h	14 ^h	0.7	6d	
LION	2-6	08 ^h	16 ^h	0.28	6d	
EPHIN	4-8	08 ^h	15 ^h	2.11	6d	
EPHIN	8-25	07 ^h	15 ^h	0.47	6d	
EPHIN	25-41	07 ^h	14 ^h	0.08	6d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 May 20

2001 May 20 • AR9415 To event 397

Hα	6563 Å	No Flare			s18w90*		
1 – 12	keV	0600	0603	0606		M6.4	1.2E-2
53 – 93	keV	054613	060253	060739		52	HXT Y

15.4	GHz	0601.0	0602.0	0603.0		2.43	
8.8	GHz	0601.0	0602.0	0605.0		2.54	
5	GHz	0601.0	0602.0	0605.0	U0.5 P5	2.61	
2.7	GHz	0602.0	0602.0	0606.0		2.56	
1.4	GHz	0602.0	0602.0	0607.0		2.40	
900	MHz	0601.5	0603.4	0604.2		2.41	
610	MHz	0602.0	0603.0	0605.0		1.88	
500	MHz	0601.0	0604.0	0613.0		1.65	
410	MHz	0602.0	0602.0	0603.0		1.68	
245	MHz	0602.0	0605.0	0607.0		2.66	
200	MHz	0604.0	0606.0	0606.0		3.09	
DS II	25-340	0605		0624	SH	3	
DS III	25-300	0602		0604	G	3	
DS CONT	23-180	0613		0655		1	
DS DCIM	800-2000	0601		0605	G	2	
CME	WL	0626	546 km/s	-0.1 km/s ²	179°	231°	

* – probable localization of the flare event



Particle event: To(Ep>10 MeV) – 15d16^h

Tmax₁(Ep>10 MeV) – 15d20^h, Jmax₁(Ep>10 MeV) – 5 /cm².s.sr

Tmax₂(Ep>10 MeV) – 16d06^h, Jmax₂(Ep>10 MeV) – 8.1 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 335 MeV


– Eqm₂ = 120 MeV

Sources: ☐ back side solar flare event < 15d15^h56^m, AR unknown, behind W_L



CME 15d15^h56^m, V=1701 km/s, Δφ = 360°; dA = 255°;

o solar flare 15d10^h01^m, M6.3/1N, S26E42 AR9502

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 June 15



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 June 15

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	16 ^h	20 ^h /16d06 ^h	5.9/25.1	3d	
EPS	>10	16 ^h	20 ^h /16d06 ^h	5.0/8.1	3d	
EPS	>30	16 ^h	18 ^h /16d05 ^h	1.9/0.65	3d	
EPS	>50	16 ^h	17 ^h /16d05 ^h	0.9/0.14	2d	
EPS	>60	16 ^h	17 ^h /16d05 ^h	0.59/0.1	2d	
EPS	>100	16 ^h	17 ^h /16d05 ^h	0.18/0.02	2d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	17 ^h	18 ^h / -	70/ -	> 2d	
MEPED	>16	17 ^h	18 ^h / -	12.5/ -	> 1d	
MEPED	>36	17 ^h	18 ^h / -	4.9/ -	> 1d	
MEPED	>70	17 ^h	18 ^h / -	0.4/ -	> 1d	
MEPED	>140	-	-	-	-	
IMP-8						
CPME	>1	17 ^h	23 ^h /16d06 ^h	218/450	5d	
CPME	>4	16 ^h	23 ^h /16d06 ^h	70/46	4.5d	
CPME	>10	16 ^h	19 ^h /16d06 ^h	18.2/8.8	4d	
CPME	>30	16 ^h	18 ^h /16d05 ^h	3.9/0.4	2d	
CPME	>60	16 ^h	18 ^h / -	1.45/ -	2d	
ACE						
SIS	>10	16 ^h	19 ^h /16d10 ^h	25.2/11.5	3 d	
SIS	>30	16 ^h	19 ^h /16d10 ^h	8.2/4	2 d	
SOHO						
EPHIN (INT)	>50	16 ^h	17 ^h /16d08 ^h	0.66/0.02	1d	

Differential fluxes of protons for the event of 2001 June 15



S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	19 ^h	23 ^h /16d10 ^h	99.5/289	-	
CPME	2-4.6	18 ^h	23 ^h /16d09 ^h	38.8/53.5	-	
CPME	4.6-15	17 ^h	23 ^h /16d09 ^h	5/3.5	-	
CPME	15-25	16 ^h	19 ^h /16d06 ^h	0.83/0.36	2d	
CPME	25-48	16 ^h	18 ^h /16d06 ^h	0.14/0.023	2d	
CPME	48-96	16 ^h	17 ^h /16d06 ^h	0.025/0.0025	1.5d	
CPME	96-145	16 ^h	-	-	-	
CPME	145-440	16 ^h	-	-	-	
SOHO						
LION	0.75-2	18 ^h	23 ^h /16d05 ^h	14.2/18.8	5d	
LION	2-6	18 ^h	23 ^h /16d05 ^h	4.2/3	5d	
EPHIN	4-8	18 ^h	22 ^h /16d08 ^h	22/14.7	5d	
EPHIN	8-25	16 ^h	20 ^h /16d08 ^h	1.7/1.24	4d	
EPHIN	25-41	16 ^h	19 ^h /16d08 ^h	0.14/0.03	3.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 June 15

2001		June 15		☐	AR		To event 398	
CME	WL	1556	1701 km/s		56.9* km/s ²	360°	255°	

2001		June 15		O	AR9502		To event 398	
H α	6563 Å	1005	1008	1108	S26E41	1N	FH	
1 – 12	keV	1001	1013	1020		M6.3	4.2E-2	
53 – 93	keV	<102103	102103	104949		13	HXT Y	
15.4	GHz	1006.0	1007.0	0000.0		1.88		
8.8	GHz	1005.0	1007.0	0000.0		2.18		
5	GHz	1004.0	1007.0	0000.0	P0.2 [U2.7 P5]\15.4	2.23		
2.7	GHz	1005.0	1011.0	0000.0		2.11		
1.4	GHz	1006.0	1012.0	1023.0		2.32		
610	MHz	1008.0	1010.0	1029.0		3.18		
410	MHz	1006.0	1012.0	0000.0		3.28		
245	MHz	1006.0	1013.0	0000.0		3.65		
204	MHz	1007.2	1008.2	1041.4		4.31		
127	MHz	1004.0	1026.0	1117.0		3.76		
DS II	110-460	1006		1011	H	3		
DS IV	45-270	1009		~1026		2		
DS III	25-270	1007		1011	G,HARM	2		

DS CONT	45-240	1008		~1010		2	
DS DCIM	2000-4500	1004		1038	GG	2	
DS DCIM	800-2000	1005		1037	GG	2	
DS DCIM	110-2800	1008		1121	P,S,F	3	
5	GHz	1047.0	1049.0	0000.0		2.28	
2.7	GHz	1047.0	1049.0	0000.0		2.41	
1.4	GHz	1047.0	1048.0	0000.0	P0.2 U0.6 P1.4	2.56	
610	MHz	1047.0	1048.0	0000.0		2.15	
410	MHz	1047.0	1049.0	0000.0		2.83	
245	MHz	1006.0	1048.0	0000.0		3.76	
204	MHz	1041.9	1048.5	1129.2		3.63	
DS II	54-65	1047		1049		2	
DS IV	45-270	1046		~1120		2	
DS III	25-100	1047		1048	G	2	
DS DCIM	2000-4500	1047		1118	GG	2	
DS DCIM	800-2000	1047		1121	GG	2	
CME	WL	1032	1090 km/s	9.7 km/s ²	119°	153°	



Particle event: To($E_p > 10$ MeV) – 09d20^h

Tmax($E_p > 10$ MeV) – 10d11^h, Jmax ($E_p > 10$ MeV) – 6 /cm².s.sr

Duration of the event – 1.5 days


Quasimaximal energy of protons in the event – $E_{qm} = 85$ MeV

Sources: ☉ solar flare 09d11^h16^m, C3.7/SF (DSF)



Main X-ray burst 1-8 Å: onset – 09d11^h16^m, max – 09d11^h22^m, $\Phi = 0.0021$ J/m²

CME 09d10^h32^m, V=0479 km/s, $\Delta\phi = 175^\circ$, dA = 255°

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 August 09



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 August 09

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	20 ^h	10d11 ^h	25.2	1.5d	
EPS	>10	20 ^h	10d11 ^h	6	1d	
EPS	>30	21 ^h	10d11 ^h	0.26	1d	
EPS	>50	-	10d11 ^h	0.1	0.5d	
EPS	>60	-	10d12 ^h	0.04	0.3d	
EPS	>100	-	10d12 ^h	0.013	0.3d	
POES-16						
MEPED	>0.24	-	10d10 ^h	3304	-	
MEPED	>0.8	-	10d10 ^h	1919	-	
MEPED	>2.5	-	10d10 ^h	437	-	
MEPED	>6.9	-	10d10 ^h	69	-	
MEPED	>16	-	10d10 ^h	0.4	-	
MEPED	>36	-	10d10 ^h	-	-	
MEPED	>70	-	10d10 ^h	-	-	
MEPED	>140	-	10d10 ^h	-	-	
IMP-8						
CPME	>1	18 ^h	10d12 ^h	970	5 d	
CPME	>4	18 ^h	10d12 ^h	98.7	4 d	
CPME	>10	18 ^h	10d11 ^h	8.8	3 d	
CPME	>30	18 ^h	10d11 ^h	0.2	0.5d	
CPME	>60	18 ^h	10d11 ^h	0.03	0.3d	

ACE						
SIS	>10	18 ^h	10d11 ^h	18.8	2d	
SIS	>30	18 ^h	10d11 ^h	0.4	5h	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 2001 August 09

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	23 ^h	10d09 ^h	820	5d	
CPME	2-4.6	23 ^h	10d09 ^h	109	5d	
CPME	4.6-15	23 ^h	10d12 ^h	5.9	5d	
CPME	15-25	23 ^h	10d11 ^h	0.26	2d	
CPME	25-48	23 ^h	10d11 ^h	0.011	1d	
CPME	48-96	23 ^h	10d09 ^h	0.0005	1d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	18 ^h	10d12 ^h	57.4	5d	
LION	2-6	18 ^h	10d12 ^h	11.9	5d	
EPHIN	4-8	17 ^h	10d11 ^h	32,3	5d	
EPHIN	8-25	17 ^h	10d11 ^h	2,04	5d	
EPHIN	25-41	17 ^h	10d11 ^h	0,02	3d	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 August 09

2001	August 09	☉	AR	To event 399			
H α	6365 Å	1128	1136	1150	N10E54	SF	F
DSF		<1032					
1 – 12	keV	1116	1122	1127		C3.7	2.1E-3
3	GHz	1118.5	1118.9	1120.2		1.46	
DS CONT	2273-4500	1118		1119		1	
DS DCIM	2000-4000	1118		1119	C	2	
CME	WL	1032	479 km/s	4.4 km/s ²	175°	255°	

2001	August 09	☐	AR9557	To event 399			
SPY	6365 Å	1045		>1101	S14W90		
CME	WL	1032	479 km/s	4.4 km/s ²	175°	255°	

Particle event: To(Ep>10 MeV) – 16d00^h

Tmax₁(Ep>10 MeV) – 16d03^h, Jmax₁(Ep>10 MeV) – 87 /cm².s.sr

Tmax₂(Ep>10 MeV) – 17d12^h, Jmax₂(Ep>10 MeV) – 75 /cm².s.sr

Duration of the event – 8 days

Quasimaximal energy of protons in the event – Eqm₁ = 600 MeV


– Eqm₂ = 475 MeV

Sources: ☐ back side solar flare event < 15d23^h54^m; AR9557?, 5d behind W-limb



CME: 15d23^h54^m, V=1575 km/s, Δφ = 360°; dA = 189°

▲ SC 17d11h03^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 August 16



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 August 16

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Duration	Comments
GOES-10						
EPS	>5	00 ^h	03 ^h /17d12 ^h	87.3/136	10d	
EPS	>10	00 ^h	03 ^h /17d12 ^h	87/75	8d	
EPS	>30	00 ^h	03 ^h /17d12 ^h	86.7/18	8d	
EPS	>50	00 ^h	03 ^h /17d12 ^h	60.6/6.5	8d	
EPS	>60	00 ^h	03 ^h /17d12 ^h	47.0/3.5	8d	
EPS	>100	00 ^h	03 ^h /17d12 ^h	19.8/1.1	7d	
POES-16						
MEPED	>0.24	-	02 ^h /17d12 ^h	2080/820	>2d	
MEPED	>0.8	-	02 ^h /17d12 ^h	1370/510	>2d	
MEPED	>2.5	-	02 ^h /17d12 ^h	900/310	>2d	
MEPED	>6.9	-	02 ^h /17d12 ^h	820/250	>2d	
MEPED	>16	-	02 ^h / -	400/56	-	
MEPED	>36	-	02 ^h / -	205.5/23	-	
MEPED	>70	-	02 ^h / -	59.8/2.9	-	
MEPED	>140	-	02 ^h / -	11.8/ -	-	
IMP-8						
CPME	>1	01 ^h	04 ^h /17d12 ^h	331/940	15d	
CPME	>4	00 ^h	04 ^h /17d12 ^h	289/132	14d	
CPME	>10	00 ^h	03 ^h /17d12 ^h	241/59	13d	
CPME	>30	00 ^h	03 ^h /17d12 ^h	144/15	11d	
CPME	>60	00 ^h	03 ^h /17d12 ^h	71.9/6.6	10d	

ACE						
SIS	>10	00 ^h	03 ^h /17d12 ^h	331/80	11d	
SIS	>30	00 ^h	03 ^h /17d12 ^h	203/25	10d	
SOHO						
EPHIN (INT)	>50	00 ^h	03 ^h	53.8/2.7	3d	
BALLOONS						
Mi	>227		07 ^h 50 ^m -08 ^h 23 ^m / -	1.4/ -		
Mi	>268		07 ^h 50 ^m -08 ^h 23 ^m / -	0.66/ -		
Mi	>439		07 ^h 50 ^m -08 ^h 23 ^m / -	0.11/ -		

Differential fluxes of protons for the event of 2001 August 16

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	05 ^h	05 ^h /17d11 ^h	20.6/616	7d	
CPME	2-4.6	01 ^h	05 ^h /17d11 ^h	16.1/72.4	7d	
CPME	4.6-15	01 ^h	04 ^h /17d11 ^h	7.15/7	7d	
CPME	15-25	01 ^h	04 ^h /17d11 ^h	6.5/2.46	7d	
CPME	25-48	01 ^h	04 ^h /17d11 ^h	3.1/0.49	7d	
CPME	48-96	01 ^h	04 ^h /17d11 ^h	0.98/0.11	7d	
CPME	96-145	01 ^h	03 ^h /17d12 ^h	0.44/0.083	7d	
CPME	145-440	01 ^h	03 ^h /17d12 ^h	0.085/0.0125	6d	
SOHO						
LION	0.75-2	03 ^h	07 ^h /17d12 ^h	93.8/36.5	7d	
LION	2-6	02 ^h	04 ^h /17d12 ^h	36.8/4.3	7d	
EPHIN	4-8	01 ^h	04 ^h /17d11 ^h	8.5/15.3	7d	
EPHIN	8-25	01 ^h	03 ^h /17d11 ^h	5.6/2.5	7d	
EPHIN	25-41	01 ^h	03 ^h /17d10 ^h	2.7/0.43	7d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.
Tylka A.J., O.E. Malandraki, G. Dorrian et al., 2013.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 August 16

2001		August 15		☐	AR		To event 400	
CME	WL	2354	1575 km/s	31.7 km/s ²	360°	189°		

Particle event: To($E_p > 10$ MeV) – 15d12^h

Tmax($E_p > 10$ MeV) – 15d15^h, Jmax($E_p > 10$ MeV) – 6 /cm².s.sr

Duration of the event – 2 days


Quasimaximal energy of protons in the event – $E_{qm} = 150$ MeV

Sources: • solar flare 15d11^h03^m, 1N/M1.5, S21W49, AR9608



Main X-ray burst 1-8 Å: onset – 15d11^h04^m, max – 15d11^h28^m, $\Phi = 0.037$ J/m²

• CME: 15d11^h54^m, $V = 0478$ km/s, $\Delta\phi = 130^\circ$, $dA = 248^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 September 15



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 September 15

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	12 ^h	16 ^h	12.3	2d	
EPS	>10	12 ^h	15 ^h	6	2d	
EPS	>30	12 ^h	15 ^h	0.68	1.5d	
EPS	>50	12 ^h	14 ^h	0.20	1.5d	
EPS	>60	12 ^h	14 ^h	0.13	1d	
EPS	>100	12 ^h	14 ^h	0.04	1d	
POES-16						
MEPED	>0.24	15 ^h	15 ^h	61	2d	
MEPED	>0.8	15 ^h	15 ^h	41	2d	
MEPED	>2.5	15 ^h	15 ^h	22	2d	
MEPED	>6.9	15 ^h	15 ^h	9	2d	
MEPED	>16	15 ^h	15 ^h	3.5	2d	
MEPED	>36	15 ^h	15 ^h	0.9	2d	
MEPED	>70	15 ^h	-	-	-	
MEPED	>140	15 ^h	-	-	-	
IMP-8						
CPME	>1	12 ^h	16 ^h	65	2d	
CPME	>4	12 ^h	16 ^h	20	2d	
CPME	>10	12 ^h	15 ^h	5.2	1.5d	
CPME	>30	12 ^h	15 ^h	0.4	0.7d	
CPME	>60	12 ^h	15 ^h	0.11	0.5d	
ACE						
SIS	>10	12 ^h	16 ^h	8	1d	
SIS	>30	12 ^h	14 ^h	1.1	0.7d	
SOHO						
EPHIN (INT)	>50	12 ^h	13 ^h	0.05	0.3d	




Differential fluxes of protons for the event of 2001 September 15

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	15 ^h	18 ^h	42	3d	
CPME	2-4.6	14 ^h	18 ^h	16	3d	
CPME	4.6-15	13 ^h	18 ^h	1.4	3d	
CPME	15-25	13 ^h	16 ^h	0.19	2d	
CPME	25-48	12 ^h	15 ^h	0.02	1,5d	
CPME	48-96	12 ^h	15 ^h	0.002	1d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	16 ^h	18 ^h	4.3	3d	
LION	2-6	14 ^h	17 ^h	1	3d	
EPHIN	4-8	13 ^h	16 ^h	5.3	3d	
EPHIN	8-25	13 ^h	15 ^h	0.51	3d	
EPHIN	25-41	12 ^h	14 ^h	0.03	2d	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 September 15

2001	September 15	•	AR9608	To event 401		
H α		1103	1109	1237	S21W49	1N
H α		<1124	~1139	1230	S27W53	SF
1 – 12	keV	1004	1128	1154		M1.5
53 – 93	keV	<113224	113332	114002		8
						HXT Y
15.4	GHz	1107.0	1108.0	1109.0	P5	2.00
8.8	GHz	1106.0	1108.0	1119.0		2.36
5	GHz	1107.0	1108.0	1111.0		2.43
3	GHz	1106.6	1108.9	1154.1		2.25
1.4	GHz	1109.0	1109.0	1110.0		1.52
DS DCIM	2000-4500	1106		1143	G	2
DS DCIM	800-2000	1107		1141	G	1
8.8	GHz	1119.0	1121.0	1132.0	U2.7 / 8.8	2.00
5	GHz	1120.0	1122.0	1128.0		1.98
2.7	GHz	1120.0	1122.0	1129.0		1.79
245	MHz	1121.0	1124.0	1131.0		3.04
204	MHz	1124.5	1124.9	1126.2		3.26
DS II	30-220	1121		1128	GG,FS	2
DS III	30-270	1123		1126	GG	2
DS CONT	25-180	1124		1459		1

8.8	GHz	1132.0	1134.0	1139.0		1.78	
5	GHz	1132.0	1134.0	1137.0		1.79	
2.7	GHz	1132.0	1134.0	1139.0	[U1.4] P2.7	2.00	
1.4	GHz	1134.0	1136.0	1137.0		1.46	
245	MHz	1138.0	1138.0	~1138.0		1.57	
DS III	25-90	1142		>1200	N	2	
CME	WL	1154	0478 km/s	-4.0 km/s ²	130°	248°	



Particle event: To($E_p > 10$ MeV) – 24d11^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 24\text{d}18^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 390 \text{ /cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 25\text{d}07^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 1.1 \cdot 10^3 \text{ /cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 3}(E_p > 10 \text{ MeV}) - 25\text{d}22^{\text{h}}$, $J_{\max 3}(E_p > 10 \text{ MeV}) - 9.5 \cdot 10^3 \text{ /cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 5 days

Quasimaximal energy of protons in the event – $E_{qm1} = 470 \text{ MeV}$

– $E_{qm2} = 580 \text{ MeV}$

– $E_{qm3} = 580 \text{ MeV}$

Sources: ● solar flare 24d09^h32^m, X2.6/2B, S17E26, AR9632;


○ solar flare 25d04^h24^m, M7.6/1N, S18W01, AR9628

Main X-ray burst 1-8 Å: onset – 24d09^h32^m, max – 24d10^h38^m, $\Phi = 0.63 \text{ J/m}^2$



CME: 24d10^h31^m, $V = 2402 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 142^\circ$

▲ SC 25d20^h25^m, Δ SC 29d09^h40^m,

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 September 24



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 September 24

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES -10						
EPS	>5	11 ^h	18 ^h /25d07 ^h /25d22 ^h	458/1430/21530	5d	
EPS	>10	11 ^h	18 ^h /25d07 ^h /25d22 ^h	390/1100/9500	5d	
EPS	>30	11 ^h	18 ^h /25d07 ^h /25d22 ^h	161/437/1190	5d	
EPS	>50	11 ^h	18 ^h /25d08 ^h /25d22 ^h	55/173/250	5d	
EPS	>60	11 ^h	18 ^h /25d08 ^h /25d22 ^h	32.4/103/105	5d	
EPS	>100	11 ^h	18 ^h /25d08 ^h /25d22 ^h	7.5/27.7/15	5d	
POES-16						
MEPED	>0.24	11 ^h	18 ^h /25d08 ^h /25d22 ^h	2694/6424/115014	5d	
MEPED	>0.8	11 ^h	18 ^h /25d08 ^h /25d22 ^h	2444/5464/65054	5d	
MEPED	>2.5	11 ^h	18 ^h /25d08 ^h /25d22 ^h	2227/3937 /29387	5d	
MEPED	>6.9	11 ^h	18 ^h /25d08 ^h /25d22 ^h	1770/2060/9920	5d	
MEPED	>16	11 ^h	18 ^h /25d08 ^h /25d22 ^h	990/1480/3130	5d	
MEPED	>36	11 ^h	18 ^h /25d08 ^h /25d22 ^h	402.5/620/680	5d	
MEPED	>70	11 ^h	18 ^h /25d08 ^h /25d22 ^h	31.5/62.8/33.8	5d	
MEPED	>140	11 ^h	18 ^h /25d08 ^h /25d22 ^h	2.43/4.63/1.33	5d	
ACE						
SIS	>10	11 ^h	23 ^h /25d08 ^h /25d22 ^h	1700/2200/3540	5d	
SIS	>30	11 ^h	23 ^h /25d08 ^h /25d22 ^h	500/700/860	5d	
SOHO						
EPHIN (INT)	>50	12 ^h	23 ^h /25d07 ^h /25d21 ^h	81/203/322	6d	

BALLOONS						
Mu	>95		18 ^h / - / -	4.3/ - / -		
Mu	>138		18 ^h / - / -	1.27/ - / -		
Mu	>202		18 ^h / - / -	0.15/ - / -		

Differential fluxes of protons for the event of 2001 September 24

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	13 ^h	22 ^h /25d04 ^h /25d22 ^h	329/890/13200	6.5d	
CPME	2-4.6	11 ^h	22 ^h /25d04 ^h /25d21 ^h	238/560/2800	6.5d	
CPME	4.6-15	11 ^h	22 ^h /25d05 ^h /25d21 ^h	75.6/107/220	6.5d	
CPME	15-25	11 ^h	22 ^h /25d08 ^h /25d21 ^h	55.5/68.3/106	6.5d	
CPME	25-48	11 ^h	22 ^h /25d08 ^h /25d21 ^h	11.6/15.2/16.8	6.5d	
CPME	48-96	11 ^h	22 ^h /25d08 ^h /25d21 ^h	2.4/3.6/6.7	6.5d	
CPME	96-145	11 ^h	-	-	-	
CPME	145-440	11 ^h	-	-	-	
SOHO						
LION	0.75-2	13 ^h	24 ^h /25d07 ^h /25d23 ^h	60/108/2610	6d	
LION	2-6	12 ^h	24 ^h /25d07 ^h /25d23 ^h	31/53/767	6d	
EPHIN	4-8	12 ^h	23 ^h /25d07 ^h /25d22 ^h	127/216/1760	6d	
EPHIN	8-25	12 ^h	23 ^h /25d07 ^h /25d22 ^h	36.7/53/201	6d	
EPHIN	25-41	12 ^h	23 ^h /25d07 ^h /25d22 ^h	6.8/8.8/11,8	6d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 September 24


2001 September 24		• AR9632		To event 402			
Hα		<0936	1019	1217	S16E23	2B	F
1 – 12	keV	0932	1038	1109		X2.6	6.3E-1
53 – 93	keV	094456	~101350	>101350		8	HXT Y
53 – 93	keV	<105750	~105750	113236		8	HXT Y

15.4	GHz	0933.0	1029.0	1203.0		4.18	
9.1	GHz	0932.0	1029.9	>1230.0	[U1.4] P9.1	4.40	
5	GHz	0933.0	1030.0	1324.0		4.34	
3	GHz	~0900.0	1031.1	>1209.0		4.08	
1.4	GHz	0957.0	1019.0	1142.0		3.63	
900	MHz	0830.0	1025.0	>1230.0		4.16	
600	MHz	0854.7	1024.7	1200.0		3.79	
410	MHz	0930.0	1024.0	1235.0		3.23	
245	MHz	0930.0	1048.0	1150.0		3.28	
204	MHz	0931.0	1051.7	1152.0		3.02	
DS IV	1200-3800	0959		1150	P,S	3	
DS IV	25-180	1018		1624		2	
DS III	1200-1700	0935		1123	GG,RS	2	
DS III	25-270	0937		~1017	S	2	
DS III	25-270	1017		~1110	S,C	3	
DS CONT	25-180	0938		1018		2	
DS CONT	40-270	~1019		~1116		3	
DS DCIM	2000-4500	0943		1126	GG	3	
DS DCIM	800-2000	0956		1149	GG	3	
CME	WL	1031	2402 km/s	54.1 km/s ²	360°	142°	



2001	September 25	Ø	AR9628	To event 402			
H α	6563 Å	0426	0440	0506	S18W01	1N	FZ
1 – 12	keV	0424	0440	0452		M7.6	7.2E-2
53 – 93	keV	<043302	043556	051234		19	HXT Y

15.4	GHz	0434.0	0434.0	0503.0		2.71	
8.8	GHz	0425.0	0434.0	0522.0	U1.4 P8.8	2.89	
5	GHz	0429.0	0434.0	0522.0		2.59	
2.8	GHz	0433.0	0435.0	0516.0		2.8	
1.4	GHz	0433.0	0434.0	0450.0		1.4	
900	MHz	0432.7	0434.4	0501.9		900	
610	MHz	0433.0	0434.0	0443.0		610	
410	MHz	0433.0	0434.0	0444.0		410	
245	MHz	0432.0	0433.0	0437.0		245	
DS II	28-120	0440		0456		2	
DS III	80-1000	0432		0443	GG	2	
DS III	57-170	0442		0445	GG	1	
CME	WL	0604	0750km/s	—	062°	283°	



Particle event: To(Ep>10 MeV) – 01d14^hTmax₁(Ep>10 MeV) – 01d23^h, Jmax₁(Ep>10 MeV) – 370/cm².s.srTmax₂(Ep>10 MeV) – 02d07^h, Jmax₂(Ep>10 MeV) – 1.3·10³/cm².s.sr


Duration of the event – 3.5 days

Quasimaximal energy of protons in the event – Eqm₁ = 155 MeV– Eqm₂ = 150 MeV**Sources:** ■ solar flare 01d04^h41^m, M9.1/..., s18w80*, AR9628



◇ flare activity AR9628 behind W-limb

Main X-ray burst 1-8 Å: onset – 01d04^h41^m, max – 01d05^h15^m, Φ = 0.086 J/m²CME: 01d05^h30^m, V = 1405 km/s, Δφ = 360°, dA = 225°

* – probable localization of the flare event


Particle fluxes and associated phenomena

Time profiles of the proton fluxes for the event of 2001 October 01



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 October 01

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES-10						
EPS	>5	14 ^h	23 ^h /02d07 ^h	633/2186	3.5d	
EPS	>10	14 ^h	23 ^h /02d07 ^h	370/1300	3.5d	
EPS	>30	14 ^h	23 ^h /02d08 ^h	57.2/151	3d	
EPS	>50	14 ^h	22 ^h /02d08 ^h	15.4/20.6	2d	
EPS	>60	14 ^h	22 ^h /02d08 ^h	2.1/3.9	1.5d	
EPS	>100	14 ^h	21 ^h /02d08 ^h	0.09/0.12	1d	
POES-16						
MEPED	>0.24	13 ^h	21 ^h /2d08 ^h	3090/7190	6d	
MEPED	>0.8	13 ^h	21 ^h /2d08 ^h	2520/5840	6d	
MEPED	>2.5	13 ^h	21 ^h /2d08 ^h	1920/4460	6d	
MEPED	>6.9	13 ^h	21 ^h /2d08 ^h	1195/2920	6d	
MEPED	>16	13 ^h	21 ^h /2d08 ^h	414/1200	6d	
MEPED	>36	13 ^h	21 ^h /2d08 ^h	65.1/128	6d	
MEPED	>70	13 ^h	21 ^h /2d08 ^h	0.76/1.2	6d	
MEPED	>140	13 ^h	-	-	-	

CORONAS-F						
MKL	>1	14 ^h	21 ^h /2d08 ^h	1020/2400	6d	
MKL	>14	14 ^h	21 ^h /2d08 ^h	185/610	6d	
MKL	>26	14 ^h	21 ^h /2d08 ^h	136/380	6d	
MKL	>50	14 ^h	21 ^h /2d10 ^h	53/113	6d	
ACE						
SIS	>10	13 ^h	23 ^h /02d07 ^h	273/1380	3d	
SIS	>30	13 ^h	23 ^h /02d07 ^h	53/160	3d	
SOHO						
EPHIN (INT)	>50	12 ^h	23 ^h /02d06 ^h	3.5/8.6	2d	

Differential fluxes of protons for the event of 2001 October 01

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Duration	Comments
IMP-8						
CPME	1-2	14 ^h	23 ^h /02d08 ^h	620/1350	6d	
CPME	2-4.6	14 ^h	22 ^h /02d08 ^h	178/490	6d	
CPME	4.6-15	14 ^h	21 ^h /02d08 ^h	41/115	6d	
CPME	15-25	14 ^h	21 ^h /02d08 ^h	19.3/70	6d	
CPME	25-48	14 ^h	21 ^h /02d08 ^h	2.33/7.3	4d	
CPME	48-96	14 ^h	21 ^h /02d08 ^h	0.2/0.45	4d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	13 ^h	23 ^h /02d07 ^h	31.5/110	6d	
LION	2-6	13 ^h	23 ^h /02d07 ^h	10.8/38.8	6d	
EPHIN	4-8	12 ^h	23 ^h /02d07 ^h	74/210	6d	
EPHIN	8-25	12 ^h	23 ^h /02d07 ^h	19.8/53.2	6d	
EPHIN	25-41	12 ^h	23 ^h /02d06 ^h	2.1/4.9	4d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 October 01

2001 October 01


AR9628

To event 403

H α	6563 Å	No Flare Patrol			s18w80*		
1 – 12	keV	0441	0515	0523		M9.1	8.6E-2
53 – 93	keV	045617	051221	052017		8	HXT Y

15.4	GHz	0500.0	0500.0	~0500.0		1.58	
8.8	GHz	<0509.0	~0511.0	>0515.0	U0.9 P8.8	2.11	
5	GHz	0459.0	0500.0	0501.0		2.04	
2.8	GHz	0449.0	0511.9	0533.0		1.96	
1.4	GHz	<0511.0	~0513.0	>0514.0		1.61	
900	MHz	0511.7	0511.9	0534.6		1.45	
600	MHz	~0511.6	0515.2	>0533.0		1.79	
245	MHz	0524.0	0524.0	~0524.0		1.91	
DS III	57-160	0451		0529	N	1	
DS III	57-220	0522		0525	G	2	
DS III	57-750	0529		>0755	S,C	3	
CME	WL	0530	1405 km/s	97.8 km/s ²	360°	225°	

* – probable localization of the flare event



Particle event: To($E_p > 10$ MeV) – 19d02^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 19d08^h$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 3.6/\text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 19d21^h$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 8/\text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm1} = 300 \text{ MeV}$

– $E_{qm2} = 310 \text{ MeV}$

Sources: • solar flare 19d00^h47^m, X1.6/2B, N16W18, AR9661*

M1.2/2B, N16W18, AR9661*

Ø solar flare 19d16^h13^m, X1.6/2B, N15W30, AR9661


Main X-ray burst 1–8 Å: onset – 19d00^h47^m, max – 19d01^h05^m, $\Phi = 0.12 \text{ J/m}^2$

CME: 19d01^h27^m, $V = 0558 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 284^\circ$



$\Delta \text{SC } 21d16^h48^m$

* One flare event with two X-ray bursts

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 October 19



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 October 19

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Duration	Comments
GOES-10						
EPS	>5	02 ^h	09 ^h /22 ^h	6.6/16.1	3d	
EPS	>10	02 ^h	08 ^h /21 ^h	3.6/8	3d	
EPS	>30	02 ^h	08 ^h /20 ^h	0.8/1.7	3d	
EPS	>50	02 ^h	08 ^h /20 ^h	0.35/0.59	3d	
EPS	>60	01 ^h	07 ^h /19 ^h	0.25/0.39	3d	
EPS	>100	01 ^h	07 ^h /18 ^h	0.12/0.15	3d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	05 ^h /19 ^h	15/15	3d	
MEPED	>16	-	05 ^h /19 ^h	3.1/6.1	3d	
MEPED	>36	-	05 ^h /19 ^h	1.5/2.05	3d	
MEPED	>70	-	05 ^h / -	0.35/ -	3d	
CORONAS-F						
MKL	>1	14 ^h	- /18 ^h	- /51	3d	
MKL	>14	14 ^h	08 ^h / -	4/ -	3d	
MKL	>26	14 ^h	08 ^h / -	3.3/ -	3d	
MKL	>50	14 ^h	08 ^h / -	1.8/ -	3d	
ACE						
SIS	>10	02 ^h	05 ^h /19 ^h	11/9.9	3d	
SIS	>30	02 ^h	05 ^h /19 ^h	4/3	3d	
SOHO						
EPHIN (INT)	>50	02 ^h	03 ^h /19 ^h	0.21/0.3	1,5d	


Differential fluxes of protons for the event of 2001 October 19

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	07 ^h	11 ^h /20 ^h	11/24.8	3,5d	
CPME	2-4.6	06 ^h	07 ^h /20 ^h	3.5/7.5	3,5d	
CPME	4.6-15	04 ^h	07 ^h /20 ^h	0.55/1.1	3d	
CPME	15-25	03 ^h	06 ^h /20 ^h	0.18/0.3	3d	
CPME	25-48	03 ^h	06 ^h /20 ^h	0.03/0.05	3d	
CPME	48-96	03 ^h	06 ^h /19 ^h	0.0085/0.01	3d	
CPME	96-145	03 ^h	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	04 ^h	07 ^h /20 ^h	0.15/1.6	3d	
LION	2-6	04 ^h	07 ^h /20 ^h	0.12/0.3	3d	
EPHIN	4-8	04 ^h	06 ^h /20 ^h	0.7/2.94	3d	
EPHIN	8-25	03 ^h	05 ^h /20 ^h	0.1/0.57	3d	
EPHIN	25-41	02 ^h	04 ^h /19 ^h	0.016/0.06	3d	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 October 19


2001 October 19		• AR9691		To event 404			
H α	6563 Å	0049	0059	0355	N16W18	2B	EZ
1 – 12	keV	0047	0105	0113		X1.6	1.2E-1
1 – 12	keV	0220	0232	0246		M1.2	1.8E-2
15.4	GHz	0053.0	0100.0	0237.0		2.83	
8.8	GHz	0049.0	0100.0	0237.0	P0.4-0.6 [U5 P8.8]\15.4	3.00	
5	GHz	0050.0	0100.0	0239.0		2.77	
2.8	GHz	0050.0	0126.0	0152.0		3.18	
1.4	GHz	0054.0	0124.0	0201.0		3.92	
610	MHz	0056.0	0124.0	0237.0		4.51	
410	MHz	0056.0	0140.0	0206.0		4.52	
245	MHz	0059.0	0140.0	0206.0		4.00	
DS II	25-220	0101		0119		3	
DS IV	57-1600	0058		>0200	FS	3	
DS III	57-180	~0120		0331	S,C	2	
CME	WL	0127	0558 km/s	-25.6 km/s ²	360°	284°	

* One flare event with two X-ray bursts



2001 October 19 Ø AR9661 To event 404

H α		1616	1636	1820	N15W30	2N	FUY
1 – 12	keV	1613	1630	1643		X1.6	1.6E-1
1,3-4,0	MeV						SONG-F
15.4	GHz	1614.0	1624.0	1705.0	[P1.4] U2.7 / 15.4	3.62	
8.8	GHz	1615.0	1625.0	1705.0		3.56	
6.7	GHz	1614.0	1625.0	1648.5		3.32	
5	GHz	1618.0	1625.0	1705.0		3.15	
2.7	GHz	1622.0	1625.0	1705.0		2.98	
1.4	GHz	1622.0	1624.0	1705.0		4.18	
410	MHz	1624.0	1624.0	1705.0		4.00	
245	MHz	1624.0	1624.0	1705.0		4.11	
DS II	25-180	1624		1642		2	
DS IV	25-180	1640		1704		2	
CME	WL	1650	0901 km/s	-0.7 km/s ²	360°	273°	



Particle event: To($E_p > 10$ MeV) – 22d16^h

Tmax($E_p > 10$ MeV) – 22d21^h, Jmax ($E_p > 10$ MeV) – 17 /cm².s.sr

Duration of the event – 2.5 days

Quasimaximal energy of protons in the event – $E_{qm} = 425$ MeV

Sources: ● solar 22d17^h44^m, X1.2/2B, S18E16, AR9672


○ solar 22d14^h16^m, 2N/M6.7, S17E20, AR9672

Main X-ray burst 1-8 Å: onset – 22d17^h44^m, max – 22d17^h59^m, $\Phi = 0.12$ J/m²



CME: 22d18^h26^m, $V = 0618$ km/s, 106°, dA = 126°

▲ SC 25d08^h50^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 October 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 October 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Duration	Comments
GOES-10						
EPS	>5	16 ^h	22 ^h	53.6	2.5d	
EPS	>10	16 ^h	21 ^h	17.0	2.5d	
EPS	>30	16 ^h	22 ^h	3.8	2.5d	
EPS	>50	16 ^h	22 ^h	1.9	2.5d	
EPS	>60	16 ^h	22 ^h	1.5	2.5d	
EPS	>100	16 ^h	22 ^h	0.7	2.5d	
POES-16						
MEPED	>0.24	18 ^h	20 ^h	663	2.5d	
MEPED	>0.8	18 ^h	20 ^h	478	2.5d	
MEPED	>2.5	18 ^h	20 ^h	236	2.5d	
MEPED	>6.9	18 ^h	20 ^h	81	2.5d	
MEPED	>16	18 ^h	20 ^h	9.5	2.5d	
MEPED	>36	18 ^h	20 ^h	5	2.5d	
MEPED	>70	18 ^h	20 ^h	0.95	2.5d	
MEPED	>140	18 ^h	20 ^h	0.45	2.5d	
KORONAS-F						
MKL	>1	18 ^h	23d02 ^h	342	2.5d	
MKL	>14	18 ^h	23d00 ^h	4.6	2.5d	
MKL	>26	18 ^h	23d00 ^h	3.5	2.5d	
MKL	>50	18 ^h	23d00 ^h	2.0	2.5d	

ACE						
SIS	>10	16 ^h	21 ^h	20.4	3d	
SIS	>30	16 ^h	22 ^h	6.9	3d	
SOHO						
EPHIN (INT)	>50	16 ^h	21 ^h	0.73	3d	

Differential fluxes of protons for the event of 2001 October 22


S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
IMP-8						
CPME	1-2	18 ^h	23 ^h	690	>1.5d	
CPME	2-4.6	18 ^h	23 ^h	136	>1.5d	
CPME	4.6-15	17 ^h	23 ^h	6	>1.5d	
CPME	15-25	16 ^h	22 ^h	0.5	>1.5d	
CPME	25-48	17 ^h	23 ^h	0.08	>1.5d	
CPME	48-96	17 ^h	23 ^h	0.028	>1.5d	
CPME	96-145	-	-	-	-	
CPME	145-440	-	-	-	-	
SOHO						
LION	0.75-2	18 ^h	23d01 ^h	18.5	7d	
LION	2-6	18 ^h	23d01 ^h	3.7	7d	
EPHIN	4-8	17 ^h	21 ^h	12.2	7d	
EPHIN	8-25	17 ^h	21 ^h	0.9	7d	
EPHIN	25-41	16 ^h	21 ^h	0.06	7d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 October 22

2001 October 22 • AR9672 To event 405


Hα		1744	1758	1911	S18E16	2B	UZ
1 – 12	keV	1744	1759	1814		X1.2	1.2E-1
53 – 93	keV	<174951	175509	>180251		108	HXT Y

15.4	GHz	1747.0	1751.0	1809.0	U0.4 / 15.4	3.57	
9.5	GHz	1740.2	1750.8	1757.5		3.41	
6.7	GHz	1739.3	1751.0	1758.8		3.39	
5	GHz	1746.0	1751.0	1809.0		3.23	
2.7	GHz	1747.0	1751.0	1808.0		2.90	
1.4	GHz	1747.0	1751.0	1808.0		2.76	
610	MHz	1751.0	1757.0	1808.0		2.78	
410	MHz	1754.0	1757.0	1801.0		2.69	
245	MHz	1755.0	1755.0	1808.0		3.20	
DS II	25-180	1759		1810		1	
DS III	25-180	1754		1856	N	1	
DSF		~1648	1930	1930		14°	
CME	WL	1826	0618 km/s	-7.6 km/s ²	106°	126°	



2001	October 22	Ø	AR9661	To event 405			
H α		1416	1455	1552	S17E20	2N	F
1 – 12	keV	1427	1508	1531		M6.7	1.5E-1
53 – 93	keV	<143823	~144811	>144931		10	HXT Y

15.4	GHz	1444.0	1500.0	1519.0		15.4	
8.8	GHz	1444.0	1457.0	1526.0		8.8	
6.7	GHz	1450.0	1457.4	1511.2	U0.6 P6.7	6.7	
5	GHz	1444.0	1457.0	1526.0		5	
2.7	GHz	1445.0	1457.0	1522.0		2.7	
1.4	GHz	1445.0	1500.0	1522.0		1.4	
610	MHz	1446.0	1451.0	1522.0		610	
410	MHz	1445.0	1452.0	1513.0		410	
245	MHz	1444.0	1452.0	1522.0		245	
DS II	110-150	1452		1457	SH	DS II	
DS II	25-142	1453		1513		DS II	
DS II	30-80	1456		1520		DS II	
DS IV	40-400	1445		1521		DS IV	
DS III	40-250	1445		1452	GG	DS III	
DS DCIM	100-4000	1446		1509	CS	DS DCIM	
CME	WL	1506	1336 km/s	-8.0 km/s ²	360°	122	



Particle event: To($E_p > 10$ MeV) – 28d02^h

Tmax($E_p > 10$ MeV) – 28d07^h, Jmax ($E_p > 10$ MeV) – 1.1 /cm².s.sr

Duration of the event – 1 day

Quasimaximal energy of protons in the event – $E_{qm} = 60$ MeV


Sources: unknown

○ solar flare 28d04^h36^m, M1.3/1F, N12E40, AR9682



Main X-ray burst 1-8 Å: onset – 28d04^h36^m, max – 28d04^h50^m, $\Phi = 0.013$ J/m²

Δ SC 28d13^h19^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 October 28



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 October 28

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	02 ^h	08 ^h	11.6	1.5d	
EPS	>10	02 ^h	07 ^h	1.14	1d	
EPS	>30	02 ^h	07 ^h	0.06	0.5d	
EPS	>50	02 ^h	07 ^h	0.02	-	
EPS	>60	02 ^h	07 ^h	0.01	-	
EPS	>100	-	-	-	-	
ACE						
SIS	>10	06 ^h	07 ^h	1.4	0.5d	
SIS	>30	-	08 ^h	0.1	-	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 2001 October 28

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
SOHO						
LION	0.75-2	26d18 ^h	28d09 ^h	11.4	3d	
LION	2-6	26d18 ^h	28d08 ^h	1.2	3d	
EPHIN	4-8	26d18 ^h	28d08 ^h	1.9	3d	
EPHIN	8-25	-	28d09 ^h	0.045	3d	
EPHIN	25-41	-	28d08 ^h	0.0006	-	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2001 October 28**

2001	October 28	○				AR9682	To event 406
H α		0443	0448	0531	N12E40	1F	F
1 – 12	keV	0436	0450	0458		M1.3	1.3E-2
8.8	GHz	0442.0	0444.0	0502.0		1.75	

Particle event: To(Ep>10 MeV) – 04d16^h

Tmax₁(Ep>10 MeV) – 04d20^h, Jmax₁(Ep>10 MeV) – 540 /cm².s.sr

Tmax₂(Ep>10 MeV) – 06d00^h, Jmax₂(Ep>10 MeV) – 2.4·10⁴ /cm².s.sr

Duration of the event – 5 days

Quasimaximal energy of protons in the event – Eqm₁ = 750 MeV

– Eqm₂ = 685 MeV

Sources: ● solar flare 04d16^h03^m, X1.0/3B, N07W19, AR9684

○ solar flare 04d06^h38^m, C8.4/1N, N14W57, AR9682


○ solar flare 05d09^h07^m, M2.1/1N, N03W37, AR9684

Main X-ray burst 1-8 Å: onset – 04d16^h03^m, max – 04d16^h20^m, Φ = 0.22 J/m²



CME 04d16^h35^m, V = 1810 km/s, Δφ = 360°, dA = 239°,

▲ SC 06d01^h52^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 November 04



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 November 04

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr	Dura-tion	Comments
GOES-10						
EPS	>5	16 ^h	20 ^h /06d00 ^h	570/3.5·10 ⁴	5d	
EPS	>10	16 ^h	20 ^h /06d00 ^h	540/2.3·10 ⁴	5d	
EPS	>30	16 ^h	20 ^h /06d00 ^h	360/5.1·10 ³	5d	
EPS	>50	16 ^h	20 ^h /06d00 ^h	210/1.1·10 ³	5d	
EPS	>60	16 ^h	20 ^h /06d00 ^h	148/670	5d	
EPS	>100	16 ^h	20 ^h /06d00 ^h	59/126	5d	
POES-16						
MEPED	>0.24	16 ^h	22 ^h /06d03 ^h	5.4·10 ³ /5·10 ⁴	5d	
MEPED	>0.8	16 ^h	22 ^h /06d03 ^h	3.8·10 ³ /4·10 ⁴	5d	
MEPED	>2.5	16 ^h	22 ^h /06d03 ^h	3.1·10 ³ /3.2·10 ⁴	5d	
MEPED	>6.9	16 ^h	22 ^h /06d03 ^h	2.6·10 ³ /2.1·10 ⁴	5d	
MEPED	>16	16 ^h	22 ^h /06d03 ^h	1.3·10 ³ /9.3·10 ³	5d	
MEPED	>36	16 ^h	22 ^h /06d03 ^h	840/3·10 ³	5d	
MEPED	>70	16 ^h	22 ^h /06d03 ^h	121/230	5d	
MEPED	>140	16 ^h	22 ^h /06d03 ^h	27.6/29.4	5d	
CORONAS-F						
MKL	>1.	16 ^h	22 ^h /06d02 ^h	4.3·10 ³ /3.8·10 ⁴	5d	
MKL	>14	16 ^h	22 ^h /06d02 ^h	750/5.5·10 ³	5d	
MKL	>26	16 ^h	22 ^h /06d02 ^h	515/4.4·10 ³	5d	
MKL	>50	16 ^h	22 ^h /06d02 ^h	252/1.5·10 ³	5d	

ACE						
SIS	>10	16 ^h	23 ^h /6d00 ^h	1.6·10 ³ /4.6·10 ³	5d	
SIS	>30	16 ^h	23 ^h /6d00 ^h	700/1.9·10 ³	5d	
SOHO						
EPHIN (INT)	>50	16 ^h	20 ^h /06d02 ^h	170/3.2·10 ³	4d	
BALLOONS						
Mu	>254		05d(08 ^h 08 ^m -08 ^h 37 ^m)	1.2/ -		
Mu	>317		05d(08 ^h 08 ^m -08 ^h 37 ^m)	0.32/ -		
Mu	>409		05d(08 ^h 08 ^m -08 ^h 37 ^m)	0.14/ -		
Mu	>184		06d(02 ^h 16 ^m -02 ^h 57 ^m)	- /2.0		
Mu	>265		06d(02 ^h 16 ^m -02 ^h 57 ^m)	- /0.73		
Mu	>390		06d(02 ^h 16 ^m -02 ^h 57 ^m)	- /0.17		
NM						
Network	>500		20 ^h /-	0.3/ -		

Differential fluxes of protons for the event of 2001 November 04

S/c, instru-ments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura-tion	Comments
SOHO						
LION	0.75-2	17 ^h	05d02 ^h /06d02 ^h	49.4/1800	8d	
LION	2-6	17 ^h	05d02 ^h /06d02 ^h	3.4/810	8d	
EPHIN	4-8	17 ^h	06d03 ^h /06d01 ^h	130/1080	8d	
EPHIN	8-25	17 ^h	06d03 ^h /06d01 ^h	46.3/180	8d	
EPHIN	25-41	17 ^h	05d20 ^h /06d02 ^h	12.4/44.1	8d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Struminsky A.B., 2003.
 Kuwabara T., J.W. Bieber, J. Clem, et.al., 2006.
 Rawat R., S. Alex, and G.S. Lakhina, 2006.
 Miroshnichenko L.I. and J. Perez-Peraza, 2008.
 Lario D., A. Aran, R.B. Decker, 2009.
 Tylka A.J., O.E. Malandraki, G. Dorrian et al., 2013.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 November 04


2001 November 04		•		AR9684		To event 407	
Hα		1603	1614	>2340	N07W19	3B	FUZ
1 – 12	keV	1603	1620	1657		X1.0	2.2E-1
53 – 93	keV	160256	161254	>171336		38	HXT Y
15.4	GHz	1605.0	1617.0	1725.0		3.20	
8.8	GHz	1604.0	1616.0	0000.0	U1.4 P5-8.8	3.45	
5	GHz	1604.0	1617.0	0000.0		3.45	
2.7	GHz	1602.0	1624.0	0000.0		3.28	
1.4	GHz	1602.0	1607.0	1727.0		3.20	
610	MHz	1604.0	1609.0	1709.0		3.56	
410	MHz	1605.0	1617.0	1714.0		3.57	

DS II	29-180	1610		1621		2	
DS IV	36-180	1612		1850		1	
DS III	25-180	1611		1715	N	2	
CME	WL	1635	1810 km/s	-63.4 km/s ²	360°	239°	



2001 November 04 Ø AR9682 To event 407

H α		0641	0643	0700	N14W57	1N	C
1 – 12	keV	0638	0643	0646		C8.4	2.3E-3
53 – 93	keV	064056	064306	065144		13	HXT Y
15.4	GHz	0640.0	0641.0	0641.0	3 \ 15.4	2.18	
8.8	GHz	0640.0	0641.0	0643.0		2.28	
5	GHz	0640.0	0641.0	0643.0		2.40	
3	GHz	0640.7	0641.3	0644.1		2.45	
DS I	110-270	0650		>1200	N	1	
DS III	25-132	0639		0640		1	
CME	WL	0750	259 km/s	0.1 km/s ²	136°	011°	



2001


November 05

Ø

AR9684

To event 407

H α		0908	0911	0947	N03W37	1N	EFU
1 – 12	keV	0907	0915	0922		M2.1	1.6E-2
53 – 93	keV	092530	092548	092602		8	HXT Y
8.8	GHz	0907.0	0909.0	0915.0	[U0.6 P0.9] U4 P8.8	1.78	
5	GHz	0912.0	0912.0	~0912.0		1.54	
3	GHz	0912.4	0915.0			1.34	
1.4	GHz	0908.0	0909.0	0909.0		1.48	
900	MHz	0909.0	0909.9			1.85	
600	MHz	0909.0	0909.4	0912.4		1.30	
410	MHz	0922.0	0922.0	0924.0		1.63	
245	MHz	0922.0	0924.0	0926.0		1.78	
DS II	190-2200	0907		0928		2	
DS I	90-180	~0906		~0955	N	1	
DS III	45-150	~0910		~0957	N	1	
DS III	30-220	0933		0937	GG	2	
DS DCIM	800-2000	0907		0910	GG	2	
DS UNCLF	200-380	0910		0936		1	
CME	WL	0943	0997km/s	-8.1km/s ²	040°	259°	



Particle event To(Ep>10 MeV) – 17d10^h

Tmax(Ep>10 MeV) – 19d22^h, Jmax (Ep>10 MeV) – 13/cm².s.sr

Duration of the event – 5 days

Quasimaximal energy of protons in the event – E_{qm} = 45 MeV


Sources: • solar flare 17d04^h48^m, M2.8/2N, S13E42, AR9704

Main X-ray burst 1–8 Å: onset – 17d04^h49^m, max – 17d05^h25^m, Φ = 0.1 J/m²



CME: 05^h30^m, V = 1379 km/s, Δφ = 360°, dA = 058°

▲ SC19d18^h15^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 November 17



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 November 17

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm².s.sr ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	10 ^h	20d01 ^h	139	5d	
EPS	>10	10 ^h	19d22 ^h	13	5d	
EPS	>30	10 ^h	19d19 ^h	0.25	4d	
EPS	>50	-	-	-	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	10 ^h	19d20 ^h	9932	6d	
MEPED	>0.8	10 ^h	19d20 ^h	4978	6d	
MEPED	>2.5	10 ^h	19d20 ^h	1299	5d	
MEPED	>6.9	10 ^h	19d20 ^h	156	5d	
MEPED	>16	10 ^h	19d20 ^h	1.85	4d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	10 ^h	20d01 ^h	3440	5d	
MKL	>14	10 ^h	20d01 ^h	0.72	4d	
MKL	>26	10 ^h	20d01 ^h	0.57	4d	
MKL	>50	10 ^h	20d01 ^h	0.21	4d	
ACE						
SIS	>10	-	20d01 ^h	13	5d	
SIS	>30	-	-	-	-	
SOHO						
EPHIN (INT)	>50	19d20 ^h	20d00 ^h	0.17	1d	

Differential fluxes of protons for the event of 2001 November 17

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Duration	Comments
SOHO						
LION	0.75-2	12 ^h	19d21 ^h	150	5d	
LION	2-6	11 ^h	19d21 ^h	23.6	5d	
EPHIN	4-8	09 ^h	20d01 ^h	95.4	5d	
EPHIN	8-25	09 ^h	20d01 ^h	2.8	5d	
EPHIN	25-41	09 ^h	20d00 ^h	0.006	5d	
EPHIN	41-53	- " -	- " -	- " -	- " -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 November 17


2001 November 17

•

AR9704

To event 408

H α	6563 Å	0448	0505	0938	S13E42	2N	FZE
DSF	6563 Å	0404		0435	S10E47	13°	
1 – 12	keV	0449	0525	0611		M2.8	1.0E-1
53 – 93	keV	<050656	~050706	053910		8	HXT Y
15.4	GHz	0447.0	0533.0	0630.0		1.89	
8.8	GHz	0446.0	0459.0	0614.0		1.95	
5	GHz	0446.0	0459.0	0651.0		2.08	
2.7	GHz	0449.0	0459.0	0653.0	U1.4 P2.7	2.18	
1.4	GHz	0447.0	0449.0	0449.0		1.36	
610	MHz	0512.0	0533.0	0547.0		2.36	
410	MHz	0447.0	0450.0	0454.0		2.00	
245	MHz	0451.0	0455.0	0458.0		2.30	
DS II	45-145	0450		0455		1	
DS II	25-55	0500		0507		3	
DS IV	57-500	0444		0555		1	
DS IV	25-180	0455		1025		2	
DS III	57-800	0448		0553	S	2	
CME	WL	0530	1379 km/s	-22.5 km/s ²	360°	058°	



Particle event: To(Ep>10 MeV) – 22d21^h

Tmax₁(Ep>10 MeV) – 23d10^h, Jmax₁(Ep>10 MeV) – $2.7 \cdot 10^3 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Tmax₂(Ep>10 MeV) – 24d06^h, Jmax₂(Ep>10 MeV) – $1.1 \cdot 10^4 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 4 days

Quasimaximal energy of protons in the event – Eqm₁ = 390 MeV

– Eqm₂ = 350 MeV

Sources: ● solar flare 22d<22^h09^m, M9.9/3B, S13W38, AR9704


Ø solar flare 22d20^h18^m, M3.8/2B, S26W68, AR9704

Main X-ray burst 1-8 Å: onset – 22d22^h32^m, max – 22d23^h30^m, $\Phi = 0.31 \text{ J/m}^2$



CME: 22d23^h30^m, V = 1437 km/s, $\Delta\phi = 360^\circ$, dA = 341^o

▲ SC 24d05^h56^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 November 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 November 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	21 ^h	23d11 ^h /24d06 ^h	3.6·10 ³ /2.8·10 ⁴	4d	
EPS	>10	21 ^h	23d10 ^h /24d06 ^h	2.7·10 ³ /1.1·10 ⁴	4d	
EPS	>30	21 ^h	23d10 ^h /24d05 ^h	624/923	3d	
EPS	>50	21 ^h	23d10 ^h /24d05 ^h	137/145	3d	
EPS	>60	21 ^h	23d10 ^h /24d05 ^h	43/46	2.5d	
EPS	>100	21 ^h	23d10 ^h /24d02 ^h	4/5.6	2d	
POES-16						
MEPED	>0.24	21 ^h	23d07 ^h /24d07 ^h	8.3·10 ³ /3.6·10 ⁵	4d	
MEPED	>0.8	21 ^h	23d07 ^h /24d07 ^h	8·10 ³ /1.5·10 ⁵	4d	
MEPED	>2.5	21 ^h	23d07 ^h /24d07 ^h	7.3·10 ³ /6.7·10 ⁴	4d	
MEPED	>6.9	21 ^h	23d07 ^h /24d07 ^h	5.3·10 ³ /2.1·10 ⁴	4d	
MEPED	>16	21 ^h	23d07 ^h /24d07 ^h	2.3·10 ³ /4.7·10 ³	3d	
MEPED	>36	21 ^h	23d07 ^h /24d07 ^h	550/458	3d	
MEPED	>70	21 ^h	23d07 ^h /24d07 ^h	13.1/12	2d	
MEPED	>140	21 ^h	23d07 ^h /24d07 ^h	0.67/0.67	2d	
CORONAS-F						
MKL	> 1	21 ^h	23d11 ^h /24d06 ^h	2.9·10 ³ /4.3·10 ⁴	4d	
MKL	>14	21 ^h	23d11 ^h /24d06 ^h	920/1.6·10 ³	4d	
MKL	>26	21 ^h	23d11 ^h /24d06 ^h	520/595	4d	
MKL	>50	21 ^h	23d11 ^h /24d06 ^h	150/121	4d	
ACE						
SIS	>10	21 ^h	23d11 ^h /24d06 ^h	2.7·10 ³ /3.3·10 ³	4d	
SIS	>30	21 ^h	23d11 ^h /24d06 ^h	575/688	3d	

SOHO						
EPHIN (INT)	>50	21 ^h	23d11 ^h /24d05 ^h	68.7/108	3.2d	
BALLOONS						
Mu	>150		23d(10 ^h -11 ^h)/ -	1.3/ -		
Mu	>323		23d(10 ^h -11 ^h)/ -	0.45/ -		
Mu	>633		23d(10 ^h -11 ^h)/ -	0.14/ -		

Differential fluxes of protons for the event of 2001 November 22

S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
SOHO						
LION	0.75-2	23d01 ^h	23d11 ^h /24d05 ^h	73.5/7560	8d	
LION	2-6	23d01 ^h	23d11 ^h /24d05 ^h	50.4/3210	8d	
EPHIN	4-8	23 ^h	23d11 ^h /24d05 ^h	132/1170	8d	
EPHIN	8-25	22 ^h	23d11 ^h /24d05 ^h	73.5/200	8d	
EPHIN	25-41	21 ^h	23d11 ^h /24d05 ^h	12.7/7.4	8d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:



Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Lario D, A. Aran, R.B. Decker, 2009
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 November 22


2001 November 22 • AR9704 To event 409

Hα	6563Å	<2209	2345	0401	S13W38	3B	FHZE
DSF	6563Å	~2048		~1135	S18W36	17°	
1 – 12	keV	2232	2330	2406		M9.9	3.1E-1
53 – 93	keV	222939	225749	>000351		14	HXT Y
15.4	GHz	2204.0	2204.0	2205.0		1.49	
8.8	GHz	2203.0	2203.0	0000.0		2.08	
5	GHz	2201.0	2203.0	0000.0		2.26	
2.8	GHz	2200.0	2204.0	2209.0		3.64	
1.4	GHz	2158.0	2204.0	0024.0	U0.5 P1.4	4.63	
610	MHz	2201.0	2202.0	2204.0		2.32	
500	MHz	2201.0	2202.0	2203.0		2.22	
410	MHz	2202.0	2202.0	2203.0		2.53	
245	MHz	2201.0	2202.0	2202.0		3.43	
200	MHz	2202.0	2202.0	2203.0		2.48	
DS II	25-116	2231		2241		1	
DS IV	25-180	2235		0012		2	
DS IV	57-1300	2245		>2400	FS	2	
DS III	140-700	2201		2202	B	2	
DS III	57-1300	2253		2346	N	2	
DS CONT	74-180	2235		2320		1	

15.4	GHz	2203.0	2300.0	0024.0		2.51	
8.8	GHz	2202.0	2305.0	0055.0		2.90	
5	GHz	2202.0	2305.0	0055.0		3.04	
2.7	GHz	2159.0	2257.0	0024.0	U0.6 P2.7	3.99	
610	MHz	2201.0	~2255.0	0021.0		3.08	
410	MHz	2201.0	~2257.0	0024.0		3.36	
245	MHz	2201.0	~2315.0	0024.0		3.46	
200	MHz	2232.0	2317.0	2340.0		2.94	
610	MHz	2201.0	2357.0	2358.0		3.54	
410	MHz	2202.0	2355.0	2357.0		3.49	
DS IV	57-300	<0000		0010	FS	1	
DS I	60-170	0010		0202	S	1	
CME	WL	2330	1437 km/s	-12.9 km/s ²	360°	349°	



2001	November 22	Ø			AR9698	To event 409	
H α	6563 Å	2022	2026	>2126	S26W68	2B	FHZ
1 – 12	keV	2018	2036	2052		M3.8	5.1E-2
53 – 93	keV	<204429	204433	204639		8	HXT Y
9.5	GHz	2022.0	2027.4	2037.5		2.72	
5	GHz	2023.0	2027.0	0000.0		2.76	
2.7	GHz	2023.0	2027.0	2039.0	U1.4 P2.7	2.80	
1.4	GHz	2023.0	2023.0	2037.0		2.11	
245	MHz	2038.0	2039.0	2039.0		2.41	
DS II	25-180	2022		2047		3	
DS III	25-180	2021		2056	N	1	
DS III	57-200	2023		2026	G	3	
DS UNCLF	57-100	2038		2044		3	
CME	WL	2030	1443 km/s	-43.3 km/s ²	360°	221°	



Particle event: To($E_p > 10$ MeV) – 26d05^h

Tmax($E_p > 10$ MeV) – 26d11^h, Jmax ($E_p > 10$ MeV) – 336/cm².s.sr

Duration of the event – 3 days


Quasimaximal energy of protons in the event – $E_{qm} = 800$ MeV

Sources: • solar flare 26d04^h32^m, M7.1/1B, N08W54, AR9742



Main X-ray burst 1-8 Å: onset – 26d04^h32^m, max – 26d05^h40^m, $\Phi = 0.34$ J/m²

CME: 26d05^h30^m, $V = 1446$ km/s, $\Delta\phi = 212^\circ$, $dA = 266^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 December 26



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 December 26

S/c, Instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	05 ^h	11 ^h	432	3d	
EPS	>10	05 ^h	11 ^h	340	3d	
EPS	>30	05 ^h	07 ^h	154	3d	
EPS	>50	05 ^h	07 ^h	106	3d	
EPS	>60	05 ^h	07 ^h	86	2.5d	
EPS	>100	05 ^h	07 ^h	40	2d	
POES 16						
MEPED	>0.24	05 ^h	07 ^h	3350	3d	
MEPED	>0.8	05 ^h	07 ^h	2150	3d	
MEPED	>2.5	05 ^h	07 ^h	1500	3d	
MEPED	>6.9	05 ^h	07 ^h	1300	3d	
MEPED	>16	05 ^h	07 ^h	563	3d	
MEPED	>36	05 ^h	07 ^h	448	3d	
MEPED	>70	05 ^h	07 ^h	94	2d	
MEPED	>140	05 ^h	07 ^h	27	2d	
ACE						
SIS	>10	05 ^h	09 ^h	987	2.5d	
SIS	>30	05 ^h	07 ^h	565	2d	
SOHO						
EPHIN (INT)	>50	05 ^h	07 ^h	84	2d	

BALLOONS						
Mu	>139		07 ^h 07 ^m -08 ^h 44 ^m	9.9		
Mu	>323		07 ^h 07 ^m -08 ^h 44 ^m	1.31		
Mu	>633		07 ^h 07 ^m -08 ^h 44 ^m	0.23		
NM						
Network	>500		07 ^h	0.74		

Differential fluxes of protons for the event of 2001 December 26

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
SOHO						
LION	0.75-2	05 ^h	12 ^h	29.4	3d	
LION	2-6	05 ^h	12 ^h	17.3	3d	
EPHIN	4-8	05 ^h	11 ^h	107	3d	
EPHIN	8-25	05 ^h	11 ^h	28.2	3d	
EPHIN	25-41	05 ^h	09 ^h	4.7	2.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Struminsky A.B, 2003.
 Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
 Miroshnichenko L.I. and J. Perez-Peraza, 2008.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 December 26

2001 December 26		●		AR9742		To event 410	
H α		0432	0514	0823	N08W54	1B	EPT
1 – 12	keV	0432	0540	0647		M7.1	3.4E-01
15.4	GHz	0453.0	0507.0	0642.0		3.26	
8.8	GHz	0439.0	0512.0	0706.0		3.58	
5	GHz	0435.0	0512.0	0700.0	P5 - 8.8	3.59	
2.8	GHz	0430.0	0514.0	0644.0		3.42	
1.4	GHz	0447.0	0514.0	0648.0		2.81	
610	MHz	0501.0	0624.0	0644.0		2.48	
410	MHz	0455.0	0557.0	0643.0		2.26	
200	MHz	0446.0	0453.0	0659.0		2.02	

DS II	25-180	0502		0519		3	
DS IV	57-1000	0501		>0810		2	
DS IV	25-180	0520		1049		3	
DS I	60-270	<0652		~1200	N	2	
DS III	57-250	0455		0511	G	2	
DS III	25-72	0557		0557		3	
CME	WL	0530	1446 km/s	-39.9 km/s ²	212°	266°	



Particle event: To(Ep>10 MeV) – 29d05^h

Tmax(Ep>10 MeV) – 29d08^h, Jmax (Ep>10 MeV) – 40/cm².s.sr

Duration of the event – 1.5 days

Quasimaximal energy of protons in the event – Eqm = 195 MeV

Sources: ■ solar flare 28d20^h02^m, X3.4/..., s26e90, AR9767*

○ solar flare 28d09^h38^m, M9.3/..., S08W88, AR9748


Main X-ray burst 1-8 Å: onset – 28d20^h02^m, max – 28d20^h45^m, $\Phi = 1.3 \text{ J/m}^2$

CME: 28d20^h30^m, V=2216 km/s, $\Delta\phi = 360^\circ$, dA = 115°;



▲ SC 29d05^h38^m

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 December 29



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 December 29

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	05 ^h	08 ^h	115	1.5d	
EPS	>10	05 ^h	08 ^h	40	1.5d	
EPS	>30	05 ^h	08 ^h	2	1.5d	
EPS	>50	05 ^h	08 ^h	0.44	1d	
EPS	>60	05 ^h	08 ^h	0.24	1d	
EPS	>100	05 ^h	08 ^h	0.05	1d	
POES-16						
MEPED	>0.24	04 ^h	09 ^h	1200	1.5d	
MEPED	>0.8	04 ^h	09 ^h	640	1.5d	
MEPED	>2.5	04 ^h	09 ^h	274	1.5d	
MEPED	>6.9	04 ^h	09 ^h	106	1.5d	
MEPED	>16	04 ^h	09 ^h	10.7	1.5d	
MEPED	>36	04 ^h	09 ^h	1.4	1d	
MEPED	>70	04 ^h	09 ^h	0.18	1d	
MEPED	>140	04 ^h	09 ^h	0.018	1d	
ACE						
SIS	>10	03 ^h	08 ^h	37.7	1.5d	
SIS	>30	04 ^h	08 ^h	2.7	1d	
SOHO						
EPHIN (INT)	>50	01 ^h	07 ^h	0.08	1d	

Differential fluxes of protons for the event of 2001 December 29



S/c, instruments	ΔE, MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
SOHO						
LION	0.75-2	02 ^h	08 ^h	10.6	1.5d	
LION	2-6	02 ^h	08 ^h	2.7	1.5d	

EPHIN	4-8	02 ^h	08 ^h	16.8	1d	
EPHIN	8-25	01 ^h	07 ^h	1.9	1d	
EPHIN	25-41	01 ^h	07 ^h	0.07	1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2001 December 29**


2001 December 28		■			AR9767		To event 411	
H α	6563 Å	No Flare			s26e90*			
1 – 12	keV	2002	2045	2132		X3.4	1.3E00	
8.8	GHz	1950.0	2004.0	2021.0		3.28		
5	GHz	1946.0	2004.0	2024.0	U0.6 P5	3.43		
2.7	GHz	1942.0	2005.0	2024.0		2.88		
1.4	GHz	2011.0	2012.0	2015.0		1.76		
610	MHz	2004.0	2006.0	2007.0		1.60		
410	MHz	2003.0	2003.0	~2003.0		1.74		
245	MHz	1959.0	2000.0	2000.0		2.85		
DS II	25-180	2003		2006		2		
DS III	57-130	2005		2014	GG	3		
DS III	25-180	2007		2040	N	2		
15.4	GHz	1944.0	2017.0	2105.0		3.15		
8.8	GHz	1957.0	2015.0	2033.0		3.04		
5	GHz	1949.0	2015.0	2105.0	P5	3.46		
2.7	GHz	1946.0	2015.0	2054.0		3.20		
1.4	GHz	1949.0	2023.0	2045.0		2.58		
DS II	57-130	2018		2037	SH	3		
DS III	57-110	2015		2019	GG	3		
CME	WL	2030	2216 km/s	6.9 km/s ²	360°	115°		

* – probable localization of the flare event



2001	December 29	Ø		AR9748	To event 411		
Hα	6563 Å	0941	0942	0952	S08W88	SF	
1 – 12	keV	0938	0945	1006		M9.3	1.1E-1
15.4	GHz	0939.0	0939.0	0945.0		1.88	
8.8	GHz	0939.0	0943.0	0951.0	U0.9 P8.8	2.34	
5	GHz	0939.0	0941.0	0951.0		2.15	
2.7	GHz	0939.0	0939.0	0943.0		2.00	
1.4	GHz	0939.0	0940.0	0943.0		1.76	

900	MHz	0939.3	0942.3			1.65	
600	MHz	0939.2	0942.0			1.88	
410	MHz	0940.0	0942.0	0948.0		2.56	
245	MHz	0939.0	0944.0	0946.0		3.90	
204	MHz	0943.2	0944.2	0947.5		4.11	
DS II	40-280	0943		1015		3	
DS IV	40-450	0939		1020		2	
DS III	25-180	0939		0943		3	
DS III	40-270	0943		0949	GG	2	
DS III	30-155	0950		0957	N	2	
DS V	45-165	0939		0940	G	2	
DS DCIM	2000-4500	0939		0944	G	1	
DS DCIM	800-2000	0939		0945	GG	1	
DS III/V	40-700	0939		0942	G,U	3	
CME	WL	0954	0634 km/s	-10.8 km/s ²	150°	271°	



Particle event: To(Ep>10 MeV) – 30d20^h

Tmax₁(Ep>10 MeV) – 31d02^h, Jmax₁(Ep>10 MeV) – 25.5 /cm².s.sr

Tmax₂(Ep>10 MeV) – 31d16^h, Jmax₂(Ep>10 MeV) – 75 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 190 MeV

– Eqm₂ = 170 MeV

Sources: ☐ solar flare 29d19^h50^m, M1.8/, s08w90, AR9748


Main X-ray burst 1-8 Å: onset – 29d19^h50^m, max – 29d21^h27^m, Φ = 0.19 J/m²

CME: 29d20^h30^m, V=0819 km/s, Δφ = 211°; dA = 281°;



ΔSC 30d20^h29^m

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2001 December 30



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 December 30

S/c, instruments	Ep, MeV	To	Tmax	Jmax, /cm ² .s.sr	Dura- tion	Comments
GOES-10						
EPS	>5	20 ^h	31d02 ^h /31d16 ^h	63.2/177	3d	
EPS	>10	20 ^h	31d02 ^h /31d16 ^h	25.5/75	3d	
EPS	>30	20 ^h	31d02 ^h /31d16 ^h	2.2/4.1	3d	
EPS	>50	20 ^h	31d02 ^h /31d16 ^h	0.4/0.5	3d	
EPS	>60	20 ^h	31d02 ^h /31d16 ^h	0.17/0.23	3d	
EPS	>100	20 ^h	31d02 ^h /31d16 ^h	0.05 /0.1	3d	
POES-16						
MEPED	>0.24	20 ^h	31d02 ^h /31d16 ^h	1830/3.6·10 ³	3d	
MEPED	>0.8	20 ^h	31d02 ^h /31d16 ^h	625/1.5·10 ³	3d	
MEPED	>2.5	20 ^h	31d02 ^h /31d16 ^h	220/610	3d	
MEPED	>6.9	20 ^h	31d02 ^h /31d16 ^h	110/194	3d	
MEPED	>16	20 ^h	31d02 ^h /31d16 ^h	13/32.6	3d	
MEPED	>36	20 ^h	31d02 ^h /31d16 ^h	2.5/2.8	3d	
MEPED	>70	20 ^h	31d02 ^h /31d16 ^h	0.25 /0.18	3d	
MEPED	>140	-	-	-	-	
ACE						
SIS	>10	20 ^h	31d00 ^h /31d16 ^h	24/63.4	3d	
SIS	>30	20 ^h	31d00 ^h /31d16 ^h	2.3 /3.6	3d	
SOHO						
EPHIN (INT)	>50	-	31d00 ^h / -	0.06/ -	-	

Differential fluxes of protons for the event of 2001 December 30

S/c, instruments	ΔE , MeV	To	Tmax	Jmax. /cm ² .s.sr.MeV	Dura- tion	Comments
SOHO						
LION	0.75-2	22 ^h	31d02 ^h /31d15 ^h	6.7/25.7	3d	
LION	2-6	20 ^h	31d02 ^h /31d15 ^h	1.7/6.4	3d	
EPHIN	4-8	21 ^h	31d02 ^h /31d15 ^h	8.4/27.6	3d	
EPHIN	8-25	20 ^h	31d02 ^h /31d14 ^h	1.3/3.9	3d	
EPHIN	25-41	20 ^h	31d02 ^h /31d14 ^h	0.08/0.15	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2001 December 30

2001 December 29		☐		AR9767		To event 412	
H α	6563 Å	No	Flare	Patrol	s08w90*		
1 – 12	keV	1950	2127	2355		M1.8	1.9E-01
410	MHz	2007.0	2009.0	2010.0		2.30	
DS I	70-170	2013		2228	S	1	
DS III	57-180	2140		2148	G	1	
DS CONT	78-180	2158		0538		1	
CME	WL	2030	0819 km/s	9.3 km/s ²	211°	281°	

* – probable localization of the flare event

Events in 2002

				Page
1.	Event 2002.01.10 – (2002-010)	№ 413	413
2.	Event 2002.01.15 – (2002-015)	№ 414	417
3.	Event 2002.02.20 – (2002-051)	№ 415	421
4.	Event 2002.03.16 – (2002-075)	№ 416	426
5.	Event 2002.03.18 – (2002-077)	№ 417	432
6.	Event 2002.03.20 – (2002-079)	№ 418	437
7.	Event 2002.03.22 – (2002-081)	№ 419	441
8.	Event 2002.04.17 – (2002-107)	№ 420	445
9.	Event 2002.04.19 – (2002-109)	№ 421	450
10.	Event 2002.04.21 – (2002-111)	№ 422	454
11.	Event 2002.05.22 – (2002-142)	№ 423	459
12.	Event 2002.07.07 – (2002-188)	№ 424	463
13.	Event 2002.07.16 – (2002-197)	№ 425	467
14.	Event 2002.07.19 – (2002-200)	№ 426	474
15.	Event 2002.07.22 – (2002-203)	№ 427	479
16.	Event 2002.08.14 – (2002-226)	№ 428	485
17.	Event 2002.08.17 – (2002-229)	№ 429	490
18.	Event 2002.08.18 – (2002-230)	№ 430	495
19.	Event 2002.08.20 – (2002-232)	№ 431	500
20.	Event 2002.08.22 – (2002-234)	№ 432	505
21.	Event 2002.08.24 – (2002-236) – GLE 64	№ 433	510
22.	Event 2002.09.06 – (2002-249)	№ 434	515
23.	Event 2002.11.09 – (2002-313)	№ 435	520

Particle event: To($E_p > 10$ MeV) – 10d02^h

Tmax($E_p > 10$ MeV) – 11d01^h, Jmax($E_p > 10$ MeV) – 70 /cm².s.sr

Duration of the event – 3 days


Quasimaximal energy of protons in the event – $E_{qm} = 85$ MeV

Sources: ☉ solar flare 09d17^h42^m, M9.5/2B, N13W02, AR9773



Main X-ray burst 1-8 Å: onset – 09d17^h42^m, max – 09d18^h01^m, $\Phi = 0.091$ J/m²

CME: 09d18^h54^m, V = 0113 km/s, $\Delta\phi = 026^\circ$, dA = 353°

Particle fluxes and associated phenomena



Time profiles of the proton fluxes for the event of 2002 January 10



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum




Table of the fluxes integral of proton for the event 2002 January 10

S/c, instru-ments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura-tion	Comments
GOES 10						
EPS	>5	02 ^h	11d01 ^h	190	3d	
EPS	>10	02 ^h	11d01 ^h	70	3d	
EPS	>30	02 ^h	11d01 ^h	2.8	3d	
EPS	>50	-	11d00 ^h	0.27	3d	
EPS	>60	-	11d00 ^h	0.14	3d	
POES 16						
MEPED	>0.24	-	11d05 ^h	2080	3d	
MEPED	>0.8	-	11d05 ^h	970	3d	
MEPED	>2.5	-	11d04 ^h	410	3d	
MEPED	>6.9	-	11d03 ^h	250	3d	
MEPED	>16	-	11d02 ^h	23	3d	
MEPED	>36	-	11d02 ^h	2	3d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1	-	11d02 ^h	1020	3d	
MKL	>14	-	11d01 ^h	12	3d	
MKL	>26	-	11d01 ^h	7	3d	
MKL	>50	-	11d01 ^h	2.5	3d	
ACE						
SIS	>10	08 ^h	11d00 ^h	63	3d	
SIS	>30	08 ^h	11d00 ^h	2.6	3d	
SOHO						
EPHIN (INT)	>50	22 ^h	11d 04 ^h	0.2	1d	

Fluxes differential of proton for the event 2002 January 10

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Duration	Comments
SOHO						
LION	0.75-2	14 ^h	11d06 ^h	46.2	3d	
LION	2-6	14 ^h	11d06 ^h	10.4	3d	
EPHIN	4-8	14 ^h	11d05 ^h	60.6	3d	
EPHIN	8-25	14 ^h	11d05 ^h	6.7	2d	
EPHIN	25-41	09 ^h	11d00 ^h	0.16	2d	
ERHIN	41-53	- " -	- " -	- " -	- " -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 January 10


2002 January 09

⊙

AR9773

To event 413

H α		1744	1800	1900	N13W02	2B	FU
1 – 12	keV	1742	1801	1812		M9.5	9.1E-02
15.4	GHz	1757.0	1757.0	1758.0		2.79	
8.8	GHz	1751.0	1757.0	0000.0	U2.7 P8.8	2.91	
5	GHz	1751.0	1757.0	0000.0		2.77	
2.7	GHz	1753.0	1757.0	0000.0		2.40	
1.4	GHz	1751.0	1757.0	1759.0		2.80	
DS III	25-90	1805		1806		1	
CME	WL	1854	0113 km/s	2.3 km/s ²	026°	353°	



Particle event: To($E_p > 10$ MeV) – 15d07^h
 Tmax($E_p > 10$ MeV) – 15d18^h, Jmax($E_p > 10$ MeV) – 7.5 /cm².s.sr
 Duration of the event – 4 days
 Quasimaximal energy of protons in the event – Eqm = 80 MeV

Sources: ☐ solar flare 14d05^h29^m, M4.4/..., s23w90*, AR9767, 1.5d behind W-limb


○ solar flare 14d01^h48^m, 2N/M1.7, N05E44, AR9782

Main X-ray burst 1-8 Å: onset – 14d05^h29^m, max – 14d06^h27^m, $\Phi = 0.34$ J/m²



CME: 14d05^h35^m; V = 1492 km/s; $\Delta\phi = 360^\circ$; dA = 246°

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 January 15



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 January 15

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	07 ^h	19 ^h	16.2	4d	
EPS	>10	07 ^h	18 ^h	7.5	4d	
EPS	>30	07 ^h	18 ^h	1.0	3d	
EPS	>50	07 ^h	15 ^h	0.22	2d	
EPS	>60	-	15 ^h	0.12	2d	
POES-16						
MEPED	>0.24	-	22 ^h	310	4d	
MEPED	>0.8	-	22 ^h	250	4d	
MEPED	>2.5	-	22 ^h	105	4d	
MEPED	>6.9	-	22 ^h	52	3d	
MEPED	>16	-	22 ^h	5.7	2d	
MEPED	>36	-	21 ^h	1.2	2d	
MEPED	>70	-	20 ^h	0.2	4d	
ACE						
SIS	>10	14d23 ^h	22 ^h	14.5	5d	
SIS	>30	14d11 ^h	18 ^h	2	5d	
SOHO						
EPHIN (INT)	>50	14d08 ^h	16 ^h	0.06	2d	

Differential fluxes of protons for the event of 2002 January 15

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Duration	Comments
SOHO						
LION	0.75-2	14 ^h	21 ^h	9.1	5d	
LION	2-6	11 ^h	21 ^h	1.8	5d	
EPHIN	4-8	13 ^h	20 ^h	6.8	5d	
EPHIN	8-25	13 ^h	20 ^h	0.7	5d	
EPHIN	25-41	14d09 ^h	20 ^h	0.04	5d	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2002 January 15**


2002 January 14 ☐ AR9767 To event 414

H α	6563 Å	No Flare			s23w90*		
1 – 12	keV	0529	0627	0825		M4.4	3.4E-01
610	MHz	0555.0	0555.0	~0555.0		1.82	
500	MHz	0518.0	0521.0	0532.0		1.70	
245	MHz	0537.0	0540.0	0540.0		1.85	
200	MHz	0524.0	0535.0	0548.0		1.90	
DS II	85-180	0608		0611	SH	1	
DS III	30-170	0536		0643	N	1	
CME	WL	0535	1492km/s	52.3km/s ²	360°	246°	

* – probable localization of the flare event

2002 January 14 Ø AR9782 To event 414

H α		0148	0155	0320	N05E44	2N	FH
1 – 12	keV	0152	0156	0203		M1.7	8.4E-03
15.4	GHz	0154.0	0155.0	0158.0	[U1.4] / 15.4	2.54	
8.8	GHz	0153.0	0155.0	0158.0		2.45	
5	GHz	0153.0	0155.0	0159.0		2.41	
2.8	GHz	0151.0	0155.4	0210.0		2.27	
1.4	GHz	0155.0	0155.0	~0155.0		1.46	
410	MHz	0153.0	0153.0	~0153.0		1.89	
200	MHz	0153.0	0153.0	0154.0		1.54	
CME	WL						gap



Particle event: To($E_p > 10$ MeV) – 20d07^h

Tmax($E_p > 10$ MeV) – 20d08^h, Jmax ($E_p > 10$ MeV) – 3.3 /cm².s.sr

Duration of the event – 2 days


Quasimaximal energy of protons in the event – $E_{qm} = 145$ MeV

Sources: ● solar flare 20d05^h52^m, M5.1/1N, N12W72, AR9825



Main X-ray burst 1-8 Å: onset – 20d05^h52^m, max – 20d06^h12^m, $\Phi = 0.022$ J/m²

CME: 20d06^h30^m, $V = 952$ km/s, $\Delta\phi = 360^\circ$, $dA = 263^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 February 20



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 February 20

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	07 ^h	08 ^h	5.8	2d	
EPS	>10	07 ^h	08 ^h	3.3	2d	
EPS	>30	04 ^h	07 ^h	0.74	1d	
EPS	>50	02 ^h	07 ^h	0.21	1d	
EPS	>60	02 ^h	07 ^h	0.12	1d	
EPS	>100	02 ^h	07 ^h	0.03	1d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	07 ^h	10	1d	
MEPED	>36	-	07 ^h	3.3	1d	
MEPED	>70	-	07 ^h	0.2	1d	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	-	07 ^h	55	1d	
MKL	>14	-	07 ^h	2.5	1d	
MKL	>26	-	07 ^h	1.5	1d	
MKL	>50	-	07 ^h	1	1d	
ACE						
SIS	>10	06 ^h	07 ^h	8.6	3d	
SIS	>30	06 ^h	07 ^h	2.0	2d	
SOHO						
EPHIN (INT)	>50	02 ^h	06 ^h	0.13	0.5d	

Differential fluxes of protons for the event of 2002 February 20

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	08 ^h	11 ^h	2.1	9d	
LION	2-6	08 ^h	10 ^h	1	9d	
EPHIN	4-8	07 ^h	08 ^h	7.1	9d	
EPHIN	8-25	06 ^h	07 ^h	0.7	9d	
EPHIN	25-41	06 ^h	07 ^h	0.046	5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 February 20

2002 February 20


•

AR9825

To event 415

H α	6563 Å	0555	0611	0628	N12W72	1N	EF
1 – 12	keV	0552	0612	0616		M5.1	2.2E-2
12 – 25	keV	061624	061630	062912		984336	RHESSI
4–7	MeV						SONG F
15.4	GHz	0608.0	0611.0	0613.0	U1.4/15.4	2.98	
8.8	GHz	0608.0	0609.0	0612.0		2.89	
5	GHz	0608.0	0609.0	0611.0		2.36	
2.7	GHz	0606.0	0610.0	0611.0		2.00	
1.4	GHz	0608.0	0611.0	0613.0		1.68	
610	MHz	0609.0	0610.0	0611.0		2.04	
410	MHz	0608.0	0610.0	0611.0		2.18	
245	MHz	0608.0	0608.0	0610.0		2.97	
DS II	57-190	0615		0630	SH	2	
DS II	25-57	0620		0625		1	
DS IV	25-71	0621		0637		1	
DS III	25-800	0555		0558	G	3	
DS III	25-1000	0608		0615	G	3	
DS V	25-180	0555		0558		3	
DS V	25-180	0608		0610		1	

n°							Armenia(S)
CME	WL	0630	0952 km/s	-17.1 km/s^2	360°	263°	



Particle event: To(Ep>10 MeV) – 16d02^h

Tmax₁(Ep>10 MeV) – 16d13^h, Jmax₁(Ep>10 MeV) – 1.3 /cm².s.sr

Tmax₂(Ep>10 MeV) – 17d11^h, Jmax₂(Ep>10 MeV) – 1.1 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 70 MeV

– Eqm₂ = 65 MeV


Sources: • solar flare 15d22^h09^m, M2.2/1F, S08W03, AR9866

Ø solar flare 17d10^h11^m, M1.3/..., S22E14, AR9871



Main X-ray burst 1-8 Å: onset – 15d22^h09^m, max – 15d23^h10^m, Φ = 0.13 J/m²

CME: 15d23^h06^m, V=957 km/s, Δφ = 360°; dA = 309°

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 March 16



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 March 16

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	01 ^h	14 ^h /17d12 ^h	12.7/5.6	2d	
EPS	>10	02 ^h	13 ^h /17d11 ^h	1.3/1.1	2d	
EPS	>30	03 ^h	12 ^h /17d11 ^h	0.08/0.06	2d	
EPS	>50	-	12 ^h /17d11 ^h	0.01/ 0.02	2d	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	-	14 ^h /17d10 ^h	1220/1430	2d	
MEPED	>0.8	-	14 ^h /17d10 ^h	1010/1360	2d	
MEPED	>2.5	-	14 ^h /17d10 ^h	320/630	2d	
MEPED	>6.9	-	14 ^h /17d10 ^h	42/54	2d	
MEPED	>16	-	14 ^h /17d09 ^h	0.9/1.2	2d	
MEPED	>36	-	14 ^h /-	0.2/-	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	-	14 ^h /17d12 ^h	600/1980	2d	
MKL	>14	-	13 ^h /17d11 ^h	0.35/0.5	2d	
MKL	>26	-	12 ^h /17d11 ^h	0.25/0.1	2d	
MKL	>50	-	12 ^h /17d11 ^h	0.05/0.03	2d	

ACE						
SIS	>10	01 ^h	11 ^h /17d09 ^h	2.7/8	2d	
SIS	>30	01 ^h	11 ^h / -	0.2/ -	2d	
SOHO						
EPHIN (INT)	>50	-	- /17d08 ^h	- /0.08	2d	

Differential fluxes of protons for the event of 2002 March 16

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	04 ^h	11 ^h /17d09 ^h	50.5/102	2d	
LION	2-6	04 ^h	11 ^h /17d09 ^h	6.7/18.3	2d	
EPHIN	4-8	03 ^h	11 ^h /17d08 ^h	17.4/81.5	2d	
EPHIN	8-25	01 ^h	10 ^h /17d08 ^h	0.43/2.6	2d	
EPHIN	25-41	00 ^h	10 ^h /17d08 ^h	0.002/0.012	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 March 16

2002 March 15


•

AR9866

To event 416


H α	6563 Å	2220	2238	0041	S08W03	1F	FZ
1 – 12	keV	2209	2310	0042		M2.2	1.3E-01
25-50	keV	221856	224614	224700		4869744	RHESSI
12-25	keV	23:31:20	23:32:42	23:51:00		2764632	RHESSI
12-25	keV	23:51:00	23:51:58	00:26:28		2748672	RHESSI
6-12	keV	00:26:28	00:26:30	00:34:00		346968	RHESSI
15.4	GHz	2223.0	2223.0	~2223.0		1.79	
8.8	GHz	2223.0	2234.0	2304.0		2.08	
5	GHz	2222.0	2234.0	2304.0		2.20	
2.7	GHz	2220.0	2234.0	2303.0		2.28	
1.4	GHz	2220.0	2234.0	2300.0	U0.6 P1.4	2.38	
610	MHz	2220.0	2236.0	2303.0		2.28	
410	MHz	2220.0	2236.0	2306.0		2.34	
245	MHz	2306.0	2306.0	~2306.0		2.70	

DS IV	25-180	2213		0045		1	
DS IV	57-750	2218		2312	FS	1	
DS III	25-138	2208		0005	N	1	
DS III	57-200	<0000		>0800	S,C	2	
DS CONT	25-180	2219		0026		1	
CME	WL	2306	0957 km/s	-17.4 km/s ²	360°	309°	



2002		March 17		Ø		AR9871		To event 416	
Ha	6563 Å	No Flare Patrol			s20e22				
1 – 12	keV	1011	1011	1024		M1.3		6.0E-03	
50-100	keV	100840	101742	102112				RHESSI	
15.4	GHz	1014.0	1015.0	1022.0		2.76			
9.1	GHz	1013.0	1015.9	1030.0		2.86			
5	GHz	1014.0	1015.0	1022.0		2.41			
3	GHz	1012.9	1016.3			2.11			

900	MHz	1008.4	1014.7			1.73	
600	MHz	1008.4	1013.7	1032.3	U0.9 P9.1	2.08	
204	MHz	1013.8	1015.9	1016.8		2.45	
DS I	200-300	~1023		~1050	S,N	2	
DS III	100-300	1008		1016	GG,C	3	
DS III	110-400	1008		1009	G	2	
DS III	40-350	1013		1023	GG	3	
DS III	40-250	1035		1036	G	2	
DS III	40-300	1039		1039	G	1	
DS DCIM	300-4000	1008		1036	P	3	
CME	WL	1034	0989 km/s	-6.2km/s ²	187°	165°	



Particle event: To(Ep>10 MeV) – 18d00^h

Tmax₁(Ep>10 MeV) – 18d15^h, Jmax₁(Ep>10 MeV) – 14.5 /cm².s.sr

Tmax₂(Ep>10 MeV) – 19d06^h, Jmax₂(Ep>10 MeV) – 20 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Emax₁ = 155 MeV

– Emax₂ = 145 MeV

Sources: ☉ solar flare 17d19^h24^m, M4.0/SF, S22E16, AR9871


☾ solar flare 18d02^h16^m, M1.0/SN, S16E27, AR9871

Main X-ray burst 1-8 Å: onset – 17d19^h24^m, max – 17d19^h31^m, Φ = 0.011 J/m²



CME: 17d20^h06^m, V = 823 km/s, Δφ = 288°, dA = 276°

▲ SC 18d13^h23^m;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 March 18



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 March 18

S/c, instru- ments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES 10						
EPS	>5	00 ^h	15 ^h /19d06 ^h	170/110	2d	
EPS	>10	00 ^h	15 ^h /19d06 ^h	14.5/20	2d	
EPS	>30	00 ^h	17 ^h /19d07 ^h	1/1.1	2d	
EPS	>50	06 ^h	15 ^h /19d06 ^h	0.25/0.17	2d	
EPS	>60	06 ^h	14 ^h /19d06 ^h	0.15/0.1	2d	
EPS	>100	06 ^h	15 ^h /19d06 ^h	0.04/0.03	2d	
METEOR						
CBM	>5	08 ^h	18 ^h /19d07 ^h	3.4/9.8	3d	
CBM	>15	07 ^h	17 ^h /19d03 ^h	1.5/3.3	2.5d	
CBM	>25	07 ^h	17 ^h /19d07 ^h	0.5/0.8	2d	
CBM	>40	06 ^h	17 ^h /19d01 ^h	0.16/0.3	1.5d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES 16						
MEPED	>0.24	-	- /19d07 ^h	- /2530	2d	
MEPED	>0.8	-	16 ^h /19d07 ^h	15200 /910	2d	
MEPED	>2.5	-	16 ^h /19d07 ^h	1930 /490	2d	
MEPED	>6.9	-	16 ^h /19d07 ^h	70 /110	2d	
MEPED	>16	-	16 ^h /19d07 ^h	05./10/	2d	
MEPED	>36	-	16 ^h /19d06 ^h	1.4/0.9	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	15 ^h /19d09 ^h	8850/1030	2d	
MKL	>14	-	15 ^h /19d09 ^h	2.75/4.3	2d	
MKL	>26	-	15 ^h /19d09 ^h	2.1/3.2	2d	
MKL	>50	-	15 ^h /19d09 ^h	1.0/1.5	2d	
ACE						
SIS	>10	07 ^h	14 ^h /19d04 ^h	11/30	1.5d	
SIS	>30	05 ^h	14 ^h /19d04 ^h	1.2/1.5	1.5d	
SOHO						
EPHIN (INT)	>50	05 ^h	13 ^h / -	0.6/ -	1.5d	


Fluxes differential of proton for the event 2002 March 18

S/c, instru- ments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	04 ^h	13 ^h /19d07 ^h	290/42	2d	
LION	2-6	03 ^h	13 ^h /19d07 ^h	26.5/11	2d	
EPHIN	4-8	05 ^h	13 ^h /19d06 ^h	101/42	2d	
EPHIN	8-25	08 ^h	13 ^h /19d06 ^h	2.6/3.1	2d	
EPHIN	25-41	05 ^h	13 ^h /19d05 ^h	0.04/0.07	2d	
ERHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2002 March 18**

2002 March 17		☉		AR9871		To event 417	
H α	6563 Å	<1938	~1938	1948	S22E16	SF	
1 – 12	keV	1924	1931	1934		M4.0	1.1E-2
50-100	keV	192616	193014	193628		6799464	HESSI

15.4	GHz	1927.0	1928.0	1934.0		2.98	
8.8	GHz	1926.0	1928.0	1934.0	U1.4 P8.8	3.00	
5	GHz	1927.0	1928.0	1935.0		2.90	
2.8	GHz	1923.0	1929.0	~1933.0		2.54	
1.4	GHz	1927.0	1928.0	1934.0		1.97	
610	MHz	1928.0	1928.0	1929.0		2.46	
410	MHz	1928.0	1929.0	1929.0		2.26	
245	MHz	1927.0	1928.0	1934.0		2.43	
DS III	N	1926		2112	25-180	3	
DS V		1926		1931	25-180	2	
n°							Armenia(S)
CME		2006	0823 km/s	-7.0 km/s ²	153°	191°	



2002	March 18	Ø			AR9871	To event 417	
Hα	6563 Å	0222	0224	0233	S16E27	SN	E
1 – 12	keV	0216	0231	0400		M1.0	4.5E-02
6-12	keV	025912	030014	030436		60024	HESSI
6-12	keV	030436	030506	032248		166440	HESSI
CME		0254	0989 km/s	-2.9 km/s ²	360°	311°	

Particle event: To($E_p > 10$ MeV) – 20d13^h

Tmax($E_p > 10$ MeV) – 20d17^h, Jmax ($E_p > 10$ MeV) – 8 /cm².s.sr

Duration of the event – 2 days


Quasimaximal energy of protons in the event – Emax = 60 MeV

Sources: O solar flare 20d08^h08^m, C1.9/SF, S19W41, AR9873



Main X-ray burst 1-8 Å: onset – 20d08^h08^m, max – 20d08^h33^m, $\Phi = 0.0031$ J/m²

▲ SC 20d13^h29^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 March 20



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Fluxes integral of proton for the event 2002 March 20

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES 10						
EPS	>5	13 ^h	17 ^h /	60	2d	
EPS	>10	13 ^h	17 ^h	8	2d	
EPS	>30	13 ^h	15 ^h	0.16	2d	
EPS	>50	13 ^h	15 ^h	0.02	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
METEOR						
CBM	>5	-	16 ^h	2.15	2d	
CBM	>15	-	16 ^h	0.5	2d	
CBM	>25	-	-	-	-	
CBM	>40	-	-	-	-	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES 16						
MEPED	>0.24	-	16 ^h	10150	2d	
MEPED	>0.8	-	16 ^h	3070	2d	
MEPED	>2.5	-	16 ^h	1120	2d	
MEPED	>6.9	-	16 ^h	28	2d	
MEPED	>16	-	16 ^h	5	2d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

ACE						
SIS	>10	10 ^h	14 ^h	7.5	2d	
SIS	>30	12 ^h	14 ^h	0.15	-	
SOHO						
EPHIN (INT)	>50	10 ^h	15 ^h	0.025	1d	

Fluxes differential of proton for the event 2002 March 20

S/c, instru- ments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	11 ^h	15 ^h	98	2d	
LION	2-6	14 ^h	15 ^h	13.5	2d	
EPHIN	4-8	10 ^h	15 ^h	45.3	2d	
EPHIN	8-25	10 ^h	15 ^h	1.73	2d	
EPHIN	25-41	10 ^h	15 ^h	0.014	2d	
ERHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 March 20

2002	March 20	o			AR9873	To event 418	
H α	6563 Å	0823	0824	0827	S19W41	SF	
1 – 12	keV	0808	0833	0845		C1.9	3.1E-3
245	MHz	0808.0	0808.0	~0808.0		2.40	
DS III	GG	0808		0808	30-270	2	
245	MHz	0837.0	0838.0	0838.0		2.63	
DS III	G	0837		0838	30-270	2	

Particle event: To(Ep>10 MeV) – 22d12^h

Tmax₁(Ep>10 MeV) – 22d20^h, Jmax₁(Ep>10 MeV) – 1 /cm².s.sr

Tmax₂(Ep>10 MeV) – 23d13^h, Jmax₂(Ep>10 MeV) – 9 /cm².s.sr

Duration of the event – 4 days

Quasimaximal energy of protons in the event – Emax₁ = 45 MeV

– Emax₂ = 50 MeV


Sources: ■ solar flare 22d10^h12^m, M1.6/..., s10w90, AR9866

Main X-ray burst 1-8 Å: onset – 22d10^h12^m, max – 22d11^h14^m, Φ = 0.049 J/m²



CME: 22d11^h06^m; V = 1750 km/s; Δφ = 360°; dA = 259°

▲ SC 23d11^h37^m

Particle fluxes and associated phenomena



Time profiles of the proton fluxes for the event of 2002 March 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum




Table of the fluxes integral of proton for the event 2002 March 22

S/c, instru-ments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES 10						
EPS	>5	14 ^h	20 ^h /23d13 ^h	3.7/110	4d	
EPS	>10	14 ^h	20 ^h /23d13 ^h	1/9	4d	
EPS	>30	-	22 ^h /23d16 ^h	0.05/0.1	-	
EPS	>50	-	22 ^h /23d16 ^h	0.015/0.012	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES 16						
MEPED	>0.24	-	22 ^h /23d16 ^h	- /2730	4d	
MEPED	>0.8	-	22 ^h /23d16 ^h	290 /1730	4d	
MEPED	>2.5	-	22 ^h /23d16 ^h	190/510	4d	
MEPED	>6.9	-	22 ^h /23d16 ^h	15/40	4d	
MEPED	>16	-	22 ^h /23d16 ^h	0.5/1.4	4d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	-	22 ^h /23d16 ^h	470/1230	4d	
MKL	>14	-	-	-	-	
MKL	>26	-	-	-	-	
MKL	>50	-	-	-	-	
ACE						
SIS	>10	13 ^h	22 ^h /23d12 ^h	6.7/7.2	2d	
SIS	>30	10 ^h	-	-	-	

SOHO						
EPHIN (INT)	>50	-	-	-	-	

Fluxes differential of proton for the event 2002 March 22

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	15 ^h	21 ^h /23d13 ^h	29.5/80	4d	
LION	2-6	15 ^h	21 ^h /23d13 ^h	7.7/21	4d	
EPHIN	4-8	13 ^h	20 ^h /23d12 ^h	23/45.5	6d	
EPHIN	8-25	13 ^h	20 ^h /23d12 ^h	1/1.4	6d	
EPHIN	25-41	13 ^h	19 ^h /23d13 ^h	0.0016/0.0035	5d	
ERHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2002 March 22**

2002 March 22 ■ AR9866 To event 419

H α	6563 Å	No Flare Patrol			s10w90		
1 – 12	keV	1012	1114	1152		M1.6	4.9E-2
6-12	keV	1037	103730	103828		9600	HESSI
25-50	keV	104536	105838	105940		663384	HESSI
6-12	keV	113208	113250	113912		99912	HESSI
3	GHz	1047.5	~1100.0	1152.5		1.77	
204	MHz	1010.7	1010.9	1013.2		1.86	
DS II		1047		1049	40-220	2	
DS IV		1052		~1154	40-200	2	
DS I	GG	1048		1049	130-160	2	
DS III	G	1047		1048	40-210	2	
DS DCIM	G	1047		1114	2000-4500	1	
CME	1106	1750 km/s	-22.5 km/s ²	360°	259°	1106	

Particle event: To($E_p > 10$ MeV) – 17d10^h

Tmax($E_p > 10$ MeV) – 17d16^h, Jmax ($E_p > 10$ MeV) – 21 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Emax = 120 MeV


Sources: ● solar flare 17d07^h46^m, M2.6/2N, S14W36, AR9906

Main X-ray burst 1-8 Å: onset – 17d07^h46^m, max – 17d08^h24^m, $\Phi = 0.15$ J/m²



CME: 17d08^h26^m; V = 1240 km/s; $\Delta\phi = 360^\circ$; dA = 292^o

▲ SC 17d11^h09^m; ▲ SC 19d08^h36^m;

Particle fluxes and associated phenomena



Time profiles of the proton fluxes for the event of 2002 April 17



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum




Table of the fluxes integral of proton for the event 2002 April 17

S/c, instru- ments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES 10						
EPS	>5	10 ^h	16 ^h	80	2d	
EPS	>10	10 ^h	16 ^h	21	2d	
EPS	>30	10 ^h	16 ^h	1.2	2d	
EPS	>50	10 ^h	16 ^h	0.2	2d	
EPS	>60	10 ^h	16 ^h	0.07	1d	
EPS	>100	-	16 ^h	0.02	1d	
POES 16						
MEPED	>0.24	-	16 ^h	2070	2d	
MEPED	>0.8	-	16 ^h	630	2d	
MEPED	>2.5	-	16 ^h	310	2d	
MEPED	>6.9	-	16 ^h	60	2d	
MEPED	>16	-	16 ^h	6	1d	
MEPED	>36	-	16 ^h	0.7	1d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	-	16 ^h	980	2d	
MKL	>14	-	-	-	2d	
MKL	>26	-	16 ^h	0.36	2d	
MKL	>50	-	16 ^h	0.1	2d	
ACE						
SIS	>10	10 ^h	15 ^h	12	2d	
SIS	>30	10 ^h	15 ^h	1.1	1d	
SOHO						
EPHIN (INT)	>50	11 ^h	15 ^h	0.16	0.5d	


Fluxes differential of proton for the event 2002 April 17

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, $(\text{cm}^2.\text{s.sr.MeV})^{-1}$	Duration	Comments
SOHO						
LION	0.75-2	10 ^h	18 ^h	15.2	2d	
LION	2-6	10 ^h	18 ^h	2.8	2d	
EPHIN	4-8	10 ^h	16 ^h	11.5	2d	
EPHIN	8-25	10 ^h	16 ^h	1	2d	
EPHIN	25-41	10 ^h	16 ^h	0.03	2d	
ERHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 April 17

2002	April 17	●	AR9906	To event 420			
H α	6563 Å	0750	0815	>1141	S14W36	2N	FTZ
1 – 12	keV	0746	0824	0957		M2.6	1.5E-1
25-50	keV	081432	081438	081624		1125888	HESSI
15.4	GHz	0803.0	0857.0	1033.0		2.97	
8.8	GHz	0753.0	0857.0	1012.0		3.38	
5	GHz	0753.0	0857.0	1037.0	P0.6 [U1.4 P5]	3.62	
2.7	GHz	0746.0	0857.0	0000.0		3.61	
1.4	GHz	0751.0	0847.0	1045.0		3.56	
610	MHz	0753.0	0905.0	1041.0		4.30	
410	MHz	0757.0	0845.0	0953.0		3.96	
245	MHz	0801.0	0901.0	1041.0		3.65	
204	MHz	0832.0	0905.5	1056.1		3.05	

DS II		0754		0820	100-600	3	
DS II		0808		0823	25-79	2	
DS II		0845		0908	100-1400	3	
DS II		0945		1024	100-900	3	
DS IV		0756		~1100	40-800	3	
DS III	GG,RS	0749		0758	720-3300	3	
DS III	N	0826		1200	25-270	2	
DS CONT		0804		~0832	25-270	2	
DS CONT		~0840		~1032	25-270	2	
DS DCIM	P,C,S	0754		1052	150-4000	3	
DS UNCLF		0813		0816	25-70	2	
CME		0826	1240 km/s	-19.8 km/s ²	360°	292°	



Particle event: To(Ep>10 MeV) – 19d05^h

Tmax₁(Ep>10 MeV) – 19d09^h, Jmax₁(Ep>10 MeV) – 1 /cm².s.sr

Tmax₂(Ep>10 MeV) – 19d19^h, Jmax₂(Ep>10 MeV) – 2.7 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 45 MeV

– Eqm₂ = 50 MeV


Sources: ◇ Flare activity AR9906

○ solar flare 19d15^h16^m, C2.5/SF, S16W59, AR9906;



Main X-ray burst 1-8 Å: onset – 19d15^h16^m, max – 19d15^h21^m;

▲ SC 19d08^h36^m;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 April 19



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 April 19

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	05 ^h	09 ^h /19 ^h	14/31	2d	
EPS	>10	05 ^h	09 ^h /19 ^h	1/2.7	2d	
EPS	>30	05 ^h	10 ^h /19 ^h	0.04/0.05	2d	
EPS	>50	05 ^h	11 ^h /19 ^h	0.01/ -	2d	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	-	10 ^h /19 ^h	25070/3330	2d	
MEPED	>0.8	-	10 ^h /19 ^h	5040/1340	2d	
MEPED	>2.5	-	10 ^h /19 ^h	960/320	2d	
MEPED	>6.9	-	10 ^h /19 ^h	31.5/50	2d	
MEPED	>16	-	10 ^h /19 ^h	0.6/1.5	2d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	-	08 ^h /19 ^h	1960/930	2d	
MKL	>14	-	-	-	-	
MKL	>26	-	-	-	-	
MKL	>50	-	-	-	-	

ACE						
SIS	>10	05 ^h	08 ^h /19 ^h	4.2/2.5	2d	
SIS	>30	-	-	-	-	
SOHO						
EPHIN (INT)	>50	05 ^h	08 ^h / -	0.01/ -	-	

Differential fluxes of protons for the event of 2002 April 19

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	05 ^h	08 ^h /20 ^h	97/46	2d	
LION	2-6	05 ^h	08 ^h /20 ^h	10/5.8	2d	
EPHIN	4-8	05 ^h	09 ^h /19 ^h	10.1/12.3	2d	
EPHIN	8-25	05 ^h	09 ^h /19 ^h	0.31/0.32	2d	
EPHIN	25-41	05 ^h	08 ^h /18 ^h	0.0015/0.0003	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2002 April 19**

2002	April 19	○	AR9906	To event 421			
H α	6563 Å	1513	1520	1528	S16W59	SF	F
1 – 12	keV	1516	1521	1527		C2.5	1.4E-3

Particle event: To($E_p > 10$ MeV) – 21d01^h

$T_{\max 1}(E_p > 10$ MeV) – 21d03^h, $J_{\max 1}(E_p > 10$ MeV) – 915 /cm².s.sr *)

$T_{\max 2}(E_p > 10$ MeV) – 21d09^h, $J_{\max 2}(E_p > 10$ MeV) – 1730 /cm².s.sr *)

Duration of the event – 6 days

Quasimaximal energy of protons in the event – $E_{qm1} = 575$ MeV

– $E_{qm2} = 570$ MeV

*) Data from ACE (SIS)


Sources: ■ solar flare 21d00^h43^m, X1.5/1F, S14W84, AR9906

Main X-ray burst 1-8 Å: onset – 21d00^h43^m, max – 21d01^h51^m, $\Phi = 0.6$ J/m²

CME: 21d01^h27^m, $V = 2393$ km/s, $\Delta\phi = 360^\circ$, $dA = 282^\circ$



▲ SC 23d04^h48^m

Particle fluxes and associated phenomena




April - May 2002

Time profiles of the proton fluxes for the event of 2002 April 21



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 April 21

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	02 ^h	03 ^h /09 ^h	195/410	6d	
EPS	>10	01 ^h	03 ^h /09 ^h	190/365	6d	
EPS	>30	01 ^h	03 ^h /10 ^h	160/195	6d	
EPS	>50	01 ^h	03 ^h /10 ^h	95/103	5d	
EPS	>60	01 ^h	03 ^h /10 ^h	68/64	5d	
EPS	>100	01 ^h	03 ^h /10 ^h	24/16.5	5d	
METEOR						
CBM	>5	02 ^h	- /11 ^h	- /585	7d	
CBM	>15	02 ^h	- /11 ^h	- /454	6d	
CBM	>25	02 ^h	- /09 ^h	- /330	5d	
CBM	>40	02 ^h	- /09 ^h	- /190	4d	
BP	>90	02 ^h	- /09 ^h	- /12.3	3d	
ChD	>600	02 ^h	- /10 ^h	- /1.4	2d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	03 ^h /10 ^h	2030/ 3110	6d	
MEPED	>16	-	03 ^h /10 ^h	970/ 1620	6d	
MEPED	>36	-	03 ^h /10 ^h	290/ 510	5d	
MEPED	>70	-	03 ^h /10 ^h	110/ 55	5d	
MEPED	>140	-	03 ^h /10 ^h	10/3	5d	
CORONAS F						

MKL	>1.	-	03 ^h /12 ^h	2830/3330	6d	
MKL	>14	-	03 ^h /12 ^h	490/840	5d	
MKL	>26	-	03 ^h /12 ^h	220/510	5d	
MKL	>50	-	03 ^h /12 ^h	110/210	5d	
ACE						
SIS	>10	01 ^h	04 ^h /09 ^h	915/1730	6d	
SIS	>30	01 ^h	04 ^h /09 ^h	430/650	6d	
SOHO						
EPHIN (INT)	>50	01 ^h	04 ^h /08 ^h	116/107	4d	
BALLOONS						
Mu	>117	-	- /22d(09 ^h 42 ^m -10 ^h 20 ^m)	- /1.5	-	After second
Mu	>137	-	- /22d(09 ^h 42 ^m -10 ^h 20 ^m)	- /0.41	-	maximum
Mu	>169	-	- /22d(09 ^h 42 ^m -10 ^h 20 ^m)	- /0.09	-	

Differential fluxes of protons for the event of 2002 April 21

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV)) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	02 ^h	05 ^h /10 ^h	7.5/19.3	5d	
LION	2-6	02 ^h	05 ^h /10 ^h	1.2/1.7	5d	
EPHIN	4-8	02 ^h	05 ^h /08 ^h	32/53.2	5d	
EPHIN	8-25	02 ^h	05 ^h /08 ^h	22.2/32.4	5d	
EPHIN	25-41	01 ^h	05 ^h /08 ^h	7.5/9.15	5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:



Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 April 21

2002 April 21 ■ AR9906 To event 422

H α	6563 Å	0059	0131	0251	S14W84	1F	FY
1 – 12	keV	0043	0151	0238		X1.5	6.0E-1
12-25	keV	004008	004914	005116		277345	RHESSI
100-300	keV	005116	013202	013324		59430512	RHESSI
15.4	GHz	0110.0	0123.0	0252.0		3.15	
8.8	GHz	0058.0	0123.0	0318.0	P5 - 8.8	3.45	
5	GHz	0057.0	0123.0	0321.0		3.46	
2.8	GHz	0044.0	0131.0	0342.0		3.29	
610	MHz	0111.0	0121.0	0000.0		2.90	
410	MHz	0111.0	0121.0	0000.0		2.38	
200	MHz	0102.0	0131.0	0305.0		2.27	

DS II	57-130	0119		0130	SH	3	
DS II	25-80	0119		0126		3	
DS IV	30-220	0117		0154		3	
DS I	57-120	0134		0238	S,C	1	
DS III	57-240	0120		0150	S,C	1	
DS III	57-210	0130		0132	G	2	
DS CONT	300-900	0120		0124		1	
2.7	GHz	0058.0	0227.0	0328.0		3.26	
610	MHz	0145.0	0210.0	0305.0	P0.6	4.11	
410	MHz	0146.0	0220.0	0252.0		3.69	
245	MHz	0143.0	0227.0	0306.0		2.88	
DS IV		0136		0238	25-124	1	
DS IV		0148		0245	130-1800	3	
DS III	G	0210		0222	25-210	2	
^o n						Mauna Kea,Norikura	
CME	WL	0127	2393 km/s	-1.4 km/s ²	360°	282°	



Particle event: To(Ep>10 MeV) – 22d07^h

Tmax₁(Ep>10 MeV) – 23d10^h, Jmax₁(Ep>10 MeV) – 260 /cm².s.sr

Tmax₂(Ep>10 MeV) – 23d16^h, Jmax₂(Ep>10 MeV) – 87 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 125 MeV

– Eqm₂ = 175 MeV


Sources: ☉ solar flare 22d03^h18^m, C5.0/SF, S22W53, DSF;

Main X-ray burst 1-8 Å: onset – 22d03^h18^m, max – 22d03^h54^m, $\Phi = 0.025$ J/m²;



CME: 22d03^h50^m, V = 1557 km/s, $\Delta\phi = 360^\circ$, dA = 250°;

▲ SC 23d10^h51^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 May 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of May 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	07 ^h	23d11 ^h /23d16 ^h	2120/520	2d	
EPS	>10	07 ^h	23d10 ^h /23d16 ^h	260/87	2d	
EPS	>30	07 ^h	23d10 ^h /23d16 ^h	7.2/4.4	2d	
EPS	>50	07 ^h	23d10 ^h /23d16 ^h	0.8/0.67	2d	
EPS	>60	07 ^h	23d10 ^h /23d16 ^h	0.3/0.22	2d	
EPS	>100	07 ^h	23d10 ^h /23d16 ^h	0.04/0.012	2d	
METEOR						
CBM	>5	08 ^h	23d11 ^h / -	83/ -	3d	
CBM	>15	08 ^h	23d11 ^h / -	27/ -	3d	
CBM	>25	08 ^h	23d11 ^h / -	5.3/ -	3d	
CBM	>40	08 ^h	23d11 ^h / -	1/ -	2d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	

POES-16						
MEPED	>0.24	-	23d11 ^h /23d16 ^h	35280/ 17140	2d	
MEPED	>0.8	-	23d11 ^h /23d16 ^h	12310/ 6130	2d	
MEPED	>2.5	-	23d11 ^h /23d16 ^h	4240/1610	2d	
MEPED	>6.9	-	23d11 ^h /23d16 ^h	830/190	2d	
MEPED	>16	-	23d11 ^h /23d16 ^h	60/15	2d	
MEPED	>36	-	23d11 ^h /23d16 ^h	4.2/1.3	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	-	23d12 ^h /23d16 ^h	9970/4080	2d	
MKL	>14	-	23d12 ^h /23d16 ^h	12/7.4	2d	
MKL	>26	-	23d12 ^h /23d16 ^h	8.2/5.5	2d	
MKL	>50	-	23d12 ^h /23d16 ^h	3.14/2.85	2d	
ACE						
SIS	>10	06 ^h	23d10 ^h /23d15 ^h	320/55	3d	
SIS	>30	07 ^h	23d09 ^h /23d15 ^h	9.7/3.6	3d	
SOHO						
EPHIN (INT)	>50	07 ^h	23d09 ^h /23d15 ^h	1/0.34	1d	

Differential fluxes of protons for the event of 2002 May 22

S/c. instruments	ΔE. MeV	To	Tmax	Jmax. (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	07 ^h	23d10 ^h /23d16 ^h	730/270	3d	
LION	2-6	05 ^h	23d10 ^h /23d16 ^h	224/47	3d	
EPHIN	4-8	03 ^h	23d10 ^h /23d16 ^h	1390/179	3d	
EPHIN	8-25	02 ^h	23d09 ^h /23d15 ^h	61/7.3	3d	
EPHIN	25-41	02 ^h	23d09 ^h /23d16 ^h	0.56/0.12	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 May 22

2002	May 22	☉	AR	To event 423			
Hα	6563 Å	<0400	~0400I	0437	S22W53	SF	U
DSF	6563 Å	21 ^d >1929		22 ^d <1135	S30W34	43°	
1 – 12	keV	0318	0354	0502		C5.0	2.5E-2
6-12	keV	04:00:44	04:00:54	04:16:32		74256	HESSI
2.8	GHz	0323.0	0337.6	0408.0		1.26	
200	MHz	0329.0	0329.0	0330.0		1.54	
DS III	57-180	0320		0353	GG	2	
DS III	25-180	0348		0352		2	
DS CONT	25-180	0321		0353		1	
CME	WL	0350	1557 km/s	-10.4 km/s ²	360°	250°	

Particle event: To($E_p > 10$ MeV) – 07d13^h

Tmax($E_p > 10$ MeV) – 07d20^h, Jmax ($E_p > 10$ MeV) – 26 /cm².s.sr *)

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm} = 85$ MeV

*) Data from ACE (SIS)


Sources: ■ solar flare 07d11^h15^m, M1.0/..., s19w90*, AR10017

Main X-ray burst 1-8 Å: onset – 07d11^h15^m, max – 07d11^h43^m, $\Phi = 0.062$ J/m²



CME: 07d11^h30^m; $V = 1423$ km/s; $\Delta\phi = 228^\circ$; $dA = 260^\circ$

* Localization from EPL observtion

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 July 07



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 July 07

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	13 ^h	20 ^h	6.6	3d	
EPS	>10	13 ^h	20 ^h	5.2	3d	
EPS	>30	13 ^h	19 ^h	1.3	3d	
EPS	>50	13 ^h	21 ^h	0.27	3d	
EPS	>60	13 ^h	23 ^h	0.13	3d	
EPS	>100	-	-	-	-	
METEOR						
CBM	>5	12 ^h	19 ^h	14.7	3d	
CBM	>15	12 ^h	19 ^h	5.2	3d	
CBM	>25	12 ^h	21 ^h	1.2	2d	
CBM	>40	12 ^h	22 ^h	0.34	1d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	22 ^h	280	3d	
MEPED	>0.8	-	22 ^h	140	3d	
MEPED	>2.5	-	22 ^h	80	3d	
MEPED	>6.9	-	22 ^h	20	3d	
MEPED	>16	-	22 ^h	12	3d	
MEPED	>36	-	22 ^h	1	3d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
ACE						
SIS	>10	12 ^h	20 ^h	26.6	3d	
SIS	>30	12 ^h	20 ^h	3.3	3d	

SOHO						
EPHIN (INT)	>50	12 ^h	19 ^h	0.1	1d	

Differential fluxes of protons for the event of 2002 July 07

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	12 ^h	22 ^h	5.3	3d	
LION	2-6	12 ^h	22 ^h	1.2	3d	
EPHIN	4-8	12 ^h	21 ^h	5.9	3d	
EPHIN	8-25	12 ^h	21 ^h	1.3	3d	
EPHIN	25-41	12 ^h	20 ^h	0.1	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2002 July 07**

2002 July 07



AR10017

To event 424

H α	6563 Å	No Flare			s13w90*		
EPL	6563 Å	1039		1059	S13W90		
1 – 12	keV	1115	1143	1317		M1.0	6.2E-2
12-25	keV	112956	113050	113908		339912	RHESSI
12-25	keV	113908	113910	114956		83664	RHESSI
12-25	keV	121304	121314	121624		56784	RHESSI
12-25	keV	121624	121842	123516		247872	RHESSI
6-12	keV	123516	123518	123852		35808	RHESSI
204	MHz	1112.6	1114.7	1115.1		1.85	
DS I	130-320	~1110		~1145	S,N	1	
DS I	45-90	1122		1137	S	2	
DS III	40-350	1111		1119	GG	2	
DS III	45-90	1152		1158	N	1	
DS III	40-65	1243		1243	B	2	
CME	WL	1130	1423 km/s	22.0 km/s ²	228°	260°	

* Localization from EPL observtion.

Particle event: To(Ep>10 MeV) – 16d12^h

Tmax₁(Ep>10 MeV) – 16d22^h, Jmax₁(Ep>10 MeV) – 27 /cm².s.sr

Tmax₂(Ep>10 MeV) – 17d14^h, Jmax₂(Ep>10 MeV) – 85 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 125 MeV

– Eqm₂ = 135 MeV

Sources: • solar flare 15d19^h50^m, 3B/X3.0, N19W01, AR10030


Ø solar flare 17d06^h58^m, M8.5/1B, N20W16 AR10030

Main X-ray burst 1-8 Å: onset – 15d19^h59^m, max – 15d20^h08^m, Φ = 0.14 J/m²



CME: 15d20^h30^m; V = 1151 km/s; Δφ = 360°; dA = 035°

Δ SC 17d16^h04^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 July 16



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 July 16

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm2.s.sr)-1	Dura- tion	Comments
GOES-10						
EPS	>5	12 ^h	22 ^h /17d14 ^h	48.5/430	3d	
EPS	>10	12 ^h	22 ^h /17d14 ^h	27/85	3d	
EPS	>30	12 ^h	22 ^h /17d14 ^h	3/1.5	2d	
EPS	>50	12 ^h	22 ^h /17d15 ^h	0.64/0.3	2d	
EPS	>60	12 ^h	22 ^h /17d14 ^h	0.32/0.2	1d	
EPS	>100	13 ^h	22 ^h /17d15 ^h	0.03/0.06	1d	
METEOR						
CBM	>5	<22 ^h	22 ^h /17d15 ^h	18/28	3d	
CBM	>15	<22 ^h	22 ^h /17d15 ^h	6.5/5	3d	
CBM	>25	<22 ^h	22 ^h /17d15 ^h	2/0.9	2d	
CBM	>40	<22 ^h	23 ^h /17d14 ^h	0.6/0.3	1d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	22 ^h /17d17 ^h	265/31200	-	
MEPED	>0.8	-	22 ^h /17d17 ^h	210/8930	-	
MEPED	>2.5	-	21 ^h /17d17 ^h	110/2480	-	
MEPED	>6.9	-	20 ^h /17d16 ^h	50/350	-	
MEPED	>16	-	20 ^h /17d16 ^h	13/15	-	
MEPED	>36	-	19 ^h /17d16 ^h	2/1	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	12 ^h /17d16 ^h	108/11980	-	
MKL	>14	-	12 ^h /17d16 ^h	10/6.7	-	
MKL	>26	-	12 ^h /17d16 ^h	6.4/4	-	
MKL	>50	-	12 ^h /17d16 ^h	2.4/1.14	-	
ACE						
SIS	>10	12 ^h	22 ^h /17d14 ^h	26/107	2d	
SIS	>30	12 ^h	21 ^h /17d14 ^h	3.8/1.65	2d	
SOHO						
EPHIN (INT)	>50	14 ^h	20 ^h /17d15 ^h	0.1/0.4	2d	

Differential fluxes of protons for the event of 2002 July 16

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	08 ^h	22 ^h /17d16 ^h	5.2/470	2,5d	
LION	2-6	08 ^h	22 ^h /17d16 ^h	1.8/85	2,5d	
EPHIN	4-8	08 ^h	21 ^h /17d15 ^h	8.4/329	2,5d	
EPHIN	8-25	08 ^h	21 ^h /17d15 ^h	1.5/15	2,5d	
EPHIN	25-41	08 ^h	21 ^h /17d14 ^h	0.11/0.065	2,5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:


Kuwabara T., J.W. Bieber, J. Clem et al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 July 16

2002 July 15 • AR10030 To event 425




Hα	6563 Å	1950	2011	>2334	N19W01	3B	FU
1 – 12	keV	1959	2008	2014		X3.0	1.4E-1
1 – 12	keV	2103	2132	2148		M1.8	4.3E-2
25-50	keV	202512	202802	205040		6989876	RHESSI
25-50	keV	205040	211414	212740		5761680	RHESSI
15.4	GHz	2003.0	2004.0	2156.0	P0.4 U1.4/15	4.30	
8.8	GHz	2001.0	2004.0	2146.0		4.18	
5	GHz	1958.0	2004.0	2204.0		3.81	
2.7	GHz	1957.0	2004.0	2221.0		3.28	
1.4	GHz	2001.0	2004.0	2228.0		3.04	
610	MHz	2003.0	2005.0	2225.0		3.70	
410	MHz	2001.0	2003.0	2304.0		4.15	
200	MHz	2003.0	2004.0	2015.0		3.56	
DS IV	30-80	1955		2213		3	
DS IV	25-180	2023		2319		2	
DS III	25-600	1954		2030	G	3	

15.4	GHz	2002.0	2230.0	2234.0	U0.5/15	4.30	
8.8	GHz	2001.0	2141.0	2234.0		4.15	
5	GHz	1957.0	2142.0	2234.0		3.70	
2.7	GHz	1955.0	2142.0	2234.0		3.32	
1.4	GHz	1956.0	2142.0	2234.0		2.70	
500	MHz	2003.0	2113.0	2148.0		2.58	
245	MHz	2001.0	2129.0	2221.0		4.40	
DS IV	57-1500	<2110		2214	F,S	3	
DS III	57-180	2214		>2400	S,C	1	
CME	WL	2030	1151 km/s	-25.6 km/s ²	360°	035°	
CME	WL	2130	1300 km/s	-7.3 km/s ²	188°	045°	



2002	July 17	Ø			AR10030	To event 425	
H α	6563 Å	0701	0704	0753	N20W16	1B	FU
1 – 12	keV	0658	0713	0753		M8.5	5.3E-2
50-100	keV	065812	070650	071044		17337330	RHESSI
1300-4000	keV						SONG F

9.1	GHz	0658.5	0703.2	>0730.0	U1.4 / 9.1	3.02	
5	GHz	0700.0	0702.0	0716.0		3.00	
3	GHz	0700.2	0702.8	0715.9		2.90	
1.4	GHz	0700.0	0703.0	0714.0		2.34	
600	MHz	0700.1	0701.4	0725.5		2.68	
245	MHz	0657.0	~0702.0	0721.0		3.20	
DS II		0706		0710		2	
DS IV		0700		0722		3	
DS III	G	0700		0713	G	3	
DS V		0700		0706		2	
DS UNCLF	DC	0705		0708	DC	2	
15.4	GHz	0700.0	0712.0	0721.0		3.26	
8.8	GHz	0700.0	0712.0	0725.0	P0.6 U1.4 P8.8	3.28	
2.7	GHz	0708.0	0712.0	0713.0		2.11	
1.4	GHz	0711.0	0712.0	0713.0		1.95	
900	MHz	0658.0	0713.0			2.45	
600	MHz	0700.1	0711.8			2.87	
410	MHz	0701.0	0709.0	0721.0		2.82	
DS III	25-270	0709		0713	S,C,F	2	
DS III	25-115	0720		0721		2	
DS V	25-82	0720		0721		2	
DS UNCLF	190-270	0711		0712	C	2	
15.4	GHz	0742.0	0743.0	0748.0	U0.9 [P8.8-15]	3.23	
8.8	GHz	0741.0	0743.0	0750.0		3.23	
5	GHz	0740.0	0743.0	0751.0		3.08	
2.7	GHz	0741.0	0743.0	0744.0		2.79	
1.4	GHz	0741.0	0743.0	0750.0		2.64	
900	MHz	0740.3	0743.5	0755.3		~2.00	
500	MHz	0740.0	0743.0	0758.0		3.30	
204	MHz	0738.8	0742.5	0751.8		3.63	
DS II	F,H	0742		0758	F,H	3	
DS II		0755		0802		2	
DS IV		~0756		~1055		2	
DS I	GG,DC	0744		0745	GG,DC	2	
DS III	G	0740		0757	G	3	
DS V		0741		0744		3	
DS UNCLF	DC	0744		0746	DC	2	
CME	WL	0731	0716 km/s	-17.8 km/s ²	177°	032°	



Particle event: To($E_p > 10$ MeV) – 19d05^h

Tmax($E_p > 10$ MeV) – 19d11^h, Jmax ($E_p > 10$ MeV) – $3.6 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm} = 215$ MeV

Sources: ☉ solar flare 18d07^h24^m, X1.8/2B, N19W33 AR10030;


◊ flare activity of AR10039, 3d behind E-limb

Main X-ray burst 1-8 Å: onset – 18d07^h24^m, max – 18d07^h44^m, $\Phi = 0.0056 \text{ J/m}^2$



CME: 18d08^h06^m, $V = 1099 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 354^\circ$;

▲ SC 19d11^h09^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 July 19



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 July 19

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	05 ^h	11 ^h	19.4	2d	
EPS	>10	05 ^h	11 ^h	3.6	2d	
EPS	>30	05 ^h	10 ^h	0.4	2d	
EPS	>50	-	10 ^h	0.25	2d	
EPS	>60	-	10 ^h	0.2	2d	
EPS	>100	-	10 ^h	0.1	2d	
METEOR						
CBM	>5	02 ^h	09 ^h	4.2	2d	
CBM	>15	02 ^h	09 ^h	1.8	1.5d	
CBM	>25	01 ^h	09 ^h	0.7	1.5d	
CBM	>40	01 ^h	09 ^h	0.35	1.4d	
BP	>90	01 ^h	09 ^h	0.06	1.0d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	14 ^h	5140	2d	
MEPED	>0.8	-	14 ^h	2520	2d	
MEPED	>2.5	-	12 ^h	510	2d	
MEPED	>6.9	-	12 ^h	50	2d	
MEPED	>16	-	10 ^h	4	2d	
MEPED	>36	-	10 ^h	1	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	15 ^h	1910	2d	
MKL	>14	-	15 ^h	1.6	2d	
MKL	>26	-	15 ^h	1.2	2d	
MKL	>50	-	15 ^h	0.54	2d	
ACE						
SIS	>10	05 ^h	09 ^h	8	2d	
SIS	>30	05 ^h	07 ^h	1.5	2d	
SOHO						
EPHIN (INT)	>50	18d21 ^h	07 ^h	0.14	1.5d	

Differential fluxes of protons for the event of 2002 July 19


S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	-	12 ^h	60.2	2d	
LION	2-6	-	12 ^h	7.7	2d	
EPHIN	4-8	04 ^h	11 ^h	18	2d	
EPHIN	8-25	02 ^h	10 ^h	0.56	2d	
EPHIN	25-41	03 ^h	07 ^h	0.028	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 July 19

2002	July 18	☉	AR10030	To event 426			
H α	6563 Å	0741	0743	0801	N19W33	2B	FU
1 – 12	keV	0724	0744	0749		X1.8	5.6E-2
CME	WL	0806	1099 km/s	-30.2km/s ²	360°	354°	0806

2002	July 18	Ø	AR10039	To event 426			
H α	6563 Å	No Flare Patrol					
1 – 12	keV	1952	1957	2006		C2.1	1.5E-3
12 - 25	keV	195256	195438	200308		137568	RHESSI

9.5	GHz	1953.7	1954.7	1955.6		1.18	
610	MHz	1949.0	2014.0	2045.0	P0.6	2.93	
410	MHz	1950.0	2000.0	2045.0		2.48	
245	MHz	2033.0	2033.0	2034.0		2.63	
DS III	25-300	2037		2039	B	3	
DS III	25-130	2047		2047	B	1	
DS V	25-180	2033		2035		2	
CME	WL	1931	2191 km/s	-129.3km/s ²	360°	104°	



Particle event: To(Ep>10 MeV) – 22d01^h

Tmax(Ep>10 MeV) – 22d11^h, Jmax(Ep>10 MeV) – 18.5 /cm².s.sr

Duration of the event – 10 days

Quasimaximal energy of protons in the event – E_{qm} = 85 MeV

Sources: ☉ solar flare 20d21^h04^m, X3.3/..., s13e90*, AR10039


☽ solar flare 23d00^h18^m, X4.8/2B, S12E70, AR10039

Main X-ray burst 1-8 Å: onset – 20d21^h04^m, max – 20d21^h30^m, Φ = 0.72 J/m²



CME: 20d22^h06^m, V = 1941 km/s, Δφ = 360°, = 91°

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 July 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 July 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm2.s.sr)-1	Dura- tion	Comments
GOES-10						
EPS	>5	01 ^h	11 ^h	42.4	10d	
EPS	>10	01 ^h	11 ^h	18.5	10d	
EPS	>30	01 ^h	12 ^h	1.9	9d	
EPS	>50	01 ^h	12 ^h	0.5	3d	
EPS	>60	01 ^h	12 ^h	0.33	3d	
EPS	>100	-	-	-	-	
METEOR						
CBM	>5	-	14 ^h	23.3	11d	
CBM	>15	-	12 ^h	5.6	9d	
CBM	>25	-	12 ^h	1.3	7d	
CBM	>40	-	12 ^h	0.37	3d	
BP	>90	-	12 ^h	0.02	-	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	10 ^h	150	10d	
MEPED	>0.8	-	10 ^h	93	9d	
MEPED	>2.5	-	10 ^h	72	8d	
MEPED	>6.9	-	10 ^h	20	7d	
MEPED	>16	-	10 ^h	8	3d	
MEPED	>36	-	10 ^h	1	3d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	21 ^h	190	8d	
MKL	>14	-	21 ^h	6.8	7d	
MKL	>26	-	21 ^h	4.5	3d	
MKL	>50	-	21 ^h	1.5	3d	
ACE						
SIS	>10	04 ^h	10 ^h	17	4d	
SIS	>30	04 ^h	09 ^h	1.7	4d	
SOHO						
EPHIN (INT)	>50	05 ^h	11 ^h	0.2	2d	

Differential fluxes of protons for the event of 2002 July 22

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	06 ^h	11 ^h	2.6	11d	
LION	2-6	06 ^h	11 ^h	1	11d	
EPHIN	4-8	05 ^h	10 ^h	7	10d	
EPHIN	8-25	05 ^h	11 ^h	1.1	10d	
EPHIN	25-41	05 ^h	10 ^h	0.07	10d	


References:

Yurchyshyn V., H. Wang, V. Abramenko et al., 2004.
 Gan W.Q, 2004.
 Kichigin G.N., L.I. Miroshnichenko, V.I. Sidorov et al., 2010.
 Struminsky A.B., 2011.
 Miroshnichenko L.I. and W.Q. Gan, 2012.
 Kichigin G.N., L.I. Miroshnichenko, V.I. Sidorov et al., 2012.
 Somov B.V., 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 July 22

2002 July 20		• AR10039			To event 427		
H α	6563 Å	No Flare Patrol			s13e87*		
1 – 12	keV	2104	2130	2154		X3.3	7.2E-1
100-300	keV	213000	213810	215300		50909556	RHESSI
500-1300							SONG F
15.4	GHz	2105.0	2128.0	2203.0	U0.4 / 15	4.63	
8.8	GHz	2105.0	2130.0	2203.0		4.30	
5	GHz	2104.0	2130.0	2203.0		3.81	
2.7	GHz	2104.0	2129.0	2203.0		3.41	
1.4	GHz	2105.0	2127.0	2203.0		3.18	
610	MHz	2106.0	2109.0	2203.0		2.63	
410	MHz	2106.0	2130.0	2155.0		2.52	
245	MHz	2106.0	2106.0	2154.0		4.08	

DS II	30-240	2107		2128		3	
DS II	110-400	2110		2120	S,H	3	
DS IV	57-1300	2107		2126		1	
DS IV	60-1700	<2128		2152		1	
DS I	140-180	2140		2213	S	1	
DS III	57-330	2107		2111	G	3	
DS III	25-180	2107		2129	N	2	
DS III	57-1700	2127		2130	G	2	
15.4	GHz	2105.0	2128.0	2203.0	U0.4 / 15	4.63	
8.8	GHz	2105.0	2130.0	2203.0		4.30	
5	GHz	2104.0	2130.0	2203.0		3.81	
2.7	GHz	2104.0	2129.0	2203.0		3.41	
1.4	GHz	2105.0	2127.0	2203.0		3.18	
610	MHz	2106.0	2109.0	2203.0		2.63	
410	MHz	2106.0	2130.0	2155.0		2.52	
245	MHz	2106.0	2106.0	2154.0		4.08	
DS II	30-240	2107		2128		3	
DS II	110-400	2110		2120	S,H	3	
DS IV	57-1300	2107		2126		1	
DS IV	60-1700	<2128		2152		1	
DS I	140-180	2140		2213	S	1	
DS III	57-330	2107		2111	G	3	
DS III	25-180	2107		2129	N	2	
DS III	57-1700	2127		2130	G	2	
n°						Mauna Kea, Haleakala	
CME	WL	2206	1941 km/s	–	360°	091°	




2002 July 23

Ø

AR10039

To event 427

H α	6563 Å	0023	0029	0240	S12E70	2B	EF
1 – 12	keV	0018	0035	0047		X4.8	4.6E-1
300-800	keV	001816	003038	011608		188025680	RHESSI
6 – 12	keV	015216	015222	015428		95520	RHESSI
15.4	GHz	0023.0	0031.0	0145.0	U0.4 / 15	4.00	
8.8	GHz	0022.0	0038.0	0145.0		3.97	
5	GHz	0021.0	0038.0	0144.0		3.63	
2.7	GHz	0022.0	0029.0	0144.0		3.26	
1.4	GHz	0025.0	0028.0	0000.0		3.20	
610	MHz	0025.0	0029.0	0133.0		3.18	
410	MHz	0025.0	0029.0	0000.0		3.15	
245	MHz	0026.0	0029.0	0000.0		3.90	
200	MHz	0026.0	0029.0	0118.0		3.66	
DS II	25-260	0029		0053		3	
DS IV	57-1500	0028		0046		1	
DS IV	25-180	0050		0226		1	
DS I	75-160	0046		~0130	S,C	1	
DS III	57-1300	0027		0031	G	1	
DS III	25-180	0028		0031		2	
DS III	57-300	0038		0040	G	2	
CME	WL	0042	2285 km/s	–	360°	087°	



Particle event: To($E_p > 10$ MeV) – 14d06^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 14d09^h$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 6.7 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 14d16^h$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 6.9 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm1} = 85 \text{ MeV}$


– $E_{qm2} = 50 \text{ MeV}$

Sources: • solar flare 14d01^h47^m, M2.3/1N, N10W54, AR10061



Main X-ray burst 1-8 Å: onset – 14d01^h47^m, max – 14d02^h12^m, $\Phi = 0.06 \text{ J/m}^2$

CME: 14d02^h30^m, $V = 1309 \text{ km/s}$, $\Delta\phi = 133^\circ$, $dA = 282^\circ$;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 August 14



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 August 14

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm2.s.sr)-1	Dura- tion	Comments
GOES 10						
EPS	>5	02 ^h	09 ^h /16 ^h	26.5/44	2d	
EPS	>10	02 ^h	09 ^h /16 ^h	6.7/6.9	2d	
EPS	>30	02 ^h	09 ^h /16 ^h	0.3/0.12	1d	
EPS	>50	02 ^h	09 ^h /16 ^h	0.1/0.015	1d	
EPS	>60	-	09 ^h / -	0.07/ -	1d	
EPS	>100	-	-	-	-	
POES 16						
MEPED	>0.24	-	10 ^h /16 ^h	- /1720	2d	
MEPED	>0.8	-	10 ^h /16 ^h	310 /1220	2d	
MEPED	>2.5	-	10 ^h /16 ^h	170/630	2d	
MEPED	>6.9	-	10 ^h /16 ^h	40/70	2d	
MEPED	>16	-	11 ^h /16 ^h	5/2	1d	
MEPED	>36	-	10 ^h /16 ^h	0.6 /0.4	1d	
MEPED	>70	-	10 ^h /16 ^h	0.2/ -	1d	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	-	08 ^h / -	250/ -	2d	
MKL	>14	-	08 ^h / -	2.7/ -	1d	
MKL	>26	-	08 ^h / -	1.85/ -	1d	
MKL	>50	-	08 ^h / -	0.46/ -	1d	

ACE						
SIS	>10	02 ^h	09 ^h /15 ^h	9.5/23.8	2d	
SIS	>30	02 ^h	09 ^h /15 ^h	0.9/0.1	2d	
SOHO						
EPHIN (INT)	>50	-	08 ^h / -	0.02/ -	-	



Differential fluxes of protons for the event of 2002 August 14

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	03 ^h	09 ^h /16 ^h	21.6/105	2.5d	
LION	2-6	03 ^h	09 ^h /16 ^h	4.6/28.6	2d	
EPHIN	4-8	03 ^h	09 ^h /16 ^h	8.6/77.9	2.5d	
EPHIN	8-25	03 ^h	09 ^h /16 ^h	0.4/3.8	2.5d	
EPHIN	25-41	02 ^h	07 ^h	0.027/0.009	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 August 14

2002 August 14		● AR10039		To event 428			
H α	6563 Å	0147	0201	0343	N09W54	1N	EF
1 – 12	keV	0147	0212	0246		M2.3	6.0E-2
12-25	keV	021400	021422	023428		4238062	RHESSI
6-12	keV	023428	023606	031500		1477953	RHESSI
8.8	GHz	0201.0	0209.0	0210.0		1.66	
5	GHz	0149.0	0201.0	0308.0		2.04	
2.8	GHz	0143.0	0202.2	0259.0	U0.5 P2.8	2.26	
1.4	GHz	0144.0	0203.0	0000.0		2.08	
500	MHz	0145.0	0203.0	0245.0		1.81	
245	MHz	0152.0	0200.0	0223.0		2.88	
DS II	25-157	0157		0208		3	
DS III	25-140	0155		0158	G	1	
DS III	57-160	0157		0159	G	2	
DS V	25-144	0159		0205		3	

15.4	GHz	0232.0	0235.0	0246.0		2.00	
8.8	GHz	0232.0	0235.0	0246.0	[U0.6 P8.8]	2.23	
5	GHz	0232.0	0235.0	0246.0		2.08	
2.7	GHz	0232.0	0235.0	0246.0		1.85	
1.4	GHz	0239.0	0239.0	~0239.0		2.08	
610	MHz	0235.0	0235.0	~0235.0		1.52	
245	MHz	0234.0	0239.0	0246.0		2.08	
DS III	25-180	0300		0304		1	
DS III	25-400	0304		0305	B	2	
CME	WL	0230	1309 km/s	-28.5	133°	282°	



Particle event: To(Ep>10 MeV) – 17d00^h

Tmax(Ep>10 MeV) – 17d10^h, Jmax (Ep>10 MeV) – 1.7 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – E_{qm} = 70 MeV


Sources: ● solar flare 16d11^h11^m, 2N/M5.2, S14E20, AR10069

Main X-ray burst 1-8 Å: onset – 16d11^h32^m, max – 16d12^h32^m, Φ = 0.16 J/m²



CME: 16d12^h30^m; V = 1585 km/s, Δφ = 360°, dA = 121°.

▲ SC 18d18^h46^m;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 August 17



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 August 17

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm2.s.sr)-1	Dura- tion	Comments
GOES 10						
EPS	>5	00 ^h	10 ^h	9	2d	
EPS	>10	00 ^h	10 ^h	1.7	2d	
EPS	>30	00 ^h	09 ^h	0.12	1d	
EPS	>50	-	09 ^h	0.06	-	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
ACE						
SIS	>10	16d18 ^h	09 ^h	2.6	2d	
SIS	>30	-	-	-	-	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 2002 August 17



S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm².s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	16d19 ^h	08 ^h	4.3	2d	
LION	2-6	16d19 ^h	08 ^h	1	2d	
EPHIN	4-8	16d18 ^h	08 ^h	3.7	2d	
EPHIN	8-25	16d18 ^h	08 ^h	0.23	2d	
EPHIN	25-41	16d18 ^h	08 ^h	0.0025	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.
Tylka A.J., O.E. Malandraki, G. Dorrian et al., 2013.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2002 August 17**

2002	August 16	•	AR10069			To event 429	
H α	6563 Å	1111	1213	1409	S14E20	2N	FH
1 – 12	keV	1132	1232	1307		M5.2	1.6E-1
50-100	keV	120604	122118	123156		20782080	RHESSI
25 – 50	keV	133824	134654	140616		1091448	RHESSI
15.4	GHz	1133.0	1135.0	1143.0		2.28	
8.8	GHz	1132.0	1135.0	1143.0	U0.4 P8.8	2.38	
5	GHz	1133.0	1135.0	0000.0		2.11	
2.7	GHz	1133.0	1136.0	1143.0		1.95	
1.4	GHz	1132.0	1135.0	1143.0		2.00	
610	MHz	1132.0	1134.0	1143.0		1.96	
410	MHz	1133.0	1135.0	1143.0		1.92	
204	MHz	1134.2	1135.3	1138.0		3.23	
DS II	45-65	1144		1147		2	
DS I	45-270	1156		~1335	S,C	2	
DS III	25-95	1127		1128	B	2	
DS III	55-270	1133		1141	GG,C	2	
DS III	30-75	1143		1205	N	2	
DS III	25-270	1145		1150	GG,FS	2	
DS III	25-180	1149		1205	N	2	
DS DCIM	130-4000	1132		1145	S,P	3	
15.4	GHz	1133.0	1213.0	1441.0		3.00	
8.8	GHz	1132.0	1213.0	1428.0		3.28	
5	GHz	1133.0	1213.0	1448.0	P5	3.36	
2.7	GHz	1134.0	1214.0	1426.0		3.20	
1.4	GHz	1132.0	1217.0	1412.0		3.11	
DS II	100-600	1205		1317		3	
DS II	25-180	1206		1224		3	
DS IV	25-180	1205		1726		3	
DS III	25-270	1203		1215	GG,FS	2	
DS III	25-270	1215		~1335	S	2	
DS CONT	45-270	1204		~1335	GG,FS	2	
DS DCIM	100-4000	1208		1317	P,C,S	3	
CME		1230	1585km/s	-67.1	360°	121°	



Particle event: To($E_p > 10$ MeV) – 18d22^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 19\text{d}03^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 2.3 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 19\text{d}12^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 1.8 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm1} = 75 \text{ MeV}$


– $E_{qm2} = 70 \text{ MeV}$

Sources: • solar flare 18d21^h11^m, 1N/M2.2, S10W20, AR10069



Main X-ray burst 1–8 Å: onset – 18d21^h12^m, max – 18d21^h25^m, $\Phi = 0.22 \text{ J/m}^2$

CME: 18d21^h54^m, $V = 0682 \text{ km/s}$, $\Delta\phi = 140^\circ$, $dA = 202^\circ$.

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 August 18



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 August 18

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm2.s.sr)-1	Dura- tion	Comments
GOES 10						
EPS	>5	22 ^h	19d03 ^h /19d11 ^h	11.3/9.8	2d	
EPS	>10	22 ^h	19d03 ^h /19d12 ^h	2.3/1.8	2d	
EPS	>30	22 ^h	19d02 ^h /19d12 ^h	0.3/0.1	1.5d	
EPS	>50	22 ^h	19d02 ^h /19d12 ^h	0.12/0.03	1.5d	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
POES 16						
MEPED	>0.24	-	19d07 ^h / -	3120/ -	2d	
MEPED	>0.8	-	19d07 ^h / -	910/ -	2d	
MEPED	>2.5	-	19d05 ^h / -	84/ -	2d	
MEPED	>6.9	-	19d04 ^h / -	10/ -	2d	
MEPED	>16	-	19d02 ^h / -	1.5/ -	1.5d	
MEPED	>36	-	19d02 ^h / -	0.7/ -	1.5d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	-	19d00 ^h / -	445/ -	2d	
MKL	>14	-	19d00 ^h / -	1.6/ -	1.5d	
MKL	>26	-	19d00 ^h / -	1.14/ -	1.5d	
MKL	>50	-	19d00 ^h / -	0.46/ -	1.5d	

ACE						
SIS	>10	22 ^h	19d02 ^h /19d12 ^h	3.5/2.25	2d	
SIS	>30	22 ^h	19d02 ^h /19d11 ^h	0.4/0.1	1d	
SOHO						
EPHIN (INT)	>50	22 ^h	19d02 ^h /19d11 ^h	0.015/0.05	-	


Differential fluxes of protons for the event of 2002 August 18

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	22 ^h	19d01 ^h /19d11 ^h	22/25.2	2d	
LION	2-6	22 ^h	19d01 ^h /19d11 ^h	1.6/2	2d	
EPHIN	4-8	22 ^h	19d03 ^h /19d11 ^h	3.2/3.6	2d	
EPHIN	8-25	22 ^h	19d03 ^h /19d11 ^h	0.2/0.21	2d	
EPHIN	25-41	22 ^h	19d02 ^h /19d11 ^h	0.01/0.005	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 August 18

2002	August 18	•	AR10069	To event 430			
H α	6563 Å	2111	2121	2200	S10W20	1N	FH
1 – 12	keV	2112	2125	2137		M2.2	2.2E-2
15.4	GHz	2115.0	2121.0	2131.0		3.00	
8.8	GHz	2113.0	2121.0	2134.0	U0.6 P8.8	3.41	
5	GHz	2113.0	2121.0	2132.0		3.28	
2.7	GHz	2114.0	2121.0	2130.0		2.83	
1.4	GHz	2114.0	2115.0	2125.0		2.66	
610	MHz	2113.0	2116.0	2124.0		2.60	
410	MHz	2113.0	2115.0	2123.0		2.73	
245	MHz	2111.0	2116.0	2139.0		2.72	

DS II	28-180	2124		2155		1	
DS II	25-130	2124		2135		3	
DS IV	25-180	2136		0140		1	
DS IV	30-80	2142		2320		2	
DS III	25-2000	2110		2122	G	3	
DS III	25-80	2142		2142	B	1	
DS III	330-530	2144		2148	G	1	
DS III	25-500	2159		2203	G	2	
DS CONT	60-110	2135		>2400		3	
CME	WL	2154	0682km/s	1.9 km/s ²	140°	202°	



Particle event: To($E_p > 10$ MeV) – 20d09^h

Tmax($E_p > 10$ MeV) – 20d10^h, Jmax ($E_p > 10$ MeV) – 2.5 /cm².s.sr *)

Duration of the event – 1 day *)

Quasimaximal energy of protons in the event – $E_{qm} = 80$ MeV


*) Data from ACE (SIS)

Sources: • solar flare 20d01^h33^m, M5.0/1B, S10W35, AR10069



Main X-ray burst 1-8 Å: onset – 20d01^h33^m, max – 20d01^h40^m, $\Phi = 0.12$ J/m²

CME: 20d01^h54^m; $V = 0961$ km/s, $\Delta\phi = 360^\circ$, $dA = 121^\circ$.

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 August 20



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 August 20

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES 10						
EPS	>5	09 ^h	10 ^h	0.7	2d	
EPS	>10	09 ^h	10 ^h	0.4	2d	
EPS	>30	08 ^h	11 ^h	0.07	2d	
EPS	>50	08 ^h	11 ^h	0.05	2d	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
METEOR						
CBM	>5	09 ^h	10 ^h	17.3	1.2d	
CBM	>15	09 ^h	10 ^h	3.2	1.2d	
CBM	>25	09 ^h	10 ^h	0.5	0.7d	
CBM	>40	09 ^h	10 ^h	0.2	0.7d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
ACE						
SIS	>10	09 ^h	10 ^h	2.5	1d	
SIS	>30	08 ^h	10 ^h	0.5	1d	
SOHO						
EPHIN (INT)	>50	08 ^h	10 ^h	0.3	1d	



Differential fluxes of protons for the event of 2002 August 20

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	08 ^h	14 ^h	4.7	2d	
LION	2-6	08 ^h	13 ^h	0.75	2d	
EPHIN	4-8	08 ^h	11 ^h	2	2d	
EPHIN	8-25	08 ^h	11 ^h	0.2	2d	
EPHIN	25-41	08 ^h	10 ^h	0.03	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 August 20

2002 August 20		• AR10069		To event 431			
H α	6563 Å	0135	0140	0202	S10W35	1B	EF
1 – 12	keV	0133	0140	0143		M5.0	1.2E-2
25 – 50	keV	013044	014022	014836		6439679	RHESSI
12 – 25	keV	014836	015022	015816		697067	RHESSI
15.4	GHz	0149.0	0149.0	0151.0	U5 / 15	2.65	
8.8	GHz	0139.0	0140.0	0141.0		2.00	
5	GHz	0139.0	0139.0	0140.0		1.80	
2.8	GHz	0137.0	0139.5	0152.0		2.01	
1.4	GHz	0138.0	0139.0	0139.0		2.04	
610	MHz	0138.0	0139.0	0000.0		3.28	
410	MHz	0134.0	0139.0	0142.0		3.90	
245	MHz	0134.0	0139.0	0143.0		4.58	
200	MHz	0137.0	0139.0	0147.0		3.43	
DS III	25-1600	0134		0149	G	3	
8.8	GHz	0208.0	0209.0	0210.0		2.15	
5	GHz	0207.0	0209.0	0214.0	[P5]	2.28	
2.8	GHz	0204.0	0209.1	0230.0		2.15	
1.4	GHz	0208.0	0209.0	0210.0		1.77	
610	MHz	0209.0	0209.0	0210.0		1.84	
410	MHz	0209.0	0209.0	~0209.0		1.73	
245	MHz	0210.0	0210.0	~0210.0		1.71	
DS I	100-180	0247		0350	S	1	
DS III	57-160	0208		0209	G	3	
DS III	25-270	0208		0212	G	2	
DS III	57-1000	0210		0210	G	1	
DS III	57-130	0211		0212	G	2	

15.4	GHz	0254.0	0255.0	0301.0		2.26	
8.8	GHz	0255.0	0257.0	0258.0		1.86	
200	MHz	0258.0	0258.0	0312.0		2.18	
DS I	100-180	0247		0350	S	1	
CME	WL	0154	0961 km/s	1.7 km/s ²	157°	210°	



Particle event: To($E_p > 10$ MeV) – 22d03^h

Tmax($E_p > 10$ MeV) – 22d05^h, Jmax ($E_p > 10$ MeV) – 16 /cm².s.sr

Duration of the event – 2 days


Quasimaximal energy of protons in the event – $E_{qm} = 450$ MeV

Sources: • solar flare 22d01^h47^m, M5.4/2B, S07W62, AR10069



Main X-ray burst 1-8 Å onset – 22d01^h47^m, max – 22d01^h57^m, $\Phi = 0.033$ J/m²

CME: 22d02^h06^m; V = 998 km/s, $\Delta\varphi = 360^\circ$, dA = 231°.

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 August 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 August 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm2.s.sr)-1	Dura- tion	Comments
GOES-10						
EPS	>5	03 ^h	05 ^h	18.3	2d	
EPS	>10	03 ^h	05 ^h	16	2d	
EPS	>30	03 ^h	05 ^h	7.4	2d	
EPS	>50	03 ^h	05 ^h	3.9	2d	
EPS	>60	03 ^h	05 ^h	2.9	2d	
EPS	>100	03 ^h	05 ^h	1.2	2d	
METEOR						
CBM	>5	03 ^h	06 ^h	15	2d	
CBM	>15	03 ^h	05 ^h	9.5	2d	
CBM	>25	03 ^h	05 ^h	6.5	2d	
CBM	>40	03 ^h	05 ^h	4.4	2d	
BP	>90	03 ^h	05 ^h	0.9	1.5d	
ChD	>600	03 ^h	05 ^h	0.07	1d	
POES-16						
MEPED	>0.24	-	05 ^h	120	2d	
MEPED	>0.8	-	05 ^h	110	2d	
MEPED	>2.5	-	05 ^h	100	2d	
MEPED	>6.9	-	05 ^h	70	2d	
MEPED	>16	-	05 ^h	23	2d	
MEPED	>36	-	05 ^h	10	2d	
MEPED	>70	-	05 ^h	2.6	2d	
MEPED	>140	-	05 ^h	1	2d	

CORONAS F						
MKL	>1.	-	05 ^h	140	2d	
MKL	>14	-	05 ^h	15	2d	
MKL	>26	-	05 ^h	10	2d	
MKL	>50	-	05 ^h	5.5	2d	
ACE						
SIS	>10	02 ^h	04 ^h	22.6	2d	
SIS	>30	02 ^h	04 ^h	11	2d	
SOHO						
EPHIN (INT)	>50	02 ^h	04 ^h	1.7	1d	

Differential fluxes of protons for the event of 2002 August 22

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	02 ^h	06 ^h	0.5	2d	
LION	2-6	02 ^h	06 ^h	0.17	2d	
EPHIN	4-8	02 ^h	05 ^h	1.23	2d	
EPHIN	8-25	02 ^h	04 ^h	0.75	2d	
EPHIN	25-41	02 ^h	04 ^h	0.19	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 August 22

2002	August 22	•	AR10069	To event 432			
H α	6563 Å	0151	0153	0225	S07W62	2B	EF
1 – 12	keV	0147	0157	0205		M5.4	3.3E-2
50-100	keV	014508	015422	022212		9830232	RHESSI

15.4	GHz	0153.0	0156.0	0208.0		2.18	
8.8	GHz	0151.0	0156.0	0206.0		2.36	
5	GHz	0150.0	0151.0	0200.0	P5	2.71	
2.8	GHz	0136.0	0152.3	0240.0		2.46	
1.4	GHz	0150.0	0151.0	0153.0		2.46	
610	MHz	0150.0	0153.0	0154.0		2.48	
410	MHz	0139.0	0140.0	0140.0		2.18	
200	MHz	0153.0	0154.0	0155.0		1.78	
DS II	25-200	0156		0205		2	
DS IV	25-180	0209		0218		1	
DS IV	57-280	~0210		>0358	FS	1	
DS III	57-1300	0151		0155	GG	3	
DS III	25-280	0151		0156	G	3	
DS III	57-130	0202		0203	G	3	
DS III	30-190	0206		0214	G	2	
CME	WL	0206	0998 km/s	-32.8km/s ²	360°	153°	



Particle event: To(Ep> 10MeV) – 24d01^h

Tmax(Ep>10 MeV) – 24d03^h, Jmax (Ep>10 MeV) – 92 /cm².s.sr

Duration of the event – 4 days

Quasimaximal energy of protons in the event – E_{qm} = 775 MeV


Sources: ● solar flare 24d00^h49^m, X3.1/1F, S02W81, AR10069

Main X-ray burst 1–8 Å: onset – 24d00^h49^m, max – 24d01^h12^m, Φ = 0.46 J/m²;



CME: 24d01^h27^m; V = 1913 km/s, Δφ = 360°, dA = 270°;

▲ SC 26d11^h31^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 August 24



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 August 24

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm2.s.sr)-1	Dura- tion	Comments
GOES-10						
EPS	>5	01 ^h	03 ^h	96	4d	
EPS	>10	01 ^h	03 ^h	92	4d	
EPS	>30	01 ^h	03 ^h	87	4d	
EPS	>50	01 ^h	02 ^h	56	3.5d	
EPS	>60	01 ^h	02 ^h	47	3.5d	
EPS	>100	01 ^h	02 ^h	26	3.5d	
METEOR						
CBM	>5	02 ^h	03 ^h	187	4d	
CBM	>15	02 ^h	03 ^h	134	4d	
CBM	>25	02 ^h	03 ^h	100	3d	
CBM	>40	02 ^h	03 ^h	73	3d	
BP	>90	02 ^h	03 ^h	7.1	2.5d	
ChD	>600	02 ^h	03 ^h	1.43	2d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	04 ^h	730	4d	
MEPED	>16	-	04 ^h	190.	4d	
MEPED	>36	-	03 ^h	96	3.5d	
MEPED	>70	-	02 ^h	25	3.5d	
MEPED	>140	-	02 ^h	10	3.5d	

CORONAS F						
MKL	>1.	-	05 ^h	560	4d	
MKL	>14	-	05 ^h	128	3.5d	
MKL	>26	-	05 ^h	85	3.5d	
MKL	>50	-	05 ^h	32.5	3.5d	
ACE						
SIS	>10	01 ^h	03 ^h	146.	4d	
SIS	>30	01 ^h	03 ^h	100	4d	
SOHO						
EPHIN (INT)	>50	01 ^h	02 ^h	48	2.5d	
BALLOONS						
Mi	>183	-	07 ^h 06 ^m -08 ^h 06 ^m	0.76	-	
Mi	>359	-	07 ^h 06 ^m -08 ^h 06 ^m	0.27	-	
Mi	>647	-	07 ^h 06 ^m -08 ^h 06 ^m	0.15	-	
NM						
Network	>500	01 ^h	02 ^h	0.66	-	

Differential fluxes of protons for the event of 2002 August 24

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	01 ^h	04 ^h	0.65	4d	
LION	2-6	01 ^h	04 ^h	0.67	4d	
EPHIN	4-8	01 ^h	04 ^h	2.5	4d	
EPHIN	8-25	01 ^h	03 ^h	1.8	4d	
EPHIN	25-41	01 ^h	02 ^h	1.3	4d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Struminsky A.B., 2003.

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.

Miroshnichenko L.I. and J. Perez-Peraza, 2008.


Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 August 24

2002 August 24 • AR10069 To event 433

H α	6563 Å	0055	0103	0123	S02W81	1F	F
1 – 12	keV	0049	0112	0131		X3.1	4.6E-1
50–100	keV	004348	005730	005828		4036893	RHESSI
25–50	keV	012824	012954	023428		51893096	RHESSI
4–7	MeV						SONG F

15.4	GHz	0052.0	0100.0	0221.0	U1.4/15	4.26	
8.8	GHz	0051.0	0104.0	0219.0		4.11	
5	GHz	0050.0	0109.0	0219.0		3.77	
2.7	GHz	0050.0	0102.0	0207.0		3.08	
1.4	GHz	0054.0	0056.0	0146.0		2.62	
500	MHz	0053.0	0105.0	0119.0		2.88	
245	MHz	0101.0	0102.0	0128.0		2.45	
DS II	25-180	0101		0114		2	
DS IV	25-180	0115		0136		1	
DS III	60-1600	0101		0103	G	1	
DS III	57-180	0101		0104	G	3	
DS III	57-520	0104		0107	G	2	
DS CONT	57-200	~0108		0135		1	
CME	WL	0127	1913 km/s	32.8km/s ²	360°	270°	



Particle event: To(Ep>10 MeV) – 06d06^h

Tmax₁(Ep>10 MeV) – 06d14^h, Jmax₁(Ep>10 MeV) – 3 /cm².s.sr

Tmax₂(Ep>10 MeV) – 07d17^h, Jmax₂(Ep>10 MeV) – 67 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 80 MeV

– Eqm₂ = 175 MeV


Sources: ☉ solar flare 05d16^h18^m, C5.2/SF, N12E28, AR10102

Main X-ray burst 1-8 Å: onset – 05d16^h18^m, max – 05d17^h06^m, Φ = 0.016 J/m²



CME: 05d16^h54^m; V = 1748 km/s, Δφ = 360°, dA = 114°

▲ SC 07d16^h38^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 September 06



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2001 September 06

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm2.s.sr) ⁻¹	Duration	Comments
GOES 10						
EPS	>5	06 ^h	14 ^h /07d17 ^h	6.6/360	3d	
EPS	>10	06 ^h	14 ^h /07d17 ^h	3/67	3d	
EPS	>30	07 ^h	13 ^h /07d16 ^h	0.14/1.9	3d	
EPS	>50	07 ^h	13 ^h /07d16 ^h	0.05/0.33	2d	
EPS	>60	07 ^h	13 ^h /07d16 ^h	0.04/0.2	2d	
EPS	>100	07 ^h	- /07d16 ^h	- /0.04	1d	
METEOR						
CBM	>5	-	- /07d17 ^h	- /16	2d	
CBM	>15	-	- /07d17 ^h	- /4.6	2d	
CBM	>25	-	- /07d17 ^h	- /0.83	1d	
CBM	>40	-	- /07d17 ^h	- /0.17	1d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES 16						
MEPED	>0.24	-	16 ^h /07d18 ^h	-/59300	3d	
MEPED	>0.8	-	16 ^h /07d18 ^h	-/29600	3d	
MEPED	>2.5	-	16 ^h /07d18 ^h	-/7870	3d	
MEPED	>6.9	-	16 ^h /07d18 ^h	-/790	2d	
MEPED	>16	-	16 ^h /07d18 ^h	2/30	2d	
MEPED	>36	-	16 ^h /07d18 ^h	-/1.5	1d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	18 ^h /07d18 ^h	33/8150	3d	
MKL	>14	-	18 ^h /07d18 ^h	1.5/6.4	2d	
MKL	>26	-	18 ^h /07d18 ^h	1.14/3.9	2d	
MKL	>50	-	18 ^h /07d18 ^h	0.5/1.14	1d	
ACE						
SIS	>10	03 ^h	13 ^h /07d16 ^h	5.5/72.5	3d	
SIS	>30	05 ^h	13 ^h /07d15 ^h	0.4/3.8	2d	
SOHO						
EPHIN (INT)	>50	-	18 ^h /07d15 ^h	0.03/0.25	0.3d	


Differential fluxes of protons for the event of 2001 September 06

S/c, instru-ments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	05d23 ^h	17 ^h /07d16 ^h	1.5/285	4d	
LION	2-6	05d23 ^h	17 ^h /07d16 ^h	0.58/46	4d	
EPHIN	4-8	05d22 ^h	16 ^h /07d15 ^h	2.7/178	4d	
EPHIN	8-25	05d22 ^h	16 ^h /07d15 ^h	0.44/12.2	4d	
EPHIN	25-41	05d22 ^h	15 ^h /07d15 ^h	0.016/0.21	4d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 September 06

2002	September 05	☉	AR10102	To event 434			
H α	6563 Å	1630	1645	1745	N12E28	SF	FU
DSF		~1631		~1734	N08E31	06°	
1 – 12	keV	1618	1706	1735		C5.2	1.6E-2
6-12	keV	170620	170702	173008		121416	RHESSI

2.7	GHz	1632.0	1632.0	~1632.0		1.59	
1.4	GHz	1630.0	1634.0	1635.0		1.74	
610	MHz	1629.0	1635.0	1643.0	P0.6	2.20	
410	MHz	1631.0	1634.0	1636.0		2.15	
DS II	40-250	1633		1647		2	
DS I	200-350	~1710		~1733	S,N	1	
DS III	40-65	1628		1628	B	2	
DS CONT	25-180	1635		0429		1	
DS DCIM	320-700	1632		1637		2	
CME	WL	1654	1748 km/s	43.0 km/s ²	360°	114°	



Particle event: To(Ep>10 MeV) – 09d17^h

Tmax₁(Ep>10 MeV) – 10d02^h, Jmax₁(Ep>10 MeV) – 150 /cm².s.sr

Tmax₂(Ep>10 MeV) – 10d13^h, Jmax₂(Ep>10 MeV) – 40 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 160 MeV

– Eqm₂ = 145 MeV

Sources: ● solar flare 09d13^h08^m, M4.6/2B, S04W29, AR10180


○ solar flare 10d03^h04^m, M2.4/2N, S12W37, AR10180

Main X-ray burst 1-8 Å: onset – 09d13^h08^m, max – 09d13^h23^m, Φ = 0.048 J/m²



CME: 09d13^h32^m; V = 1838 km/s, Δφ = 360°, dA = 233°.

▲ SC 09d17^h51^m; ▲ 09d18^h49^m; ▲ 11d12^h31^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2002 November 09



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2002 November 09

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	17 ^h	10d02 ^h /10d13 ^h	324/135	2d	
EPS	>10	17 ^h	10d02 ^h /10d13 ^h	150/40	2d	
EPS	>30	17 ^h	10d01 ^h /10d13 ^h	8.7/1.6	2d	
EPS	>50	17 ^h	10d02 ^h /10d13 ^h	1.2/0.32	1d	
EPS	>60	17 ^h	10d02 ^h /10d13 ^h	0.6/0.26	1d	
EPS	>100	17 ^h	10d02 ^h /10d13 ^h	0.07 /0.08	1d	
METEOR						
CBM	>5	13 ^h	10d05 ^h / -	104/ -	3d	
CBM	>15	13 ^h	10d05 ^h / -	34/ -	2d	
CBM	>25	13 ^h	10d05 ^h / -	7.3/ -	2d	
CBM	>40	13 ^h	10d05 ^h / -	1.2/ -	2d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	10d06 ^h /10d22 ^h	- /2120	3d	
MEPED	>0.8	-	10d06 ^h /10d22 ^h	1330/1530	2d	
MEPED	>2.5	-	10d05 ^h /10d22 ^h	1110/810	2d	
MEPED	>6.9	-	10d02 ^h /10d22 ^h	690/210	2d	
MEPED	>16	-	10d02 ^h /10d22 ^h	80/7	1d	
MEPED	>36	-	10d02 ^h /10d22 ^h	4/1	1d	
MEPED	>70	-	-	- /0.3	1d	
CORONAS F						
MKL	>1.	-	10d05 ^h /10d22 ^h	1530/2150	2d	
MKL	>14	-	10d02 ^h /10d22 ^h	40/30	1d	
MKL	>26	-	10d02 ^h /10d22 ^h	25//2	1d	
MKL	>50	-	10d02 ^h /10d22 ^h	7.4/0.64	1d	

ACE						
SIS	>10	15 ^h	10d02 ^h /10d13 ^h	270/53	2d	
SIS	>30	15 ^h	10d02 ^h /10d13 ^h	13.5/1.7	1d	
SOHO						
EPHIN (INT)	>50	15 ^h	10d03 ^h /10d15 ^h	0.38/0.16	2d	

Differential fluxes of protons for the event of 2002 November 09


S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	17 ^h	10d03 ^h /10d15 ^h	29.5/98.4	3d	
LION	2-6	17 ^h	10d03 ^h /10d15 ^h	16.7/30.5	3d	
EPHIN	4-8	15 ^h	10d03 ^h /10d14 ^h	127/179	3d	
EPHIN	8-25	14 ^h	10d03 ^h /10d14 ^h	22/13	2d	
EPHIN	25-41	14 ^h	10d03 ^h /10d14 ^h	0.62/0.13	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Struminsky A., 2003.
 Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
 Struminsky A.B., 2007.


Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2002 November 09

2002 November 09 • AR10180					To event 435		
Hα	6563 Å	1309	1322	1428	S12W29	M4.6/2B	FU
1 – 12	keV	1308	1323	1336		M4.6	4.8E-2
100–300	keV	130712	132150	132840		9681024	RHESSI
25 – 50	keV	1400	140534	140932		172848	RHESSI
12 – 25	keV	140932	141134	14262		140472	RHESSI
15.4	GHz	1312.0	1317.0	0000.0		2.28	
9.5	GHz	1310.0	1321.2	1328.1	U0.6 [P2.7 U5] P9.5	2.78	
5	GHz	1311.0	1312.0	0000.0		2.30	
2.7	GHz	1310.0	1316.0	0000.0		2.70	
1.4	GHz	1311.0	1316.0	0000.0		2.52	
610	MHz	1311.0	1311.0	~1311.0		1.46	
410	MHz	1312.0	1314.0	0000.0		2.11	
245	MHz	1309.0	1312.0	0000.0		2.59	
DS II	25-85	1317		1337		2	
DS IV	25-180	1307		1353		2	
DS IV	100-4000	1308		1324	P,F	3	
DS III	25-180	1402		1405		1	
DS DCIM	800-2000	1307		1330	GG,F,S,P	2	
DS DCIM	2000-4500	1310		1328	GG	2	
CME	WL	1332	1838km/s	35.4 km/s ²	360°	233°	



2002 November 10 Ø AR10180 To event 435

H α	6563 Å	0307	0314	0358	S12W37	2N	FU
1 – 12	keV	0304	0321	0335		M2.4	3.0E-2
100-300	keV	030520	031654	035508		7302648	HESSI
15.4	GHz	0311.0	0312.0	0313.0		2.20	
8.8	GHz	0309.0	0311.0	0315.0		2.66	
5	GHz	0308.0	0311.0	0317.0	U0.5 P5	2.85	
2.8	GHz	0301.0	0311.3	0333.0		2.75	
500	MHz	0307.0	0311.0	0321.0		1.65	
200	MHz	0307.0	0309.0	0317.0		2.00	
DS II		0314		0325	25-180	3	
DS II	FN	0317		0323	18-50	3	
DS II	FN	0320		0334	18-80	3	
DS II	SH	0320		0334	40-160	3	
DS III	GG	0307		0313	18-800	3	
DS III	G	0313		0319	23-480	1	
DS CONT		0325		0335	25-180	1	
CME		0330	1670km/s	35.4 km/s ²	360°	316°	



Events in 2003

		№			Page
1.	Event 2003.05.28 – (2003-148)	№ 436	.	.	526
2.	Event 2003.05.31 – (2003-151)	№ 437	.	.	532
3.	Event 2003.06.18 – (2003-169)	№ 438	.	.	538
4.	Event 2003.10.26 – (2003-299)	№ 439	.	.	542
5.	Event 2003.10.28 – (2003-301) – GLE-65	№ 440	.	.	547
6.	Event 2003.10.29 – (2003-302) – GLE-66	№ 441	.	.	552
7.	Event 2003.11.02 – (2003-306) – GLE-67	№ 442	.	.	555
8.	Event 2003.11.04 – (2003-308)	№ 443	.	.	563
9.	Event 2003.11.20 – (2003-324)	№ 444	.	.	569
10.	Event 2003.11.21 – (2003-325)	№ 445	.	.	572
11.	Event 2003.12.02 – (2003-336)	№ 446	.	.	579

Particle event: To($E_p > 10$ MeV) – 28d04^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 28\text{d}11^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 2 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 29\text{d}16^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 77 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{\text{qm}1} = 170 \text{ MeV}$

– $E_{\text{qm}2} = 175 \text{ MeV}$

Sun source: ● solar flare 27d22^h56^m, X1.3/2B, S06W20, AR10365*

solar flare 28d00^h 27^m, X3.6/2B, S06W20, AR10365*

Ø solar flare 29d00^h 20^m, X1.3/2B, S06W37, AR10365

Main X-ray burst 1-8 Å: onset – 27d22^h56^m, max – 27d23^h07^m, $\Phi = 0.071 \text{ J/m}^2$

– 28d00^h 17^m, max – 28d00^h27^m, $\Phi = 0.28 \text{ J/m}^2$


CME: 27d23^h50^m, $V = 0964 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 067^\circ$

CME: 28d00^h50^m, $V = 1366 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 292^\circ$



▲ SC: 29d12^h24^m

*One flare event with two X-ray bursts

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2003 May 28



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 May 28

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	04 ^h	28d11 ^h /29d16 ^h	8.7/405	3d	
EPS	>10	04 ^h	28d11 ^h /29d16 ^h	2/77	3d	
EPS	>30	03 ^h	28d11 ^h /29d16 ^h	0.5/2	3d	
EPS	>50	03 ^h	28d11 ^h /29d16 ^h	0.3/0.62	3d	
EPS	>60	03 ^h	28d11 ^h /29d16 ^h	0.25/0.52	2d	
EPS	>100	06 ^h	28d11 ^h /29d16 ^h	0.1/0.42	2d	
METEOR						
CBM	>5	03 ^h 36 ^m	- /29d16 ^h	3.1	1d	
CBM	>10	03 ^h 36 ^m	- /29d16 ^h	1.3	1d	
CBM	>25	03 ^h 36 ^m	- /29d16 ^h	0.53	1d	
CBM	>40	03 ^h 36 ^m	- /29d16 ^h	0.33	1d	
BP	>90	03 ^h 36 ^m	- /29d16 ^h	0.07	1d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	28d11 ^h /29d16 ^h	61/1.8·10 ⁴	3d	
MEPED	>0.8	-	28d11 ^h /29d16 ^h	28/7·10 ³	3d	
MEPED	>2.5	-	28d11 ^h /29d16 ^h	9.2/2·10 ³	3d	
MEPED	>6.9	-	28d11 ^h /29d16 ^h	3/320	3d	
MEPED	>16	-	28d11 ^h /29d16 ^h	0.8/16	3d	
MEPED	>36	-	28d11 ^h /29d16 ^h	0.4/0.7	3d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	28d11 ^h /29d16 ^h	25/7.2·10 ³	3d	
MKL	>14	-	28d11 ^h /29d16 ^h	1.5/6.4	3d	
MKL	>26	-	28d11 ^h /29d16 ^h	0.7/2.3	3d	
MKL	>50	-	28d11 ^h /29d16 ^h	0.5/0.9	3d	
ACE						
SIS	>10	04 ^h	28d11 ^h /29d16 ^h	2.8/70	2.5d	
SIS	>30	03 ^h	28d11 ^h /29d16 ^h	1.2/2.5	2.5d	
SOHO						
EPHIN INT	>50	01 ^h	28d11 ^h /29d16 ^h	0.1/0.4	2d	

Differential fluxes of protons for the event of 2003 May 28

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	06 ^h	28d11 ^h /29d16 ^h	40.5/428	2d	
LION	2-6	06 ^h	28d11 ^h /29d16 ^h	6.5/62.7	2d	
EPHIN	4-8	05 ^h	28d11 ^h /29d16 ^h	5.4 /180	2d	
EPHIN	8-25	02 ^h	28d11 ^h /29d16 ^h	0.12 /9.2	2d	
EPHIN	25-41	01 ^h	28d11 ^h /29d16 ^h	0.013 /0.09	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:




Kuwabara T., J.W. Bieber, J. Clem et al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2003 May 28

2003 May 28			• AR10365					To event 436
Hα	6563 Å	2258	0047	0245	S06W20	2B	EFHTU	
1 – 12	keV	2256	2307	2313		X1.3	7.1E-2	
1 – 12	keV	0017	0027	0039		X3.6	2.8E-1	
6–12	keV	230532	230610	231020		377798	RHESSI	
6–12	keV	231020	231642	232656		299981	RHESSI	
6–12	keV	23:25:56	23:27:50	23:30:20		28274	RHESSI	
6–12	keV	23:30:20	23:35:02	23:36:56		32609	RHESSI	
6–12	keV	23:36:56	23:38:30	23:41:48		28879	RHESSI	
25–50	keV	23:41:48	23:43:46	23:47:32		287482	RHESSI	
6–12	keV	23:47:32	23:48:38	23:49:20		10231	RHESSI	
6–12	keV	00:01:12	00:01:42	00:03:44		20241	RHESSI	
6–12	keV	00:42:32	00:43:22	00:45:24		1956	RHESSI	
25-50	keV	00:45:24	01:03:34	01:45:12		9968246	RHESSI	
50–150	keV	00:21		00:29			SONG F	
4–7	MeV		>0021				SONG F	

15.4	GHz	2301.0	2306.0	2318.0		3.08	
8.8	GHz	2259.0	2305.0	2327.0	P0.6 U1.4 P5-9	3.32	
5	GHz	2259.0	2306.0	2326.0		3.32	
2.8	GHz	2248.0	2306.8	2345.0		2.97	
1.4	GHz	2300.0	2301.0	2321.0		2.53	
610	MHz	2300.0	2309.0	2332.0		3.72	
410	MHz	2259.0	2304.0	2336.0		3.59	
245	MHz	2259.0	2303.0	2336.0		3.45	
DS II	50-300	2302		2310	F,N	3	
DS II	25-180	2306		2316		3	
DS IV	200-1400	2301		2317	F,S	2	
DS IV	25-180	2311		0145		2	
DS III	20-1400	2300		2307	GG	3	
15.4	GHz	0020.0	0023.0	0107.0	U1.4-2.8 / 15.4	3.76	
8.8	GHz	0019.0	0024.0	0109.0		3.72	
5	GHz	0020.0	0025.0	0154.0		3.57	
2.8	GHz	0013.0	0022.9	0036.0		3.29	
1.4	GHz	0021.0	0023.0	0121.0		3.32	
610	MHz	0021.0	0027.0	0124.0		4.91	
410	MHz	0021.0	0043.0	0207.0		4.72	
245	MHz	0022.0	0022.0	0139.0		5.85	
DS II	25-180	0026		0033		3	
DS IV	40-1600	0022		0241	F,S	3	
DS III	18-1800	0022		0027	G	3	
8.8	GHz	0228.0	0245.0	0248.0		1.87	
5	GHz	0218.0	0222.0	0246.0		1.94	
2.7	GHz	0218.0	0222.0	0225.0		2.00	
1.4	GHz	0215.0	0222.0	0229.0		2.23	
610	MHz	0207.0	0217.0	0239.0		4.00	
410	MHz	0207.0	0217.0	0300.0		4.68	
245	MHz	0207.0	0217.0	0300.0	0.2 \ 8.8	4.71	
°n						Mauna Kea, Norikura	
CME	WL	2350	0964 km/s	-9.6 km/s ²	360°	300°	2350
CME	WL	0050	1366 km/s	25.9	360	292	0050

* One flare event with two X-ray bursts




2003 May 29

Ø

AR10365

To event 436

H α	6563 Å	0020	0105	0223	S06W37	2B	FU
1 – 12	keV	0051	0105	0112		X1.3	6.8E-2
6-12	keV	004232	004322	004524		1956	RHESSI
25-50	keV	004524	010334	014512		9968246	RHESSI
12-25	keV	022040	022046	022440		106590	RHESSI
15.4	GHz	0059.0	0105.0	0213.0		3.34	
8.8	GHz	0058.0	0105.0	0217.0	[P0.4] U2.7 P8.8	3.53	
5	GHz	0058.0	0105.0	0217.0		3.40	
2.7	GHz	0058.0	0105.0	0213.0		2.91	
1.4	GHz	0058.0	0156.0	0158.0		3.23	
610	MHz	0058.0	0152.0	0215.0		3.94	
410	MHz	0059.0	0152.0	0212.0		4.51	
245	MHz	0058.0	0150.0	0222.0		4.49	
DS II	30-90	0106		0111	F,N	3	
DS IV	40-1800	0058		0213	F,S	2	
DS III	18-1000	0100		0106	G	3	
DS III	18-90	0108		0111	G	3	
DS III	30-300	0213		0216	G	3	
DS UNCLF	40-120	0113		0116		2	
CME	WL	0051	1237 km/s	-22.3km/s ²	360°	260°	



Particle event: To($E_p > 10$ MeV) – 31d03^h

Tmax($E_p > 10$ MeV) – 31d06^h, Jmax ($E_p > 10$ MeV) – 15.6 /cm².s.sr

Duration of the event – 2 days


Quasimaximal energy of protons in the event – $E_{qm} = 415$ MeV

Sources: • solar flare 31d02^h13^m, M9.3/2B, S07W65 AR10365

Main x-ray burst 1-8 Å: onset – 31d02^h13^m, max – 31d02^h24^m, $\Phi = 0.085$ Jo.m⁻²



CME: 31d02^h30^m, $V = 1835$ km/s, $\Delta\phi = 360^\circ$, $dA = 256^\circ$

Particle fluxes and associated phenomena




May - June 2003

Time profiles of the proton fluxes for the event of 2003 May 31



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 May 31

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	03 ^h	06 ^h	32.2	2d	
EPS	>10	03 ^h	06 ^h	15.6	2d	
EPS	>30	03 ^h	06 ^h	4.8	2d	
EPS	>50	03 ^h	04 ^h	2.3	2d	
EPS	>60	03 ^h	04 ^h	1.7	2d	
EPS	>100	03 ^h	04 ^h	0.72	1d	
METEOR						
CBM	>5	04 ^h	05 ^h	22	2d	
CBM	>10	04 ^h	05 ^h	10	1.5d	
CBM	>25	04 ^h	05 ^h	4.5	1d	
CBM	>40	04 ^h	05 ^h	2.7	1d	
BP	>90	04 ^h	04 ^h	0.5	1d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	16 ^h	6·10 ³	2d	
MEPED	>0.8	-	16 ^h	1.2·10 ³	2d	
MEPED	>2.5	-	16 ^h	220	2d	
MEPED	>6.9	-	08 ^h	32	2d	
MEPED	>16	-	06 ^h	15	2d	
MEPED	>36	-	06 ^h	7.5	2d	
MEPED	>70	-	04 ^h	1.2	1d	
MEPED	>140	-	04 ^h	0.4	1d	

CORONAS F						
MKL	>1.	-	08 ^h	420	2d	
MKL	>14	-	07 ^h	7.3	2d	
MKL	>26	-	05 ^h	5.4	2d	
MKL	>50	-	05 ^h	2.6	1d	
ACE						
SIS	>10	03 ^h	05 ^h	25.5	2d	
SIS	>30	03 ^h	06 ^h	9	2d	
SOHO						
EPHIN INT	>50	03h	05h	1.65	1d	


Differential fluxes of protons for the event of 2003 May 31

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	08 ^h	13 ^h	40	5d	
LION	2-6	08 ^h	13 ^h	7	5d	
EPHIN	4-8	05 ^h	10 ^h	22.8	5d	
EPHIN	8-25	05 ^h	09 ^h	1.2	4d	
EPHIN	25-41	03 ^h	05 ^h	0.15	4d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2003 May 31

2003	May 31	•	AR10365	To event 437			
H α	6563 Å	0218	0223	0338	S07W65	2B	U
1 – 12	keV	0213	2224	0240		M9.3	8.5E-2
100 – 300	keV	022536	022702	023416		15794196	RHESSI
12-25	keV	030048	030118	031840		630884	RHESSI
12-25	keV	031840	031958	032444		112050	RHESSI

15.4	GHz	0217.0	0222.0	0247.0	[U2.7]/15.4	3.93	
8.8	GHz	0216.0	0223.0	0251.0		3.90	
5	GHz	0216.0	0224.0	0248.0		3.61	
2.7	GHz	0216.0	0225.0	0247.0		3.18	
1.4	GHz	0217.0	0224.0	0234.0		3.26	
610	MHz	0218.0	0220.0	0238.0		3.43	
410	MHz	0220.0	0226.0	0248.0		3.30	
245	MHz	0214.0	0222.0	0249.0		4.36	
DS II	35-180	0223		0235	S,H	3	
DS II	75-180	0223		0000		3	
DS IV	35-1800	0218		0254		2	
DS III	350-1000	0220		0221	G	2	
DS III	18-300	0221		0223	G	3	
DS V	25-180	0220		0227		3	
CME	WL	0230	1835 km/s	-2.4 km/s ²	360°	256°	



Particle event: To($E_p > 10 \text{ MeV}$) – 18d08^h
 Tmax ($E_p > 10 \text{ MeV}$) – 20d06^h, Jmax ($E_p > 10 \text{ MeV}$) – $10.2 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$
 Duration of the event – 2 days
 Quasimaximal energy of protons in the event – $E_{qm} = 180 \text{ MeV}$

Sun source: ☉ solar flare 17d22^h 27^m, M6.8/..., s12e60*, AR10368


Main X-ray burst 1-8 Å: onset – 17d22^h 27^m, max – 17d22^h 55^m, $\Phi = 0.096 \text{ Jo} \cdot \text{m}^{-2}$

CME: 17d23^h 18^m, $V = 1813 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 117^\circ$



$\Delta\text{SC } 18\text{d}05^{\text{h}}12^{\text{m}}$

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2003 June 18



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 June 18

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	06 ^h	19d05 ^h	48	3d	
EPS	>10	06 ^h	19d05 ^h	10.2	3d	
EPS	>30	06 ^h	19d05 ^h	0.23	3d	
EPS	>50	06 ^h	19d05 ^h	0.1	3d	
EPS	>60	06 ^h	19d05 ^h	0.09	3d	
EPS	>100	06 ^h	19d05 ^h	0.06	3d	
POES-16						
MEPED	>0.24	-	19d05 ^h	1400	2d	
MEPED	>0.8	-	19d05 ^h	800	2d	
MEPED	>2.5	-	19d05 ^h	340	2d	
MEPED	>6.9	-	19d05 ^h	70	2d	
MEPED	>16	-	19d05 ^h	3	2d	
MEPED	>36	-	-	-	2d	
MEPED	>70	-	-	-	1d	
MEPED	>140	-	-	-	1d	
CORONAS F						
MKL	>1.	-	19d05 ^h	471	2d	
MKL	>14	-	19d05 ^h	1.2	2d	
MKL	>26	-	19d05 ^h	0.25	2d	
MKL	>50	-	19d05 ^h	0.21	1d	
ACE						
SIS	>10	07 ^h	19d05 ^h	15	3d	
SIS	>30	07 ^h	19d05 ^h	0.2	1d	
SOHO						
EPHIN INT	>50	-	-	-	-	

Differential fluxes of protons for the event of 2003 June 18

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	06 ^h	19d05 ^h	22.4	8d	
LION	2-6	06 ^h	19d05 ^h	6	8d	
EPHIN	4-8	06 ^h	19d05 ^h	18.9	8d	
EPHIN	8-25	06 ^h	19d05 ^h	1.2	8d	
EPHIN	25-41	06 ^h	19d03 ^h	0.07	3d	
EPHIN	41-53	- " -	- " -	- " -	- " -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2003 June 18

2003 June 17


•

AR10386

To event 438

H α	6563 Å	No Flare Patrol			s12e53		
1 – 12	keV	2227	2255	2312		M6.8	9.6E-2
50-100	keV	22232	224946	230716		35645838	RHESSI
15.4	GHz	2241.0	2253.0	2312.0		3.40	
8.8	GHz	2239.0	2253.0	2316.0		3.53	
5	GHz	2238.0	2253.0	2318.0	U0.6 P5	3.57	
2.7	GHz	2239.0	2254.0	2322.0		3.32	
1.4	GHz	2238.0	2245.0	2322.0		2.80	
610	MHz	2239.0	2246.0	2329.0		2.71	
410	MHz	2239.0	2315.0	2334.0		2.79	
245	MHz	2243.0	2245.0	2334.0		3.08	
DS II		2248		2300		2	
DS II		2255		2256		3	
DS IV		2243		2336		2	
DS III	B	2244		2244	B	3	
DS III	GG	2245		2259	GG	2	
CME	WL	2318	1825 km/s	-2.9 km/s ²	360°	117°	

* – probable localization of the flare event



Particle event: To($E_p > 10$ MeV) – 26d18^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 26\text{d}20^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 230 \text{ /cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 27\text{d}02^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 360 \text{ /cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm1} = 340 \text{ MeV}$

– $E_{qm2} = 400 \text{ MeV}$

Sources: ● solar flare 26d17^h21^m, X1.2/1N, N02W38 AR10484


○ solar flare 27d09^h21^m, M5.0/SF, S15E26 AR10486

Main X-ray burst 1-8 Å: onset – 26d17^h21^m, max – 26d18^h19^m, $\Phi = 0.63 \text{ J/m}^2$



CME: 26d17^h54^m; $V = 1754 \text{ km/s}$; $\Delta\phi = 171^\circ$; $dA = 235^\circ$

▲ SC 26d19^h08^m;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2003 October 26



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 October 26

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	18 ^h	20 ^h /27d02 ^h	354/758	2d	
EPS	>10	18 ^h	20 ^h /27d02 ^h	230/360	2d	
EPS	>30	18 ^h	20 ^h /27d02 ^h	30.3/38	2d	
EPS	>50	18 ^h	20 ^h /27d02 ^h	5.9/7.3	2d	
EPS	>60	18 ^h	19 ^h /27d02 ^h	2.5/3.7	2d	
EPS	>100	18 ^h	19 ^h /27d02 ^h	0.46/0.83	2d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	19 ^h /27d04 ^h	1.8·10 ³ /1.3·10 ³	2d	
MEPED	>6.9	-	19 ^h /27d04 ^h	1.4·10 ³ /520	2d	
MEPED	>16	-	19 ^h /27d04 ^h	230/240	2d	
MEPED	>36	-	19 ^h /27d04 ^h	30/35	2d	
MEPED	>70	-	19 ^h /27d04 ^h	2/2.2	2d	
MEPED	>140	-	- /27d04 ^h	- /0.6	2d	
CORONAS F						
MKL	>1.	-	23 ^h /27d05 ^h	2.2·10 ³ /1.4·10 ³	2d	
MKL	>14	-	23 ^h /27d05 ^h	240/220	2d	
MKL	>26	-	22 ^h /27d05 ^h	125/120	2d	
MKL	>50	-	22 ^h /27d05 ^h	40/17	2d	

ACE						
SIS	>10	17 ^h	21 ^h /27d03 ^h	273/352	2d	
SIS	>30	17 ^h	19 ^h /27d03 ^h	26/55.5	2d	
SOHO						
EPHIN (INT)	>50	17 ^h	18 ^h /27d03 ^h	16.6/4.6	1.5d	
BALLOONS						
Mu	>96		- /27d10 ^h	- /0.29		
Mu	>100		- /27d10 ^h	- /0.24		

Differential fluxes of protons for the event of 2003 October 26

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	20 ^h	23 ^h /27d06 ^h	38/42	2d	
LION	2-6	19 ^h	23 ^h /27d05 ^h	34/22.5	2d	
EPHIN	4-8	19 ^h	22 ^h /27d06 ^h	212/150	2d	
EPHIN	8-25	18 ^h	22 ^h /27d05 ^h	39.6/22.3	2d	
EPHIN	25-41	18 ^h	18 ^h /27d03 ^h	2.8/1.4	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2003 October 26


2003	October 26	•	AR10484	To event 439			
H α	6563 Å	1721	1733	2016	N02W38	1N	F
1 – 12	keV	1721	1819	1921		X1.2	6.3E-1
15.4	GHz	1715.0	1731.0	1919.0	U0.4 / 15	3.53	
8.8	GHz	1715.0	1731.0	1908.0		3.65	
5	GHz	1712.0	1731.0	1920.0		3.62	
2.7	GHz	1715.0	1731.0	1920.0		3.40	
1.4	GHz	1718.0	1756.0	1820.0		3.20	
610	MHz	1720.0	1758.0	1826.0		2.77	
410	MHz	1720.0	1811.0	1849.0		2.63	
245	MHz	1722.0	1724.0	1908.0		3.08	
DS II	25-100	1735		1743		2	
DS CONT	30-45	1804		1853		1	
CME	WL	1754	1537 km/s	4.8km/s ²	171°	235°	

HESSI - gap



2003 October 26 • AR10486 To event 439

H α	6563 Å	0923	0925	0946	S15E26	SF	F
1 – 12	keV	0921	0927	0932		M5.0	2.6E-2
15.4	GHz	0922.0	0923.0	0931.0	3 / 15	3.66	
8.8	GHz	0905.0	0923.0	0938.0		3.45	
5	GHz	0922.0	0923.0	0927.0		2.88	
3	GHz	0922.9	0923.6	0926.2		1.82	
DS III	25-155	0922		0923	G,C	2	
DS DCIM	800-4000	0917		0926	C	3	



Particle event: To($E_p > 10$ MeV) – 28d12^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 28\text{d}18^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 4.6 \cdot 10^3 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 29\text{d}02^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 1.2 \cdot 10^4 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 1 day

Quasimaximal energy of protons in the event – $E_{qm1} = 3340$ MeV

– $E_{qm2} = 1025$ MeV


Sources • solar flare 28d09^h51^m, X17.2/4B, S16E08, AR10486

Main X-ray burst 1-8 Å: onset – 28d09^h51^m, max – 28d11^h10^m, $\Phi = 1.8 \text{ J/m}^2$



CME: 28d11^h30^m; $V = 2459 \text{ km/s}$; $\Delta\phi = 360^\circ$; $dA = 015^\circ$

▲ SC 28d19^h08^m; ▲ SC 29d06^h11^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2003 October 28



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 October 28

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	12 ^h	18 ^h /29d02 ^h	5.5·10 ³ /1.4·10 ⁴	1d	
EPS	>10	12 ^h	18 ^h /29d02 ^h	4.6·10 ³ /1.2·10 ⁴	1d	
EPS	>30	12 ^h	17 ^h /29d02 ^h	1.8·10 ³ /4.4·10 ³	1d	
EPS	>50	12 ^h	17 ^h /29d01 ^h	790/1.45·10 ³	1d	
EPS	>60	12 ^h	17 ^h /29d01 ^h	510/820	1d	
EPS	>100	12 ^h	16 ^h /29d01 ^h	150/185	1d	
POES-16						
MEPED	>0.24	-	- /29d07 ^h	- /8.1·10 ⁴	1d	
MEPED	>0.8	-	- /29d07 ^h	- /5.2·10 ⁴	1d	
MEPED	>2.5	-	16 ^h /29d07 ^h	1.1·10 ⁴ /4.1·10 ⁴	1d	
MEPED	>6.9	-	16 ^h /29d07 ^h	8.7·10 ³ /3.4·10 ⁴	1d	
MEPED	>16	-	16 ^h /29d07 ^h	4.2·10 ³ /1.1·10 ⁴	1d	
MEPED	>36	-	16 ^h /29d07 ^h	1.4·10 ³ /3.8·10 ³	1d	
MEPED	>70	-	16 ^h /29d07 ^h	310/ 120	1d	
MEPED	>140	-	16 ^h /29d07 ^h	33/ 9	1d	
CORONAS F						
MKL	>1.	-	15 ^h /29d07 ^h	1.5·10 ⁴ /6.2·10 ⁴	1d	
MKL	>14	-	15 ^h /29d07 ^h	2.1·10 ³ /8.6·10 ³	1d	
MKL	>26	-	15 ^h /29d07 ^h	1.0·10 ³ /5.3·10 ³	1d	
MKL	>50	-	15 ^h /29d07 ^h	230/1.6·10 ³	1d	

ACE						
SIS	>10	12 ^h	17 ^h /29d02 ^h	3.9·10 ³ /4.9·10 ³	1d	
SIS	>30	12 ^h	17 ^h /29d01 ^h	1.6·10 ³ /1.9·10 ³	1d	
SOHO						
EPHIN (INT)	>50	12 ^h	17 ^h /29d02 ^h	704/1580	1d	
BALLOONS						
Mu	>160	-	- /23 ^h ÷29d01 ^h	- /13.4	-	
Mu	>242	-	- /23 ^h ÷29d01 ^h	- /3	-	
Mu	>348	-	- /23 ^h ÷29d01 ^h	- /0.32	-	
NM						
Network	>433	-	11 ^h 55 ^m / -	7.6/ -	-	
Network	>1000	-	11 ^h 55 ^m / -	0.44/ -	-	
Network	>2000	-	11 ^h 55 ^m / -	0.042/ -	-	
Network	>3700	-	11 ^h 55 ^m / -	0.005/ -	-	
Network	>5800	-	11 ^h 55 ^m / -	0.0009/ -	-	

Differential fluxes of protons for the event of 2003 October 28

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	14 ^h	17 ^h /29d02 ^h	50/150	1.5d	
LION	2-6	14 ^h	17 ^h /29d01 ^h	48/102	1.5d	
EPHIN	4-8	13 ^h	17 ^h /29d01 ^h	77/87	1.5d	
EPHIN	8-25	12 ^h	17 ^h /29d01 ^h	39/64	1.5d	
EPHIN	25-41	12 ^h	17 ^h /29d02 ^h	16/36	1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

- Struminsky A., 2005.
 Bieber J.W., J. Clem, P. Evenson et al., 2005.
 Miroshnichenko L.I., K.-L. Klein, G. Trotter et al., 2005.
 Plainaki C., A. Belov, E. Eroshenko et al., 2005.
 Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
 Perez-Peraza J., A. Gallegos-Cruz, E.V. Vashenyuk et al., 2006.
 Dmitriev A.V., H.-C. Yeh, J.-K. Chao et al., 2006.
 Nonaka T., Y. Hayashi, N. Ito et al., 2006.
 Miroshnichenko L.I. and J. Perez-Peraza, 2008.
 Lario D., R.B. Decker, and A. Aran, 2008.
 Struminsky A., B. Heber, R. Gomez-Herrero et al., 2008.
 Lario D, A. Aran, R.B. Decker, 2009.
 Troitskaia E.V., I.V. Arkhangelskaja, L.I. Miroshnichenko et al., 2009.
 Perez-Peraza J., A. Gallegos-Cruz, E.V. Vashenyuk, 2009.
 Kichigin G.N., L.I. Miroshnichenko, V.I. Sidorov et al., 2012.
 Miroshnichenko L.I., J.A. Pérez-Peraza, V.M. Velasco-Herrera et al., 2012.
 Miroshnichenko L.I. and W.Q. Gan, 2012.
 Mauricev E.A., Yu.V. Balabin, E.V. Vashenyuk et al., 2013.
 Velinov P., A. Mishev, 2013.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2003 October 28**


2003 October 28

•

AR10486

To event 440

H α	6563 Å	<1001	~1205	>1420	S16E08	4B	FU
1 – 12	keV	0951	1110	1124		X17.2	1.8E00
25-50	keV	094016	095322	095400		271440	RHESSI
25-50	keV	102832	103530	103700		14146876	RHESSI
800-7000	keV	103700	111410	113012		14146876	RHESSI
50–150	keV	1102		1113			SONG-F
60–100	MeV		>1103				SONG-F
15.4	GHz	1004.0	1104.0	1456.0		4.76	
8.8	GHz	1003.0	1107.0	1456.0	U1.4-2.7 P 8.8	4.85	
5	GHz	1004.0	1112.0	1448.0		4.67	
2.7	GHz	1014.0	1114.0	1459.0		4.11	
1.4	GHz	1131.0	1146.0	0000.0		4.11	
610	MHz	1014.0	1117.0	1508.0		4.49	
410	MHz	1005.0	1104.0	1508.0		4.57	
245	MHz	1015.0	1102.0	1508.0		5.69	
DS II	100-1400	1100		1129		3	
DS II	25-180	1102		1111		3	
DS IV	100-4000	1012		1515	P,C,F,S,	3	
DS IV	25-180	1033		1531		3	
DS III	400-4000	0949		0955	GG,C	3	
DS III	25-270	1013		1031	GG,F,S	2	
DS III	25-270	1058		1111	GG,C	3	
DS CONT	50-270	~1030		>1224		3	
DS DCIM	2000-4500	1005		1236	GG	3	
DS DCIM	800-2000	1058		1336	GG,S,P,F,S	3	
n°						CORONAS-F	
n°						Tsumed, (S, A, T)	
CME	WL	1130	2459 km/s	-105.2 km/s ²	360°	015°	1130



Particle event: To($E_p > 10$ MeV) – 29d22^h

Tmax($E_p > 10$ MeV) – 29d23^h, Jmax($E_p > 10$ MeV) – 2230 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm} = 810$ MeV

Sources: ● solar flare 29d20^h37^m, X10.0/2B, S15W02, AR10486


○ solar flare 30d01^h53^m, 1F/M1.6, N08W22, AR10488

Main X-ray burst 1-8 Å: onset – 29d20^h37^m, max – 29d20^h49^m, $\Phi = 0.87$ J/m²



CME: 29d20^h54^m; $V = 2459$ km/s ; $\Delta\phi = 360^\circ$; $dA = 190^\circ$

▲ SC 29d06^h11^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2003 October 29



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 October 29

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm².s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	22 ^h	30d01 ^h	3520	2.5d	
EPS	>10	22 ^h	23 ^h	2230	2.5d	
EPS	>30	21 ^h	23 ^h	707	2d	
EPS	>50	21 ^h	23 ^h	320	2d	
EPS	>60	21 ^h	23 ^h	230	2d	
EPS	>100	21 ^h	23 ^h	91	2d	
METEOR						
CBM	>5	20 ^h	30d01 ^h	920	2.5d	
CBM	>10	20 ^h	30d00 ^h	640	2.5d	
CBM	>25	20 ^h	23 ^h	460	2.5d	
CBM	>40	20 ^h	23 ^h	330	2d	
BP	>90	20 ^h	23 ^h	90	2d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	-	-	-	.
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	23 ^h	8120	2.5d	
MEPED	>16	-	23 ^h	2520	2.5d	
MEPED	>36	-	23 ^h	930	2.5d	
MEPED	>70	-	23 ^h	130	2d	
MEPED	>140	-	23 ^h	35	2d	

CORONAS F						
MKL	>1.	-	24 ^h	9240	2.5d	
MKL	>14	-	24 ^h	1550	2d	
MKL	>26	-	24 ^h	750	2d	
MKL	>50	-	24 ^h	180	2d	
ACE						
SIS	>10	22 ^h	23 ^h	1840	2d	
SIS	>30	22 ^h	23 ^h	680	2d	
SOHO						
EPHIN (INT)	>50	21 ^h	30d01 ^h	320	1.5d	
BALLOONS						
Mu	>141	-	20 ^h 30 ^m	1.9	-	Before
Mu	>183	-	20 ^h 30 ^m	1	-	maximum
Mu	>238	-	20 ^h 30 ^m	0.22	-	
NM						
Network	>500	-	21 ^h 30 ^m	2.5	-	

Differential fluxes of protons for the event of 2003 October 29

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	22 ^h	30d00 ^h	220	2d	
LION	2-6	22 ^h	30d00 ^h	70	2d	
EPHIN	4-8	22 ^h	30d00 ^h	210	2d	
EPHIN	8-25	22 ^h	30d00 ^h	48	2d	
EPHIN	25-41	22 ^h	30d00 ^h	8.7	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References;

Kuwabara T., K. Munakata, S. Yasue et al., 2004.
Dmitriev A.V., H.-C. Yeh, J.-K. Chao et al., 2006.
Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Miroshnichenko L.I. and J. Perez-Peraza, 2008.
Lario D., A. Aran, R.B. Decker, 2009.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.
Lario D., R.B. Decker, and A. Aran, 2008.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2003 October 29

2003		October 29		•	AR10486		To event 441	
Hα	6563 Å	2037	2042		2253	S15W02	2B	UZ
1 – 12	keV	2037	2049		2101		X10.0	8.7E-1
100–300	keV	203556	2048:4		210824		247490912	RHESSI
50–150	keV	2038			2055			SONG-F
4–7	MeV		>1103					SONG-F

8.8	GHz	2039.0	2043.0	2052.0	U1.4 / 8.8	4.04	
5	GHz	2039.0	2043.0	2053.0		3.67	
2.7	GHz	2039.0	2041.0	0000.0		3.40	
1.4	GHz	2039.0	2043.0	2057.0		3.23	
610	MHz	2040.0	2054.0	2059.0		4.76	
410	MHz	2041.0	2055.0	2059.0		4.64	
245	MHz	2042.0	2042.0	0000.0		5.56	
DS II	60-430	2042		~2050	F,N	3	
DS II	25-180	2044		2053		3	
DS IV	25-1200	2057		2220		3	
DS III	18-380	2042		2044	G	3	
DS III	120-1200	2049		2100	GG	3	
CME	WL	2054	2029 km/s	-146.5.2 km/s ²	360°	190°	



Particle event: To($E_p > 10$ MeV) – 02d17^h

Tmax($E_p > 10$ MeV) – 02d23^h, Jmax ($E_p > 10$ MeV) – 990 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm} = 1700$ MeV

Sources: ● solar flare 02d17^h03^m, X8.3/2B, S14W56, AR10486

○ solar flare 01d22^h26^m, M3.2/1N, S12W60, AR10486 *

○ solar flare 03d01^h06^m, 2B/X2.3, N10W83, AR10488


Main X-ray burst 1-8 Å: onset – 02d17^h03^m, max – 02d17^h25^m, $\Phi = 0.91$ J/m²

CME: 02d17^h30^m, $V = 2598$ km/s; $\Delta\phi = 360^\circ$; $dA = 265^\circ$



▲ SC: 04d06^h25^m

* - Contribution of event beginning

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2003 November 02



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 November 02

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	17 ^h	23 ^h	1220	2d	
EPS	>10	17 ^h	23 ^h	990	2d	
EPS	>30	17 ^h	22 ^h	320	2d	
EPS	>50	17 ^h	20 ^h	110	2d	
EPS	>60	17 ^h	19 ^h	89.5	2d	
EPS	>100	17 ^h	19 ^h	42	2d	
METEOR						
CBM	>5	17 ^h	21 ^h	470	2d	
CBM	>10	17 ^h	21 ^h	355	2d	
CBM	>25	17 ^h	19 ^h	270	2d	
CBM	>40	17 ^h	19 ^h	214	2d	
BP	>90	17 ^h	19 ^h	53	2d	
ChD	>600					
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	21 ^h	2070	2d	
MEPED	>16	-	19 ^h	930	2d	
MEPED	>36	-	19 ^h	420	2d	
MEPED	>70	-	18 ^h	110	2d	
MEPED	>140	-	18 ^h	40	2d	
CORONAS F						
MKL	>1.	-	19 ^h	4520	2d	
MKL	>14	-	19 ^h	940	2d	
MKL	>26	-	19 ^h	540	2d	
MKL	>50	-	19 ^h	180	2d	

ACE						
SIS	>10	17 ^h	23 ^h	1630	2d	
SIS	>30	17 ^h	20 ^h	540	2d	
SOHO						
EPHIN (INT)	>50	17 ^h	19 ^h	117	2d	
BALLOONS						
Mu	>129	-	03d(00 ^h 56 ^m ÷01 ^h 32 ^m)	3	-	After
Mu	>184	-	03d(00 ^h 56 ^m ÷01 ^h 32 ^m)	1.13	-	maximum
Mu	>263	-	03d(00 ^h 56 ^m ÷01 ^h 32 ^m)	0.19	-	
NM						
Network	>433	-	18 ^h	15.5	-	
Network	>1000	-	18 ^h	0.183	-	
Network	>2000	-	18 ^h	0.00465	-	
Network	>3700	-	18 ^h	0.00018	-	

Differential fluxes of protons for the event of 2003 November 02

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	14 ^h	23 ^h	18.6	2.5d	
LION	2-6	13 ^h	23 ^h	12.4	2.5d	
EPHIN	4-8	12 ^h	22 ^h	47	2.5d	
EPHIN	8-25	12 ^h	22 ^h	28.7	2.5d	
EPHIN	25-41	18 ^h	22 ^h	8.2	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Kuznetsov S.N., B.Yu. Yushkov, and K. Kudela, 2007.
Miroshnichenko L.I. and J. Perez-Peraza, 2008.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.
Maurchev E.A., Yu.V. Balabin, E.V. Vashenyuk et al., 2013.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2003 November 02


2003	November 02	●	AR10486	To event 442			
Hα	6563 Å	1704	1717	1954	S14W56	2B	UZ
1 – 12	keV	1703	1725	1739		X8.3	9.1E-1
800-7000	keV	170916	172246	173028		434251584	RHESSI
6-12	keV	173028	174230	174436		5220713	RHESSI
15.4	GHz	1703.0	1719.0	1830.0	U0.6 / 15	4.62	
8.8	GHz	1704.0	1721.0	1818.0		4.30	
5	GHz	1703.0	1722.0	1803.0		4.08	
2.7	GHz	1703.0	1717.0	1758.0		4.00	
1.4	GHz	1711.0	1720.0	1836.0		3.97	
610	MHz	1712.0	1718.0	1836.0		3.85	
410	MHz	1711.0	1715.0	1836.0		4.08	
245	MHz	1714.0	1715.0	1835.0		4.40	

DS II	25-180	1714		1737		3	
DS IV	30-80	1714		1824		3	
DS III	25-180	1714		1800	N	2	
°n							Bolivia
CME	WL	1730	2598 km/s	-32.4 km/s ²	360°	265°	



2003 November 01 Ø AR10486 To event 442

H α	6563 Å	2228	2234	2312	S12W60	1N	FU
1 – 12	keV	2226	2238	2249		M3.2	2.8E-2
50-100	keV	222456	223426	224812		14352336	RHESSI
15.4	GHz	2229.0	2233.0	2242.0		2.23	
8.8	GHz	2228.0	2230.0	2242.0	U1.4 P8.8	2.75	
5	GHz	2229.0	2230.0	2239.0		2.57	
2.7	GHz	2230.0	2232.0	2234.0		1.98	
1.4	GHz	2348.0	2348.0	~2348.0		1.92	
610	MHz	2345.0	2351.0	2354.0		2.20	
410	MHz	2237.0	2237.0	2242.0		2.60	
245	MHz	2234.0	2235.0	2240.0		4.95	
DS II	25-290	2224		2257		3	
DS II	20-45	2248		2302	UE	3	
DS II	25-180	2336		2255		1	
DS IV	70-150	2243		2333		2	
DS III	18-330	2234		2235	G	3	
DS III	20-200	2239		>2400	S,C	2	
DS CONT	200-750	2345		2356		1	
DS UNCLF	180-430	2237		2240		1	
DS UNCLF	18-30	2307		2313		3	
CME	WL	2306	0899km/s	-26.3km/s ²	360°	224°	




2003 November 03

Ø

AR10488

To event 442

H α	6563 Å	0106	0131	0237	N10W83	2B	EF
1 – 12	keV	0109	0130	0145		X2.7	3.6E-1
25-50	keV	010120	012526	020128		354025280	RHESSI
6-12	keV	023744	023746	023828		64704	RHESSI
50–150	keV	0117		0129			SONG F
0.5–1.3	MeV		>0117				SONG F
15.4	GHz	0101.0	0121.0	0202.0	U1.4 / 15	2.77	
8.8	GHz	0101.0	0121.0	0147.0		2.72	
5	GHz	0059.0	0121.0	0149.0		2.64	
2.7	GHz	0103.0	0121.0	0134.0		2.51	
1.4	GHz	0101.0	0103.0	0126.0		2.28	
610	MHz	0059.0	0122.0	0151.0		2.76	
410	MHz	0117.0	0148.0	0148.0		2.65	
245	MHz	0105.0	0105.0	0201.0		3.08	
DS II	50-200	0124		0129		3	
DS IV	20-850	0108		0136		1	
DS IV	25-115	0131		0406		1	
DS III	18-340	0108		0118	G	3	
DS III	18-250	0133		0226	S,C	3	
n°						Mauna Kea, Haleakala	
CME	WL	0159	0827 km/s	-28.3 km/s ²	065°	324°	



Particle event: To($E_p > 10$ MeV) – 04d22^h

Tmax($E_p > 10$ MeV) – 05d07^h, Jmax ($E_p > 10$ MeV) – 126 /cm².s.sr

Duration of the event – 4 days

Quasimaximal energy of protons in the event – $E_{qm} = 445$ MeV


Sources: ● solar flare 04d19^h29^m, X>17.5/3B, S19W83, AR10486

○ solar flare 05d10^h46^m, M5.3/SF, S16W90, AR10486



Main X-ray burst 1-8 Å: onset – 04d19^h29^m, max – 04d19^h50^m, $\Phi = 2.3$ J/m²

CME: 04d19^h54^m, V = 2657 km/s, $\Delta\phi = 360^\circ$, dA = 260°

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2003 November 04



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 November 04

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	23 ^h	05d07 ^h	212	4d	
EPS	>10	22 ^h	05d07 ^h	126	4d	
EPS	>30	22 ^h	05d07 ^h	27	4d	
EPS	>50	21 ^h	05d06 ^h	7.9	4d	
EPS	>60	21 ^h	05d06 ^h	4.2	3d	
EPS	>100	20 ^h	05d06 ^h	0.9	3d	
METEOR						
CBM	>5	23 ^h	05d08 ^h	137	4d	
CBM	>10	22 ^h	05d08 ^h	71	4d	
CBM	>25	22 ^h	05d08 ^h	29	3d	
CBM	>40	22 ^h	05d06 ^h	12	3d	
BP	>90	22 ^h	05d06 ^h	0.76	2d	
ChD	>600	22 ^h	05d06 ^h	0.07	1d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	05d06 ^h	1420	4d	
MEPED	>6.9	-	05d06 ^h	710	4d	
MEPED	>16	-	05d06 ^h	220	4d	
MEPED	>36	-	05d06 ^h	65	4d	
MEPED	>70	-	05d06 ^h	3.6	3d	
MEPED	>140	-	05d06 ^h	0.4	3d	

CORONAS F						
MKL	>1.	-	05d07 ^h	2540	4d	
MKL	>14	-	05d07 ^h	160	4d	
MKL	>26	-	05d07 ^h	90	4d	
MKL	>50	-	05d07 ^h	30	3d	
ACE						
SIS	>10	19 ^h	05d08 ^h	216	4d	
SIS	>30	22 ^h	05d08 ^h	34	3d	
SOHO						
EPHIN (INT)	>50	22 ^h	05d07 ^h	3.8	3d	
BALLOONS						
Mi	>158	-	05d07 ^h	0.56	-	
Mi	>177	-	05d07 ^h	0.33	-	
Mi	>213	-	05d07 ^h	0.12	-	

Differential fluxes of protons for the event of 2003 November 04

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	23 ^h	05d06 ^h	38	5d	
LION	2-6	23 ^h	05d06 ^h	14.5	4d	
EPHIN	4-8	23 ^h	05d06 ^h	121	4d	
EPHIN	8-25	23 ^h	05d06 ^h	17	4d	
EPHIN	25-41	22 ^h	05d06 ^h	1.1	3d	


References:

Plainaki C., A. Belov, E. Eroshenko et al., 2005.
Dmitriev A.V., H.-C. Yeh, J.-K. Chao et al., 2006.
Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2003 November 04**

2003 November 04 • AR10486 To event 443

H α	6563 Å	1932	1957	2049	S19W83	3B	FZ
1 – 12	keV	1929	1950	2006		X>17.5	2.3E00
50-100	keV	192816	193822	193916		3265828	RHESSI
25-50	keV	201512	201602	202156		7718513	RHESSI
6-12	keV	202156	202354	203100		9181674	RHESSI
25-50	keV	203228	203438	204028		6256194	RHESSI
6-12	keV	204028	204202	204500		3023174	RHESSI
12-25	keV	204500	204742	211520		12714261	RHESSI
50–150	keV	1932		1957			SONG F
100–200	MeV						SONG F
15.4	GHz	1933.0	1945.0	2121.0	/ 15	4.78	
8.8	GHz	1933.0	1946.0	2058.0		4.48	
5	GHz	1933.0	1945.0	2113.0		4.36	
2.7	GHz	1933.0	1946.0	2116.0		4.30	
1.4	GHz	1933.0	1944.0	2102.0		3.90	
610	MHz	1933.0	1944.0	2101.0		4.51	
410	MHz	<1933.0	1942.0	>1943.0		3.48	
245	MHz	1932.0	1941.0	2000.0		3.68	
DS II		1943		1947		3	
DS II	30-80	<2000		2006	UE	3	
DS II	23-170	2003		2008	F,N	3	
DS IV	20-45	1947		2017		2	
DS III	25-180	2002		2013	G	3	
DS III	23-57	2047		2056	G	3	
°n						Halea	
CME	WL	1954	2657 km/s	434.8 km/s ²	360°	260°	



2003


November 05

Ø

AR10486

To event 443

H α	6563 Å	1051	1051	1058	S16W90	SF	
1 – 12	keV	1046	1052	1056		M5.3	1.7E-02
15.4	GHz	1049.0	1049.0	1050.0	3 / 15	2.77	
8.8	GHz	<1050.0	~1050.0	>1050.0		2.15	
5	GHz	<1050.0	~1050.0	>1050.0		1.67	
3	GHz	1049.4	1049.6	1050.4		1.28	
DS DCIM	2000-4000	1049		1050	C	2	



Particle event: To($E_p > 10$ MeV) – 20d08^h

Tmax($E_p > 10$ MeV) – 20d11^h, Jmax ($E_p > 10$ MeV) – 4.4 /cm².s.sr

Duration of the event – 1 day

Quasimaximal energy of protons in the event – $E_{qm} = 140$ MeV


Sources: ● solar flare 20d07^h35^m, M9.6/2B, N01W08, AR10501

Main X-ray burst 1-8 Å: onset – 20d07^h35^m, max – 20d07^h47^m, $\Phi = 0.06$ J/m²

CME: 20d08^h06^m, $V = 669$ km/s, $\Delta\phi = 360^\circ$, $dA = 219^\circ$



▲ SC: 20d08^h03^m

Particle fluxes and associated phenomena




November 2003

Time profiles of the proton fluxes for the event of 2003 November 20



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 November 20

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	06 ^h	11 ^h	21.2	1d	
EPS	>10	08 ^h	11 ^h	4.4	1d	
EPS	>30	08 ^h	11 ^h	0.35	1d	
EPS	>50	09 ^h	11 ^h	0.09	1d	
EPS	>60	09 ^h	10 ^h	0.06	1d	
EPS	>100	09 ^h	10 ^h	0.02	1d	
METEOR						
CBM	>5	08 ^h	11 ^h	4.8	1d	
CBM	>10	09 ^h	11 ^h	1.16	0.7d	
CBM	>25	10 ^h	12 ^h	0.32	0.7d	
CBM	>40	10 ^h	12 ^h	0.14	0.5d	
BP	>90	09 ^h	10 ^h	0.023	0.3d	
ChD	>600					
POES-16						
MEPED	>0.24	-	12 ^h	2570	1d	
MEPED	>0.8	-	12 ^h	550	1d	
MEPED	>2.5	-	12 ^h	110	1d	
MEPED	>6.9	-	12 ^h	40	1d	
MEPED	>16	-	12 ^h	2	1d	
MEPED	>36	-	12 ^h	0.7	1d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	12 ^h	660	1d	
MKL	>14	-	12 ^h	1.35	1d	
MKL	>26	-	12 ^h	0.9	1d	
MKL	>50	-	12 ^h	0.3	1d	
ACE						
SIS	>10	08 ^h	10 ^h	5.7	1d	
SIS	>30	08 ^h	10 ^h	1.1	1d	
SOHO						
EPHIN (INT)	>50	08 ^h	09 ^h	0.05	0.5d	

Differential fluxes of protons for the event of 2003 November 20

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	01 ^h	11 ^h	17.5	1d	
LION	2-6	02 ^h	11 ^h	2.5	1d	
EPHIN	4-8	07 ^h	10 ^h	5.3	1d	
EPHIN	8-25	07 ^h	10 ^h	0.26	1d	
EPHIN	25-41	08 ^h	10 ^h	0.01	1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Kuwabara T., J.W. Bieber, J. Clem et al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2003 November 20

2003 November 20		•		AR10501		To event 444	
H α	6563Å	0738	0740	0836	N01W08	2B	FH
1 – 12	keV	0735	0747	0753		M9.6	6.0E-2
12-25	keV	071104	071254	071624		82992	RHESSI
12-25	keV	080008	080050	082704		335064	RHESSI
12-25	keV	082704	082842	082842		19262	RHESSI
50–150	keV	0803		0816			SONG F
0.5–1.3	MeV						SONG F

15.4	GHz	0738.0	0740.0	0753.0		3.15	
8.8	GHz	0737.0	0740.0	0757.0		3.32	
5	GHz	0737.0	0739.0	0754.0		3.36	
2.7	GHz	0736.0	0739.0	0749.0	[U0.6] P2.7	3.99	
1.4	GHz	0737.0	0739.0	0744.0		2.70	
610	MHz	0735.0	0740.0	0742.0		2.71	
410	MHz	0735.0	0743.0	0745.0		2.41	
245	MHz	0738.0	0739.0	0745.0		2.45	
127	MHz	0735.7	0739.7	0744.0		2.83	
DS IV	25-113	0741		0900		1	
DS I	50-270	0739		0743	GG,DC	2	
DS I	200-400	~0758		~1442	S,N	2	
DS III	18-700	0736		0743	G	3	
DS III	25-180	0748		1506	N	2	
DS III	40-400	<0752		~0843	CG,N	2	
DS V	25-180	0738		0741		3	
DS CONT	25-270	0738		~0744		2	
CME	WL	0806	0669 km/s	-23.8km/s ²	360°	219°	



Particle event: To($E_p > 10$ MeV) – 21d08^h

Tmax($E_p > 10$ MeV) – 22d02^h, Jmax ($E_p > 10$ MeV) – $10.7 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 4 days


Quasimaximal energy of protons in the event – $E_{qm} = 80$ MeV

Sources: • solar flare 20d23^h42^m, M5.8/2B, N02W17, AR10501

Main X-ray burst 1-8 Å: onset – 20d23^h42^m, max – 20d23^h53^m, $\Phi = 0.028 \text{ J/m}^2$



CME: 21d00^h26^m; $V = 494 \text{ km/s}$; $\Delta\phi = 052^\circ$; $dA = 237^\circ$

Particle fluxes and associated phenomena




November 2003

Time profiles of the proton fluxes for the event of 2003 November 21



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 November 21

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	08 ^h	22d02 ^h	41	5d	
EPS	>10	08 ^h	22d02 ^h	11	4d	
EPS	>30	08 ^h	22d01 ^h	0.3	3d	
EPS	>50	08 ^h	22d01 ^h	0.02	2d	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
METEOR						
CBM	>5	08 ^h	22d00 ^h	5.7	3d	
CBM	>10	09 ^h	22d00 ^h	1.17	2d	
CBM	>25	09 ^h	22d00 ^h	0.36	1.5d	
CBM	>40	09 ^h	22d00 ^h	0.14	1d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	23 ^h	1740	5d	
MEPED	>0.8	-	23 ^h	870	5d	
MEPED	>2.5	-	23 ^h	320	5d	
MEPED	>6.9	-	23 ^h	35	4d	
MEPED	>16	-	23 ^h	6.5	3d	
MEPED	>36	-	23 ^h	0.5	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	24 ^h	720	5d	
MKL	>14	-	24 ^h	2	4d	
MKL	>26	-	24 ^h	0.8	3d	
MKL	>50	-	24 ^h	0.4	2d	
ACE						
SIS	>10	08 ^h	24 ^h	16.5	4d	
SIS	>30	07 ^h	24 ^h	0.6	1d	
SOHO						
EPHIN (INT)	>50	23 ^h	22d01 ^h	0.1	0.3d	

Differential fluxes of protons for the event of 2003 November 21

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	08 ^h	22d03 ^h	22	5d	
LION	2-6	08 ^h	22d02 ^h	3.8	5d	
EPHIN	4-8	07 ^h	22d02 ^h	17.8	5d	
EPHIN	8-25	07 ^h	22d02 ^h	1.4	5d	
EPHIN	25-41	07 ^h	22d01 ^h	0.027	5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:


Kuwabara T., J.W. Bieber, J. Clem, et.al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2003 November 21

2003 November 21 • AR10501 To event 445

H α	6563 Å	2346	2354	0031	N02W17	2B	F
1 – 12	keV	2342	2353	2358		M5.8	2.8E-02
6–12	keV	235628	235754	000248		35568	RHESSI
6–12	keV	000852	000918	001024		5400	RHESSI
6–12	keV	002024	002514	003420		89832	RHESSI
12-25	keV	003420	004854	005416		530784	RHESSI

15.4	GHz	2352.0	2353.0	2354.0		2.20	
8.8	GHz	2346.0	2348.0	2355.0		2.63	
5	GHz	2346.0	2348.0	0000.0		2.85	
2.8	GHz	2345.0	2349.0	2357.0	U0.4 P2.8	2.90	
1.4	GHz	2346.0	2349.0	2352.0		2.72	
610	MHz	2346.0	2347.0	2351.0		2.43	
410	MHz	2348.0	2348.0	0000.0		2.32	
245	MHz	2345.0	2348.0	0000.0		2.53	
DS III	18-1400	2347		2351	G	3	
DS III	18-420	2352		2354	G	3	
DS III	18-50	2356		2356	B	3	
DS III	25-180	2356		0248	N	1	
DS III	18-1000	0025		0030	G	3	
CME	WL	0026	0494 km/s	-3.3 km/s ²	052°	237°	



Particle event: To($E_p > 10$ MeV) – 02d12^h

Tmax($E_p > 10$ MeV) – 02d18^h, Jmax ($E_p > 10$ MeV) – 21 /cm².s.sr

Duration of the event – 4.5 days

Quasimaximal energy of protons in the event – $E_{qm} = 100$ MeV


Sources: ☉ solar flare 02d09^h40^m, C7.2/..., s19w90* AR10508

Main X-ray burst 1-8 Å: onset – 02d09^h40^m, max – 02d09^h47^m, $\Phi = 0.0051$ J/m²



CME: 02d10^h50^m, $V = 1393$ km/s, $\Delta\phi = 150^\circ$, $dA = 267^\circ$

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2003 December 02



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2003 December 02

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm².s.sr)⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	12 ^h	18 ^h	42.6	5d	
EPS	>10	12 ^h	18 ^h	21	4.5d	
EPS	>30	12 ^h	16 ^h	0.9	3d	
EPS	>50	10 ^h	16 ^h	0.15	2d	
EPS	>60	10 ^h	16 ^h	0.09	2d	
EPS	>100	-	16 ^h	0.03	-	
METEOR						
CBM	>5	13 ^h	17 ^h	30.4	3d	
CBM	>10	13 ^h	17 ^h	9.9	3d	
CBM	>25	13 ^h	17 ^h	1.56	1.5d	
CBM	>40	13 ^h	16 ^h	0.19	1d	
BP	>90	13 ^h	15 ^h	0.015	0.5d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	18 ^h	3520	5d	
MEPED	>0.8	-	17 ^h	2610	5d	
MEPED	>2.5	-	17 ^h	1420	5d	
MEPED	>6.9	-	17 ^h	630	4.5d	
MEPED	>16	-	17 ^h	30	3d	
MEPED	>36	-	17 ^h	0.6	2d	
MEPED	>70	-	17 ^h	0.1	2d	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	22 ^h	2250	5d	
MKL	>14	-	17 ^h	17	4.5d	
MKL	>26	-	17 ^h	8.4	3d	
MKL	>50	-	17 ^h	1.6	2d	
ACE						
SIS	>10	12 ^h	17 ^h	98.5	4d	
SIS	>30	12 ^h	17 ^h	1.2	3d	
SOHO						
EPHIN (INT)	>50	16 ^h	17 ^h	0.1	1d	

Differential fluxes of protons for the event of 2003 December 02

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	12 ^h	18 ^h	140	4.5d	
LION	2-6	12 ^h	18 ^h	58	4.5d	
EPHIN	4-8	12 ^h	17 ^h	342	4d	
EPHIN	8-25	12 ^h	17 ^h	25.1	4d	
EPHIN	25-41	12 ^h	17 ^h	0.13	4d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2003 December 02

2003 December 02		☉			AR10508			To event 446	
H α	6563 Å	No Flare			s19w90				
1 – 12	keV	0940	0948	0954		C7.2	5.1E-3		
12-25	keV	094524	094602	100316		412771	RHESSI		
9.1	GHz	0942.9	0943.6	0945.0		1.11			
3	GHz	0943.0	0943.9	0944.9		1.72			
245	MHz	0943.0	0943.0	~0943.0		1.93			
DS I	170-270	0943		0944	GG,DC	2			
DS III	180-300	0943		0944	GG	1			
DS DCIM	2098-4500	0936		0944	G	1			
CME	WL	1050	1393 km/s	18.5 km/s ²	150°	267°			

* – probable localization of the flare event

Events in 2004

				Page
1.	Event 2004.04.11 – (2004-102)	№ 447	584
2.	Event 2004.07.22 – (2003-204)	№ 448	589
3.	Event 2004.07.23 – (2003-205)	№ 449	594
4.	Event 2004.07.25 – (2004-207)	№ 450	599
5.	Event 2004.08.01 – (2004-214)	№ 451	604
6.	Event 2004.09.13 – (2004-257)	№ 452	608
7.	Event 2004.09.19 – (2004-263)	№ 453	613
8.	Event 2004.11.01 – (2004-306)	№ 454	618
9.	Event: 2004.11.07 – (2004-312)	№ 455	622
10.	Event 2004.11.10 – (2004-315)	№ 456	627

Particle event: To(Ep>10MeV) – 11d06^h

Tmax₁(Ep>10MeV) – 11d12^h, Jmax₁(Ep>10MeV) – 13 /cm².s.sr

Tmax₂(Ep>10MeV) – 11d20^h, Jmax₂(Ep>10MeV) – 14.5 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – Eqm₁ = 95 MeV

– Eqm₂ = 80 MeV


Sources: • solar flare 11d03^h54^m, C9.6/1F, S14W47, AR10588

Main X-ray burst 1-8 Å: onset – 11d03^h54^m, max – 11d04^h19^m, Φ = 0.013 J/m²



CME: 11d04^h30^m, V = 1645 km/s, Δφ = 314°, dA = 237°

Ø CME 11d11^h54^m, V = 1132 km/s, Δφ = 360°, dA = 295° - back side event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2004 April 11



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 April 11

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	06 ^h	19 ^h /20 ^h	47/56.5	2d	
EPS	>10	06 ^h	12 ^h /20 ^h	13/14.5	2d	
EPS	>30	06 ^h	12 ^h /20 ^h	0.6/0.5	2d	
EPS	>50	06 ^h	11 ^h /20 ^h	0.16/0.1	1d	
EPS	>60	06 ^h	11 ^h / -	0.08/ -	1d	
EPS	>100	06 ^h	10 ^h / -	0.02/ -	1d	
METEOR						
CBM	>5	05 ^h	15 ^h / -	6.65/ -	3d	
CBM	>15	05 ^h	13 ^h / -	2.2./ -	2d	
CBM	>25	06 ^h	13 ^h / -	0.65/ -	2d	
CBM	>40	06 ^h	13 ^h / -	0.2/ -	1d	
BP	>90	06 ^h	13 ^h / -	0.014/ -	1d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	06 ^h	12 ^h /20 ^h	130/880	3d	
MEPED	>0.8	06 ^h	12 ^h /20 ^h	110/770	3d	
MEPED	>2.5	06 ^h	12 ^h /20 ^h	90/550	2d	
MEPED	>6.9	06 ^h	12 ^h /20 ^h	80/150	2d	
MEPED	>16	06 ^h	12 ^h /20 ^h	9/6	1d	
MEPED	>36	06 ^h	12 ^h /20 ^h	1.4/1	1d	
MEPED	>70	-	11 ^h / -	0.3/ -	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	13 ^h / -	180/ -	3d	
MKL	>14	-	13 ^h / -	3.7/ -	2d	
MKL	>26	-	13 ^h / -	2.5/ -	2d	
MKL	>50	-	13 ^h / -	0.9/ -	1d	
ACE						
SIS	>10	05 ^h	10 ^h /20 ^h	14.3/28.2	2d	
SIS	>30	05 ^h	10 ^h /20 ^h	1.7/0.5	1d	
SOHO						
EPHIN (INT)	>50	05 ^h	11 ^h / -	0.07/ -	2d	

Differential fluxes of protons for the event of 2004 April 11

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	06 ^h	12 ^h /22 ^h	8.9/59.2	5d	
LION	2-6	06 ^h	12 ^h /22 ^h	3.3/19.2	5d	
EPHIN	4-8	05 ^h	11 ^h /21 ^h	13.8/59.1	5d	
EPHIN	8-25	05 ^h	11 ^h /20 ^h	1.1/2.9	5d	
EPHIN	25-41	05 ^h	11 ^h /19 ^h	0.047/0.017	4d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 April 11

2004 April 11


•

AR10588

To event 447

Ha	6562 Å	0400	0421	0451	S14W47	1F	EF
DSF	Ha	~0307		0400	S18W50	16°	
1 – 12	keV	0354	0419	0435		C9.6	1.3E-2
50-100	keV	040632	041542	044344		937728	RHESSI
50-150	keV	0413		0418			SONG F
150-500	keV						SONG F
0.18-0.7	MeV						SONG F

15.4	GHz	0412.0	0415.0	0428.0		2.34	
8.8	GHz	0411.0	0415.0	0000.0		2.66	
5	GHz	0411.0	0416.0	0425.0		2.81	
2.7	GHz	0410.0	0415.0	0424.0	[U0.6] P2.7	2.99	
1.4	GHz	0407.0	0407.0	0000.0		2.38	
610	MHz	0400.0	0402.0	0404.0		2.20	
410	MHz	0411.0	0413.0	0415.0		2.76	
245	MHz	0408.0	0410.0	0411.0		2.15	
DS III	18-700	0358		0425	GG	3	
CME	WL	0430	1645 km/s	-77.6 km/s ²	314°	237°	



Particle event: To(Ep>10MeV) – 22d17^h

Tmax₁(Ep>10MeV) – 22d20^h, Jmax₁(Ep>10MeV) – 0.9 /cm².s.sr

Tmax₂(Ep>10MeV) – 23d10^h, Jmax₂(Ep>10MeV) – 2 /cm².s.sr

Duration of the event – 1 day

Quasimaximal energy of protons in the event – Eqm₁ = 70 MeV

– Eqm₂ = 155 MeV

Sources: ☉ solar flare 22d00^h14^m, M9.1/SB, n06e25*, AR10652


Main X-ray burst 1-8 Å: onset – 22d00^h14^m, max – 22d00^h32^m, Φ = 0.079 J/m²

CME: 22d01^h32^m, V = 0492 km/s, Δφ = 083°, dA = 180°



▲ SC 24d06^h13^m

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2004 July 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 July 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	17 ^h	20 ^h /23d10 ^h	3/7.1	1d	
EPS	>10	17 ^h	20 ^h /23d10 ^h	0.9/2	1d	
EPS	>30	17 ^h	19 ^h /23d10 ^h	0.07/0.42	1d	
EPS	>50	17 ^h	- /23d10 ^h	- /0.15	1d	
EPS	>60	17 ^h	- /23d10 ^h	- /0.1	1d	
EPS	>100	-	- /23d10 ^h	- /0.04	-	
METEOR						
CBM	>5	11 ^h	20 ^h /23d11 ^h	8/4.3	1d	
CBM	>15	11 ^h	20 ^h /23d09 ^h	2/1.6	1d	
CBM	>25	22 ^h	- /23d09 ^h	- /0.23	1d	
CBM	>40	22 ^h	- /23d09 ^h	- /0.12	1d	
BP	>90	22 ^h	- /23d09 ^h	- /0.01	1d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	12 ^h	- /23d10 ^h	- /420	1d	
MEPED	>0.8	12 ^h	- /23d10 ^h	- /190	1d	
MEPED	>2.5	12 ^h	- /23d10 ^h	- /90	1d	
MEPED	>6.9	-	- /23d10 ^h	- /20	1d	
MEPED	>16	-	- /23d10 ^h	- /1.4	1d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	20 ^h /23d10 ^h	95/560	1d	
MKL	>14	-	20 ^h /23d10 ^h	0.35/1.4	1d	
MKL	>26	-	20 ^h /23d10 ^h	0.14/0.9	1d	
MKL	>50	-	20 ^h /23d10 ^h	0.07/0.45	1d	
ACE						
SIS	>10	17 ^h	19 ^h /23d08 ^h	0.9/3	1d	
SIS	>30	17 ^h	20 ^h /23d08 ^h	0.2/0.9	1d	
SOHO						
EPHIN (INT)	>50	-	- /23d10 ^h	- /0.05	1d	

Differential fluxes of protons for the event of 2004 July 22


S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	11 ^h	20 ^h /23d08 ^h	1.6/9.7	1d	
LION	2-6	11 ^h	20 ^h /23d08 ^h	0.26/1.2	1d	
EPHIN	4-8	15 ^h	19 ^h /23d08 ^h	1.26/2.1	1d	
EPHIN	8-25	15 ^h	19 ^h /23d08 ^h	0.096/0.14	1d	
EPHIN	25-41	15 ^h	19 ^h /23d08 ^h	0.003/0.009	1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 July 22

2004 July 22 ☉ AR10652 To event 448

H α	6563 Å	No Flare Patrol			n06e25		
1 – 12	keV	0014	0032	0043		M9.1	7.9E-2
25-50	keV	235256	002958	005528		46462728	RHESSI

15.4	GHz	0017.0	0031.0	0126.0		2.51	
8.8	GHz	0018.0	0031.0	0124.0		2.56	
5	GHz	0015.0	0027.0	0109.0	U0.6 P5	2.60	
2.7	GHz	0021.0	0027.0	0109.0		2.56	
1.4	GHz	0024.0	0026.0	0031.0		2.32	
610	MHz	0032.0	0032.0	0123.0		1.91	
410	MHz	0015.0	0015.0	0016.0		2.77	
245	MHz	0030.0	0031.0	0031.0		3.46	
DS III	18-200	<0000		>0710	S,C	3	
DS III	100-1000	0015		0131	N	1	
CME	WL	0132	0492 km/s	0.3 km/s ²	083°	180°	



Particle event: To($E_p > 10$ MeV) – 23d16^h

Tmax($E_p > 10$ MeV) – 23d19^h, Jmax($E_p > 10$ MeV) – 1.8 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm} = 75$ MeV

Sources: ☉ solar flare 22d22^h23^m, 2N/M1.6, N05E04 AR10652*

22d23^h10^m, 2N/M1.2, N05E04 AR10652*

☾ solar flare 23d17^h07^m, M2.2/SF, N03W04 AR10652

Main X-ray burst 1–8 Å: onset–22d22^h40^m, max–22d22^h58^m, $\Phi = 0.016$ Jo.m⁻²


onset–22d23^h10^m, max–22d23^h24^m, $\Phi = 0.023$ Jo.m⁻²

CME: 22d23^h54^m; $V = 0448$ km/s; $\Delta\varphi = 046^\circ$; $dA = 200^\circ$



▲ SC 24d06^h13^m

* – One solar flare event with two X-ray burst

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2004 July 23



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 July 23

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	16 ^h	19 ^h	5.8	2d	
EPS	>10	16 ^h	19 ^h	1.8	2d	
EPS	>30	16 ^h	19 ^h	0.27	2d	
EPS	>50	16 ^h	19 ^h	0.07	2d	
EPS	>60	16 ^h	19 ^h	0.05	2d	
EPS	>100	-	-	-	-	
METEOR						
CBM	>5	16 ^h	24d01 ^h	2,2	2d	
CBM	>15	16 ^h	24d01 ^h	1	2d	
CBM	>25	16 ^h	24d01 ^h	0.36	1d	
CBM	>40	16 ^h	24d01 ^h	0.13	1d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	20 ^h	2300	2d	
MEPED	>0.8	-	20 ^h	800	2d	
MEPED	>2.5	-	20 ^h	120	2d	
MEPED	>6.9	-	20 ^h	24	2d	
MEPED	>16	-	20 ^h	3	2d	
MEPED	>36	-	20 ^h	1.4	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	20 ^h	540	2d	
MKL	>14	-	20 ^h	1.4	2d	
MKL	>26	-	20 ^h	0.9	1d	
MKL	>50	-	20 ^h	0.4	1d	
ACE						
SIS	>10	16 ^h	19 ^h	4.2	1d	
SIS	>30	16 ^h	19 ^h	0.8	1d	
SOHO						
EPHIN (INT)	>50	14 ^h	18 ^h	0.05	1d	

Differential fluxes of protons for the event of 2004 July 23

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	14 ^h	19 ^h	40	2d	
LION	2-6	14 ^h	19 ^h	4.6	2d	
EPHIN	4-8	14 ^h	18 ^h	5.3	2d	
EPHIN	8-25	14 ^h	18 ^h	0.27	2d	
EPHIN	25-41	14 ^h	18 ^h	0.013	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Kuwabara T., Bieber J.W., Clem J., et.al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 July 23


2004	July 22	☉		AR10652	To event 449		
Hα	6563 Å	2223	2256	>2333	N05E04	2N*	MZ
1 – 12	keV	2240	2258	2307		M1.6	1.6E-2
1 – 12	keV	2310	2324	2343		M1.2	2.3E-2
25-50	keV	223740	225450	230600		5664912	RHESSI
25-50	keV	230600	231514	231916		3941621	
15.4	GHz	2246.0	2255.0	2317.0	U2.7 / 15	2.70	
8.8	GHz	2244.0	2255.0	2327.0		2.58	
5	GHz	2243.0	2255.0	2327.0		2.11	
2.7	GHz	2244.0	2247.0	2255.0		1.83	
610	MHz	2253.0	2254.0	2301.0		2.51	
410	MHz	2245.0	2255.0	2321.0		2.52	
245	MHz	2243.0	2302.0	2328.0		2.84	
DS III	18-880	2253		2302	GG	3	
DS CONT	25-180	2338		0919		2	
CME	WL	2354	0448 km/s	-25.8km/s ²	046°	200°	

* – One solar flare event with two X-ray burst



2004 July 23 Ø AR10652 To event 449

H α	6563 Å	<1723	~1723	>1735	N03W04	SF	F
1 – 12	keV	1707	1728	1735		M2.2	1.5E-2
12-25	keV	172652	172726	175300		2487273	RHESSI
15.4	GHz	1716.0	1720.0	0000.0		2.68	
8.8	GHz	1719.0	1720.0	0000.0	[U0.6-2.7] P8.8	3.15	
5	GHz	1717.0	1720.0	1733.0		2.98	
2.7	GHz	1718.0	1720.0	1733.0		2.32	
1.4	GHz	1719.0	1720.0	1729.0		2.46	
610	MHz	1718.0	1719.0	1721.0		2.11	
410	MHz	1718.0	1728.0	1728.0		2.70	
245	MHz	1716.0	1720.0	0000.0		3.60	
DS III	25-180	1710		1711		2	
DS III	25-180	1716		1732	N	2	
DS DCIM	100-4000	1710		1737	P,C	3	
CME	WL	1754	0569 km/s	-8.1 km/s ²	142°	256°	



Particle event: To($E_p > 10$ MeV) – 25d17^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 25\text{d}21^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 27 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 26\text{d}23^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 430 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 4 days

Quasimaximal energy of protons in the event – $E_{\text{qm}1} = 140 \text{ MeV}$

– $E_{\text{qm}2} = 155 \text{ MeV}$

Sources: ● solar flare 25d14^h19^m, M1.1/1F, N08W33, AR10652


○ solar flare 26d17^h23^m, M1.1/2N, N03W45, AR10652

Main X-ray burst 1-8 Å: onset – 25d14^h19^m, max – 25d15^h14^m, $\Phi = 0.065 \text{ J/m}^2$



CME: 25d14^h54^m, $V = 1333 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 204^\circ$

▲ SC 26d22^h49^m;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2004 July 25



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 July 25

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	17 ^h	21 ^h /26d23 ^h	60/2280	4d	
EPS	>10	17 ^h	21 ^h /26d23 ^h	27/430	4d	
EPS	>30	17 ^h	20 ^h /26d22 ^h	1.6/8.7	3d	
EPS	>50	17 ^h	20 ^h /26d22 ^h	0.2/0.9	2d	
EPS	>60	17 ^h	19 ^h /26d22 ^h	0.12/0.42	2d	
EPS	>100	17 ^h	19 ^h /26d21 ^h	0.02/0.01	2d	
METEOR						
CBM	>5	17 ^h	21 ^h /26d23 ^h	22/167	4d	
CBM	>15	17 ^h	21 ^h /26d23 ^h	7.4/45	3.5d	
CBM	>25	17 ^h	21 ^h /26d23 ^h	1.6/9.2	2d	
CBM	>40	17 ^h	21 ^h /26d23 ^h	0.2/1.1	1.5d	
BP	>90	17 ^h	22 ^h /26d23 ^h	0.002/0.007	1.5d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	- /26d23 ^h	- /1.9·10 ⁴	4d	
MEPED	>0.8	-	21 ^h /26d23 ^h	920/1.1·10 ⁴	4d	
MEPED	>2.5	-	21 ^h /26d23 ^h	350/7·10 ³	4d	
MEPED	>6.9	-	21 ^h /26d23 ^h	130/2.8·10 ³	4d	
MEPED	>16	-	21 ^h /26d23 ^h	26/210	3d	
MEPED	>36	-	21 ^h /26d23 ^h	2.5/7	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	20 ^h /26d24 ^h	540/1.5·10 ⁴	4d	
MKL	>14	-	20 ^h /26d23 ^h	9.8//130	4d	
MKL	>26	-	20 ^h /26d22 ^h	4.5/40	4d	
MKL	>50	-	20 ^h /26d22 ^h	1.75/10	4d	
ACE						
SIS	>10	17 ^h	19 ^h /26d22 ^h	40.5/765	3d	
SIS	>30	17 ^h	19 ^h /26d21 ^h	2.8/18	2d	
SOHO						
EPHIN (INT)	>50	16 ^h	19 ^h /26d22 ^h	0.06/0.8	2d	

Differential fluxes of protons for the event of 2004 July 25

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	17 ^h	22 ^h /26d23 ^h	5.5/875	3d	
LION	2-6	17 ^h	20 ^h /26d23 ^h	0.9/240	3d	
EPHIN	4-8	17 ^h	19 ^h /26d22 ^h	6.6/1050	3d	
EPHIN	8-25	16 ^h	19 ^h /26d21 ^h	1.5/66	3d	
EPHIN	25-41	16 ^h	19 ^h /26d21 ^h	0.075/0.65	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	



References:

Kuwabara T., Bieber J.W., Clem J., et.al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 July 25

2004	July 25	•	AR10652	To event 450			
Hα	6563 Å	1433	1448	1643	N08W33	1F	F
1 – 12	keV	1419	1514	1643		M1.1	6.5E-02
6-12	keV	145220	145834	150504		646080	RHESSI
6-12	keV	150504	150750	151800		589305	RHESSI
6-12	keV	163112	163138	165420		146113	RHESSI
15.4	GHz	1447.0	1521.0	1603.0		1.70	
8.8	GHz	1441.0	1519.0	1647.0		1.94	
5	GHz	1441.0	1543.0	1634.0		2.04	
2.7	GHz	1424.0	1506.0	1603.0		2.08	
1.4	GHz	1434.0	1509.0	1603.0		2.11	
610	MHz	1424.0	1521.0	1606.0		2.78	
410	MHz	1424.0	1505.0	1551.0		2.69	
245	MHz	1424.0	1520.0	1534.0		3.32	
DS II	U2.4 \ P15	1521		1526		1	
DS IV	25-81	1415		1731		2	
DS DCIM	25-180	1421		1518	GG	2	
DS DCIM	2000-4500	1434		1543	GG	2	
DS DCIM	800-2000	1503		1530	C	3	

8.8	GHz	1641.0	1641.0	~1641.0		1.89	
5	GHz	1637.0	1641.0	1646.0		2.18	
2.7	GHz	1638.0	1639.0	1646.0	P2.7	2.23	
1.4	GHz	1641.0	1642.0	1643.0		1.96	
CME	WL	1454	1333 km/s	7.0 km/s ²	360°	204°	



2004 July 26 Ø AR10652 To event 450

H α	6563 Å	1726	1736	1752	N03W45	2N	EF
DSF	H α	~1652		~1407	N16W50	12°	
1 – 12	keV	1723	1730	1737		M1.1	6.3E-3
25-50	keV	172828	173046	173316		329366	RHESSI
15.4	GHz	1730.0	1730.0	~1730.0		1.86	
8.8	GHz	1726.0	1730.0	1731.0	U1.4 U8.8	2.28	
5	GHz	1726.0	1728.0	1731.0		2.08	
2.7	GHz	1727.0	1728.0	1729.0		1.58	
1.4	GHz	1726.0	1730.0	1730.0		1.41	
610	MHz	1725.0	1725.0	1729.0		1.60	
410	MHz	1725.0	1725.0	1729.0		2.04	
DS III	250-4000	1723		1730	GG,RS,C	3	
DS DCIM	800-2000	1725		1730	GG	1	
CME	WL	1830	0401 km/s	39.5 km/s ²	058°	221°	

Particle event: To($E_p > 10$ MeV) – 01d01^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 01\text{d}21^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 5.2 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 02\text{d}02^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 4.8 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm1} = 80$ MeV

– $E_{qm2} = 75$ MeV

Sources: \diamond flare activity AR10652 $> 1.5d$ behind W-limb

\square solar flare 31d05^h16^m, C8.4/..., n02w90*, AR10652


\emptyset solar flare 31d10^h35^m, C5.3/..., n02w90*, AR10652

Main X-ray burst 1-8 Å: onset – 31d05^h16^m, max – 31d06^h57^m, $\Phi = 0.076 \text{ J/m}^2$



CME: 05^h54^m: $V = 1192 \text{ km/s}$, $\Delta\phi = 259^\circ$, $dA = 287^\circ$

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2004 August 01



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 August 01

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	21 ^h	21 ^h /02d02 ^h	15.2/18.3	2d	
EPS	>10	21 ^h	21 ^h /02d02 ^h	5.2/4.8	2d	
EPS	>30	-	22 ^h /02d01 ^h	0.43/0.15	2d	
EPS	>50	-	- /02d01 ^h	- /0.04	2d	
EPS	>60	-	-	-	-	
EPS	>100	-	-	-	-	
METEOR						
CBM	>5	02 ^h	22 ^h / -	3/ -	2d	
CBM	>15	02 ^h	23 ^h / -	1.2/ -	1.5d	
CBM	>25	00 ^h	22 ^h / -	0.3/ -	1d	
CBM	>40	00 ^h	23 ^h / -	0.1/ -	1d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	22 ^h /02d03 ^h	620/450	2d	
MEPED	>0.8	-	22 ^h /02d03 ^h	380/220	2d	
MEPED	>2.5	-	22 ^h /02d03 ^h	210/170	2d	
MEPED	>6.9	-	22 ^h /02d03 ^h	45/30	2d	
MEPED	>16	-	22 ^h /02d03 ^h	4.8/1.8	2d	
MEPED	>36	-	22 ^h /02d03 ^h	0.6/0.3	2d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	22 ^h /02d03 ^h	430/110	2d	
MKL	>14	-	22 ^h /02d03 ^h	2.6/1.5	2d	
MKL	>26	-	22 ^h /02d03 ^h	1.5/0.9	2d	
MKL	>50	-	22 ^h /02d03 ^h	0.5/0.3	2d	
ACE						
SIS	>10	04 ^h	22 ^h /02d01 ^h	11.3/5.7	2d	
SIS	>30	09 ^h	22 ^h /02d01 ^h	0.6/0.3	2d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 2004 August 01

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.72-2	31d17 ^h	20 ^h /02d02 ^h	18.4/7.9	2d	
LION	2-6	31d17 ^h	20 ^h /02d02 ^h	3/1.3	2d	
EPHIN	4-8	04 ^h	19 ^h /02d01 ^h	6.8/3.5	2d	
EPHIN	8-25	04 ^h	19 ^h /02d01 ^h	0.45/0.2	2d	
EPHIN	25-41	04 ^h	19 ^h /02d01 ^h	0.014/0.006	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Kuwabara T., Bieber J.W., Clem J., et.al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 August 01

2004 July 31		☐		AR10652		To event 451	
H α	6563 Å	No Flare			n02w90*		
1 – 12	keV	0516	0657	0914		C8.4	7.6E-2
6-12	keV	063024	063458	063916		362241	RHESSI
12-25	keV	063916	064334	071552		1405027	RHESSI
12-25	keV	064900	065014	071600		978017	RHESSI
6-12	keV	084648	084658	085232		44076	RHESSI
CME	WL	0554	1192 km/s	46.4 km/s ²	197°	287°	

* – probable localization of the flare event

2004 July 31		Ø		AR10652		To event 451	
H α	6563 Å	No Flare			n02w90*		
1 – 12	keV	1035	1101	1149		C5.3	2.1E-2
6-12	keV	11:45:44	11:45:46	12:00:44		57313	RHESSI

* – probable localization of the flare event

Particle event: To($E_p > 10$ MeV) – 13d19^h

$T_{\max 1}(E_p > 10 \text{ MeV})$ – 13d23^h, $J_{\max 1}(E_p > 10 \text{ MeV})$ – 210 /cm².s.sr

$T_{\max 2}(E_p > 10 \text{ MeV})$ – 14d05^h, $J_{\max 2}(E_p > 10 \text{ MeV})$ – 180 /cm².s.sr

Duration of the event – 4 days

Quasimaximal energy of protons in the event – $E_{qm1} = 110$ MeV

– $E_{qm2} = 90$ MeV

Sources: ☉ solar flare 12d00^h04^m, M4.8/2N, N04E42, AR10672


☾ solar flare 12d01^h36^m, M3.2/SN, S14W61, AR10667

Main X-ray burst 1-8 Å onset – 12d00^h04^m, max – 12d00^h56^m, $\Phi = 0.15$ J/m²



CME: 12d00^h36^m, $V = 1328$ km/s, $\Delta\phi = 360^\circ$, $dA = 132^\circ$

▲ SC 13d20^h03^m;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2004 September 13



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 September 13

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	19 ^h	23 ^h /14d06 ^h	970/990	5d	
EPS	>10	19 ^h	23 ^h /14d05 ^h	210/180	4d	
EPS	>30	19 ^h	22 ^h /14d06 ^h	5.6/1.2	3d	
EPS	>50	19 ^h	22 ^h /14d05 ^h	0.5/0.15	1d	
EPS	>60	19 ^h	23 ^h /14d05 ^h	0.2/0.1	1d	
EPS	>100	-	23 ^h /14d05 ^h	0.02/0.015	-	
METEOR						
CBM	>5	18 ^h	14d01 ^h / -	70/ -	3d	
CBM	>15	18 ^h	23 ^h / -	18/ -	2d	
CBM	>25	18 ^h	23 ^h / -	3.2/ -	1d	
CBM	>40	22 ^h	22 ^h / -	0.46/ -	1d	
BP	>90	-	-	-	-	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	24 ^h /14d08 ^h	1·10 ⁵ /8.4·10 ³	-	
MEPED	>0.8	-	24 ^h /14d08 ^h	6.2·10 ³ /6.1·10 ³	-	
MEPED	>2.5	-	24 ^h /14d08 ^h	3·10 ³ /3·10 ³	-	
MEPED	>6.9	-	24 ^h /14d08 ^h	620/610	-	
MEPED	>16	-	24 ^h /14d08 ^h	70/25	-	
MEPED	>36	-	24 ^h /14d07 ^h	2.5/0.5	-	
MEPED	>70	-	24 ^h / -	0.3/ -	-	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	22 ^h /14d05 ^h	600/1·10 ⁴	-	
MKL	>14	-	22 ^h /14d05 ^h	5/18	-	
MKL	>26	-	22 ^h /14d05 ^h	2.1/11	-	
MKL	>50	-	22 ^h /14d05 ^h	0.9/3.3	-	
ACE						
SIS	>10	19 ^h	23 ^h /14d05 ^h	210/180	4d	
SIS	>30	19 ^h	22 ^h /14d06 ^h	5.6/1.2	3d	
SOHO						
EPHIN (INT)	>50	20 ^h	21 ^h /14d05 ^h	0.43/0.32	1d	

Differential fluxes of protons for the event of 2004 September 13

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.72-2	19 ^h	14d01 ^h /14d06 ^h	390/447	4d	
LION	2-6	19 ^h	14d00 ^h /14d06 ^h	75/98	4d	
EPHIN	4-8	19 ^h	23 ^h /14d05 ^h	376/443	4d	
EPHIN	8-25	19 ^h	23 ^h /14d05 ^h	25.3/24.4	4d	
EPHIN	25-41	19 ^h	22 ^h /14d04 ^h	0.3/0.12	4d	


References:

Kuwabara T., Bieber J.W., Clem J., et.al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 September 13

2004	September 12	☉	AR10672	To event 452			
Hα	6563 Å	0008	0035	0201	N04E42	2N	FHUZ
1 – 12	keV	0004	0056	0133		M4.8	1.5E-1
50-100	keV	000216	003934	010448		12612176	RHESSI
15.4	GHz	0032.0	0119.0	0000.0		2.83	
8.8	GHz	0029.0	0121.0	0256.0		3.23	
5	GHz	0023.0	0119.0	0246.0	U0.6 P2.7-5	3.45	
2.7	GHz	0026.0	0121.0	0246.0		3.45	
1.4	GHz	0024.0	0116.0	0000.0		3.28	
610	MHz	0026.0	0033.0	0205.0		2.68	
410	MHz	0028.0	0137.0	0200.0		3.15	
245	MHz	0135.0	0137.0	0140.0		2.93	

DS II	30-57	0023		0029	F,N	3	
DS II	60-110	0023		0029	S,H	3	
DS II	20-90	0029		0049	S,H	3	
DS II	30-90	0141		0150	F,N	3	
DS II	57-180	0141		0150	S,H	3	
DS IV	30-850	0013		0130		2	
DS IV	28-180	0026		0200		2	
DS III	30-130	0023		0038	GG	2	
DS III	20-180	0041		0111	GG	2	
DS III	75-440	0135		0140	G	1	
DS III	20-90	0149		0150	G	3	
DS UNCLF	50-85	0043		0047		2	
CME	WL	0036	1328 km/s	22.5 km/s ²	360°	132°	



2004 September 12		Ø		AR10667	To event 452		
Hα	6563 Å	0137	0138	0145	S14W61	SN	
1 – 12	keV	0136	0139	0141		M3.2	8.2E-3
12-25	keV	013740	013858	014008		343715	RHESSI
410	MHz	0028.0	0137.0	0200.0		3.15	
245	MHz	0135.0	0137.0	0140.0		2.93	
DS III	75-440	0135		0140	G	1	
DS III	20-90	0149		0150	G	3	

Particle event: To($E_p > 10$ MeV) – 19d18^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 20\text{d}01^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 46 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 21\text{d}02^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 10 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm1} = 390$ MeV


– $E_{qm2} = 100$ MeV

Sources: • solar flare 19d16^h46^m, M1.9/..., n03w60* AR10672



Main X-ray burst 1–8 Å: onset – 19d16^h46^m, max – 19d17^h12^m, $\Phi = 0.039 \text{ J/m}^2$

CME: gap

* – probable localization of the flare event




Time profiles of the proton fluxes for the event of 2004 September 19



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 September 19

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura-tion	Comments
GOES-10						
EPS	>5	18 ^h	20d01 ^h /20d11 ^h	98/28	2d	
EPS	>10	18 ^h	20d01 ^h /20d08 ^h	46/10	2d	
EPS	>30	18 ^h	21 ^h /20d08 ^h	7.7/1	2d	
EPS	>50	18 ^h	20 ^h /20d08 ^h	2.1/0.3	1d	
EPS	>60	18 ^h	20 ^h /20d07 ^h	1.0/0.13	1d	
EPS	>100	18 ^h	20 ^h /20d08 ^h	0.2/0.01	1d	
METEOR						
CBM	>5	17 ^h	20 ^h / -	35/ -	3d	
CBM	>15	17 ^h	20 ^h / -	14/ -	2d	
CBM	>25	17 ^h	20 ^h / -	4.9/ -	2d	
CBM	>40	17 ^h	20 ^h / -	1.7/ -	1d	
BP	>90	17 ^h	20 ^h / -	0.13/ -	1d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	22 ^h / -	140/ -	2d	
MEPED	>16	-	22 ^h / -	30/ -	2d	
MEPED	>36	-	20 ^h / -	7.2/ -	1d	
MEPED	>70	-	20 ^h / -	0.65/ -	1d	
MEPED	>140	-	20 ^h / -	0.28/ -	1d	

CORONAS F						
MKL	>1.	-	21 ^h /20d08 ^h	55/110	2d	
MKL	>14	-	21 ^h /20d08 ^h	17/12	1d	
MKL	>26	-	21 ^h /20d08 ^h	8/6.8	1d	
MKL	>50	-	21 ^h /20d08 ^h	3.5/3	1d	
ACE						
SIS	>10	17 ^h	23 ^h / -	55.2/ -	2d	
SIS	>30	17 ^h	20 ^h / -	10.3/ -	1d	
SOHO						
EPHIN (INT)	>50	17 ^h	- /20d11 ^h	-	1d	

Differential fluxes of protons for the event of 2004 September 19

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	17 ^h	- /20d17 ^h	- /16.5	3d	
LION	2-6	17 ^h	- /20d17 ^h	- /2.2	3d	
EPHIN	4-8	17 ^h	- /20d10 ^h	- /5.9	3d	
EPHIN	8-25	17 ^h	- /20d10 ^h	- /0.6	3d	
EPHIN	25-41	17 ^h	-	-	-	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 September 19

2004 September 19

☉


AR1010672

To event 453

H α	6563 Å	No Flare Patrol			n03w60*		
1 – 12	keV	1646	1712	1739		M1.9	3.9E-2
50-100	keV	16:43:16	17:01:22	17:02:52		71699552	RHESSI

15.4	GHz	1656.0	1658.0	1713.0		1.94	
8.8	GHz	1655.0	1657.0	0000.0		2.23	
5	GHz	1651.0	1657.0	0000.0	U0.6 P2.7-5	2.69	
2.7	GHz	1649.0	1657.0	0000.0		2.72	
1.4	GHz	1650.0	1654.0	0000.0		2.00	
610	MHz	1652.0	1655.0	1655.0		1.49	
410	MHz	1655.0	1656.0	1656.0		1.79	
245	MHz	1701.0	1704.0	1706.0		2.15	
DS II	25-180	1656		1713		3	
DS IV	25-76	1716		1734		1	
DS III	42-180	1650		1653		1	
CME	WL						gap

* – probable localization of the flare event



Particle event: To($E_p > 10$ MeV) – 01d06^h

Tmax ($E_p > 10$ MeV) – 01d08^h, Jmax ($E_p > 10$ MeV) – 54 /cm².s.sr

Duration of the event – 2 days


Quasimaximal energy of protons in the event – $E_{qm} = 410$ MeV

Sources: • solar flare 01d03^h04^m, M1.1/1F, N15W41, AR10691

Main X-ray burst 1-8 Å: onset – 01d03^h04^m, max – 01d03^h22^m, $\Phi = 0.0052$ J/m²



CME: 01d03^h54^m, V = 0459 km/s, $\Delta\phi = 192^\circ$, dA = 285°

Particle fluxes and associated phenomena




October - November 2004

Time profiles of the proton fluxes for the event of 2004 November 01



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 November 01

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES 10						
EPS	>5	06 ^h	08 ^h	76	2d	
EPS	>10	06 ^h	08 ^h	54	2d	
EPS	>30	06 ^h	07 ^h	15.5	2d	
EPS	>50	06 ^h	07 ^h	6.4	2d	
EPS	>60	06 ^h	07 ^h	3.9	1d	
EPS	>100	06 ^h	07 ^h	1.1	1d	
METEOR						
CBM	>5	06 ^h	08 ^h	34	3d	
CBM	>15	06 ^h	08 ^h	19	3d	
CBM	>25	06 ^h	07 ^h	11	2d	
CBM	>40	06 ^h	07 ^h	6.7	2d	
BP	>90	06 ^h	07 ^h	0.9	1d	
POES 16						
MEPED	>0.8	06 ^h	-	-	2d	
MEPED	>2.5	06 ^h	-	-	2d	
MEPED	>6.9	06 ^h	07 ^h	120	2d	
MEPED	>16	06 ^h	07 ^h	70	2d	
MEPED	>36	06 ^h	07 ^h	22	2d	
MEPED	>70	06 ^h	07 ^h	3.3	1d	
MEPED	>140	06 ^h	07 ^h	0.6	1d	
CORONAS F						
MKL	>1.	06 ^h	08 ^h	220	2d	
MKL	>14	06 ^h	08 ^h	38	2d	
MKL	>26	06 ^h	08 ^h	25	2d	
MKL	>50	06 ^h	08 ^h	12	1d	
ACE						
SIS	>10	06 ^h	07 ^h	114	2d	
SIS	>30	06 ^h	07 ^h	37.6	2d	

SOHO						
EPHIN (INT)	>50	06 ^h	07 ^h	2.5	1d	

Differential fluxes of protons for the event of 2004 November 01

S/c, instruments	ΔE , MeV	To	Tmax	J_{\max} , (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	07 ^h	12 ^h	4.3	3d	
LION	2-6	07 ^h	10 ^h	1.7	3d	
EPHIN	4-8	06 ^h	08 ^h	9.7	2d	
EPHIN	8-25	06 ^h	08 ^h	4.3	2d	
EPHIN	25-41	06 ^h	08 ^h	0.43	2d	


References:

Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 November 01

2004 November 01 • AR10691 To event 454

H α	6563 Å	0319	0320	0336	N15W41	1F	
1 – 12	keV	0304	0322	0326		M1.1	5.2E-3
50-100	keV	031724	032054	034328		2310280	RHESSI
15.4	GHz	0319.0	0319.0	~0319.0		2.04	
8.8	GHz	0319.0	0319.0	~0319.0	U2.7 P8.8	2.30	
5	GHz	0318.0	0319.0	0319.0		2.23	
2.7	GHz	0319.0	0319.0	~0319.0		1.95	
1.4	GHz	0319.0	0319.0	~0319.0		2.04	
610	MHz	0318.0	0319.0	0320.0		2.36	
500	MHz	0318.0	0319.0	0321.0		2.04	
245	MHz	<0319.0	~0322.0	>0326.0		2.87	
DS II	30-220	0321		0331		3	
DS III	23-260	0308		0309	G	1	
DS III	25-180	0312		0320		2	
DS III	25-1200	0317		0320	G	3	
CME	WL	0354	0459 km/s	-6.6 km/s ²	192°	285°	



Particle event: To($E_p > 10$ MeV) – 07d01^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 07\text{d}23^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 490 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 09\text{d}00^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 70 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm1} = 330 \text{ MeV}$

– $E_{qm2} = 100 \text{ MeV}$

Sources: • solar flare 07d15^h42^m, X2.0/..., n10w15*, AR10696

Ø solar flare 08d15^h43^m, M2.3/1N, N08W35, AR10696


Main X-ray burst 1–8 Å: onset – 07d15^h42^m, max – 07d16^h06^m, $\Phi = 0.2 \text{ J/m}^2$

CME: 07d16^h54^m, $V = 1759 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 000^\circ$



▲ SC 07d02^h57^m; ▲ 07d10^h52^m; ▲ 07d18^h27^m; ▲ 09d09^h30^m;

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2004 November 07



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 November 07

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	01 ^h	23 ^h /09d00 ^h	1.5·10 ³ /320	2d	
EPS	>10	01 ^h	23 ^h /09d00 ^h	490/70	2d	
EPS	>30	01 ^h	23 ^h /09d00 ^h	23/3.1	2d	
EPS	>50	01 ^h	23 ^h /09d00 ^h	3.7/0.37	2d	
EPS	>60	01 ^h	22 ^h /09d00 ^h	1.9/0.13	2d	
EPS	>100	-	22 ^h /08d23 ^h	0.35/0.01	2d	
METEOR						
CBM	>5	18 ^h	08d03 ^h /08d22 ^h	107/16	2d	
CBM	>15	18 ^h	08d03 ^h /08d22 ^h	35/4.4	2d	
CBM	>25	18 ^h	23 ^h /08d23 ^h	10/0.95	2d	
CBM	>40	18 ^h	23 ^h /08d23 ^h	2.7/0.24	2d	
BP	>90	18 ^h	08d00 ^h / -	0.22/ -	2d	
ChD	>600	-				
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	- /08d23 ^h	- /2.2·10 ³	2d	
MEPED	>2.5	-	24 ^h /08d23 ^h	3.3·10 ³ /1·10 ³	2d	
MEPED	>6.9	-	24 ^h /08d23 ^h	920/180	2d	
MEPED	>16	-	22 ^h /08d23 ^h	110/11	2d	
MEPED	>36	-	22 ^h /08d23 ^h	10.5/1.3	2d	
MEPED	>70	-	22 ^h / -	0.9/ -	2d	
MEPED	>140	-	-	-	-	

CORONAS F						
MKL	>1.	-	08d01 ^h / -	3.2·10 ³ / -	2d	
MKL	>14	-	08d01 ^h / -	68/ -	2d	
MKL	>26	-	08d01 ^h / -	35/ -	2d	
MKL	>50	-	08d01 ^h / -	6/ -	2d	
ACE						
SIS	>10	17 ^h	21 ^h /09d00 ^h	375/60	2d	
SIS	>30	17 ^h	21 ^h /09d00 ^h	35/1.8	2d	
SOHO						
EPHIN (INT)	>50	17 ^h	21 ^h /09d00 ^h	3.1/0.46	2d	

Differential fluxes of protons for the event of 2004 November 07

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	08 ^h	22 ^h /09d00 ^h	155/213	2d	
LION	2-6	08 ^h	22 ^h /09d00 ^h	74.2/47.3	2d	
EPHIN	4-8	08 ^h	21 ^h /09d00 ^h	400/168	2d	
EPHIN	8-25	08 ^h	21 ^h /09d00 ^h	40/10	2d	
EPHIN	25-41	08 ^h	21 ^h /09d00 ^h	0.9/0.1	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Kuwabara T., Bieber J.W., Clem J., et.al., 2006.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 November.07

2004 November 07 • AR10696 To event 455


Hα	6563 Å	No Flare			n10w15		
1 – 12	keV	1542	1606	1615		X2.0	2.0E-1
12-25	keV	160500	160818	162248		12506936	RHESSI
100-300	keV	162248	162902	163220		3415495	RHESSI
50-150	keV	1620		1627			SONG F
0,18-0,7	MeV						SONG F
15.4	GHz	1552.0	1627.0	0000.0		3.56	
8.8	GHz	1550.0	1627.0	1715.0		3.82	
5	GHz	1552.0	1627.0	0000.0	U1.4 P5	3.92	
2.7	GHz	1555.0	1630.0	0000.0		3.66	
1.4	GHz	1557.0	1625.0	0000.0		3.49	
610	MHz	1602.0	1626.0	1729.0		4.08	
410	MHz	1602.0	1625.0	0000.0		4.18	
245	MHz	1600.0	1625.0	1815.0		4.62	
DS II	25-180	1559		1616		1	
DS IV	25-180	1603		2106		1	
CME	WL	1654	1759 km/s	-19.7km/s ²	360°	000°	



2004 November 08 Ø AR10696 To event 455

H α	6563 Å	1546	1546	1556	N08W35	1N	F
1 – 12	keV	1543	1549	1552		M2.3	6.7E-3
15.4	GHz	1545.0	1546.0	1546.0		3.11	
8.8	GHz	1545.0	1546.0	1546.0	U2.7 P8.8	3.20	
5	GHz	1545.0	1546.0	1547.0		2.78	
2.7	GHz	1546.0	1546.0	~1546.0		1.75	
245	MHz	1546.0	1546.0	~1546.0		3.11	
DS V	25-180	1545		1547		2	
CME	WL						gap

RHESSI - gap



Particle event: To($E_p > 10$ MeV) – 10d02^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 10\text{d}10^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 264 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 10\text{d}16^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 193 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 3}(E_p > 10 \text{ MeV}) - 12\text{d}09^{\text{h}}$, $J_{\max 3}(E_p > 10 \text{ MeV}) - 75 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 6 days

Quasimaximal energy of protons in the event – $E_{qm1} = 485$ MeV

– $E_{qm2} = 430$ MeV

– $E_{qm3} = 110$ MeV

Sources: ● solar flare 09d16^h59^m, M8.9/2N, N07W51, AR10969


○ solar flare 10d01^h59^m, X2.5/3B, N09W49, AR10969

Main X-ray burst 1–8 Å: onset – 09d16^h59^m, max – 09d17^h19^m, $\Phi = 0.094 \text{ J/m}^2$



CME: 09d17^h26^m, $V = 2000 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 299^\circ$;

▲ SC 11d17^h10^m;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2004 November 10



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2004 November 10

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	02 ^h	10 ^h /17 ^h /12d09 ^h	531/492/236	6d	
EPS	>10	02 ^h	10 ^h /16 ^h /12d09 ^h	264/193/75	6d	
EPS	>30	02 ^h	10 ^h /16 ^h /12d08 ^h	40.4/23.7/4.3	4d	
EPS	>50	02 ^h	10 ^h /16 ^h /12d08 ^h	11.1/5.4/0.6	3d	
EPS	>60	02 ^h	10 ^h /16 ^h /12d08 ^h	6.3/2.8/0.22	3d	
EPS	>100	02 ^h	10 ^h /16 ^h /12d08 ^h	1.6/0.7/0.01	3d	
METEOR						
CBM	>5	03 ^h	11 ^h / - / -	108/ - / -	9d	
CBM	>15	03 ^h	11 ^h / - / -	64/ - / -	7d	
CBM	>25	03 ^h	11 ^h / - / -	29/ - / -	5d	
CBM	>40	03 ^h	11 ^h / - / -	11/ - / -	3d	
BP	>90	03 ^h	11 ^h / - / -	1.1/ - / -	2d	
ChD	>600	03 ^h	11 ^h / - / -	0.09/ - / -	1d	
POES-16						
MEPED	>0.24	-	11 ^h /23 ^h /12d10 ^h	- /2.2·10 ³ /1·10 ³	-	
MEPED	>0.8	-	11 ^h /21 ^h /12d10 ^h	1.2·10 ³ /2.3·10 ³ /890	-	
MEPED	>2.5	-	11 ^h /19 ^h /12d10 ^h	1·10 ³ /820/620	-	
MEPED	>6.9	-	11 ^h /17 ^h /12d10 ^h	810/610/230	-	
MEPED	>16	-	11 ^h /17 ^h /12d10 ^h	220/95/30	-	


MEPED	>36	-	11 ^h /17 ^h /12d10 ^h	45/15/3	-	
MEPED	>70	-	11 ^h /17 ^h /12d10 ^h	3.7/1.3/-	-	
MEPED	>140	-	11 ^h /17 ^h /12d10 ^h	1/0.5/-	-	
ACE						
SIS	>10	02 ^h	10 ^h /16 ^h /12d09 ^h	139/222/75	7d	
SIS	>30	02 ^h	10 ^h /16 ^h /12d09 ^h	30/28.4/5.1	4d	
SOHO						
EPHIN (INT)	>50	02 ^h	09 ^h /16 ^h /12d09 ^h	6.1/3.2/0.3	3d	

Differential fluxes of protons for the event of 2004 November 10

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	02 ^h	10 ^h /17 ^h /12d09 ^h	57.8/75.4/65.4	9d	
LION	2-6	02 ^h	10 ^h /16 ^h /12d09 ^h	16.5/22.1/14.8	9d	
EPHIN	4-8	02 ^h	09 ^h /16 ^h /12d09 ^h	96.3/134/75.6	9d	
EPHIN	8-25	02 ^h	09 ^h /16 ^h /12d09 ^h	21.5/21.9/8.0	9d	
EPHIN	25-41	02 ^h	09 ^h /16 ^h /12d09 ^h	1.8/1.0/0.26	9d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2004 November 10

2004 November 9			•	AR10696		To event 456	
H α	6563 Å	1705	1711	1829	N07W51	2N	UZ
DSF	6563 Å	1639		1731	S19W45	23°	
1 – 12	keV	1659	1719	1732		M8.9	9.4E-2
25-50	keV	174028	174110	175528		805713	RHESSI
15.4	GHz	1709.0	1711.0	1753.0		3.08	
8.8	GHz	1701.0	1710.0	0000.0		3.20	
5	GHz	1701.0	1715.0	0000.0		3.30	
2.7	GHz	1701.0	1717.0	0000.0		3.00	
1.4	GHz	1701.0	1713.0	0000.0		2.54	
610	MHz	1702.0	1716.0	0000.0		3.15	
410	MHz	1703.0	1713.0	0000.0		2.88	
245	MHz	1706.0	1714.0	1858.0		3.43	
DS II	25-56	1718		1721		1	
DS II	25-41	1724		1727		3	
DS IV	25-180	1706		2104		3	
DS III	25-150	1659		1702		1	
DS III	25-180	1711		1717		2	
410	MHz	1703.0	1747.0	0000.0		3.26	
CME	WL	1726	2000 km/s	-65.1 km/s ²	360°	299°	




2004 November 10

•

AR10696

To event 456

H α	6563 Å	0204	0210	0315	N09W49	3B	EF
1 – 12	keV	0159	0213	0220		X2.5	1.6E-1
50-100	keV	020056	021058	024024		75870032	RHESSI
2.8	GHz	0200.0	0210.3	0238.0		2.90	
1.4	GHz	0205.0	0210.0	0236.0		3.04	
610	MHz	0207.0	0208.0	0000.0		3.80	
410	MHz	0207.0	0210.0	0236.0		3.73	
245	MHz	0207.0	0209.0	0236.0		3.43	
DS II	25-180	0207		0240		3	
DS IV	130-1200	0206		0237		2	
DS IV	18-180	0247		0410	F,S	3	
DS III	18-1300	0208		0210	G	3	
DS III	25-180	0208		0230	N	3	
CME	WL	0226	3387 km/s	-108.0 km/s ²	360°	302°	



Events in 2005

			Page
1. Event 2005.01.15 – (2005-015)	№ 457	633
2. Event 2005.01.16 – (2005-016)	№ 458	638
3. Event 2005.01.17 – (2005-017) – GLE-68	№ 459	643
4. Event 2005.01.20 – (2005-020) - GLE-69	№ 460	648
5. Event 2005.05.13 – (2005-133)	№ 461	653
6. Event 2005.06.16 – (2005-167)	№ 462	658
7. Event 2005.07.10 – (2005-191)	№ 463	662
8. Event 2005.07.13 – (2005-194)	№ 464	666
9. Event 2005.07.14 – (2005-195)	№ 465	670
10. Event 2005.07.17 – (2005-198)	№ 466	676
11. Event 2005.07.25 – (2005-206)	№ 467	680
12. Event 2005.07.31 – (2005-212)	№ 468	686
13. Event 2005.08.22 – (2005-234)	№ 469	691
14. Event 2005.08.22a – (2005-234a)	№ 470	696
15. Event 2005.09.07 – (2005-250)	№ 471	701
16. Event 2005.09.14 – (2005-257)	№ 472	708

Particle event: To($E_p > 10$ MeV) – 15d07^h

Tmax($E_p > 10$ MeV) – 15d11^h, Jmax ($E_p > 10$ MeV) – $7.4 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 1 day


Quasimaximal energy of protons in the event – $E_{qm} = 300$ MeV

Sources: • solar flare 15d05^h54^m, M8.6/SF, N11E06, AR10720



Main burst X-ray: 1-8 Å: onset – 15d05^h54^m, max – 15d06^h38^m, $\Phi = 0.29 \text{ J/m}^2$

CME: 15d06^h30^m, $V = 2049 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 359^\circ$;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 January 15



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 January 15

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	07 ^h	12 ^h	22	1d	
EPS	>10	07 ^h	11 ^h	7.4	1d	
EPS	>30	07 ^h	10 ^h	1.37	1d	
EPS	>50	07 ^h	10 ^h	0.51	1d	
EPS	>60	07 ^h	10 ^h	0.33	1d	
EPS	>100	07 ^h	10 ^h	0.11	1d	
METEOR						
CBM	>5	07 ^h	11 ^h	2.57	1d	
CBM	>15	07 ^h	11 ^h	1.4	1d	
CBM	>25	07 ^h	11 ^h	0.77	1d	
CBM	>40	07 ^h	11 ^h	0.41	1d	
BP	>90	07 ^h	09 ^h	0.06	1d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	11 ^h	95	1d	
MEPED	>6.9	-	11 ^h	60	1d	
MEPED	>16	-	10 ^h	2.8	1d	
MEPED	>36	-	10 ^h	1.5	1d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
CORONAS F						
MKL	>1.	-	18 ^h	210	1d	
MKL	>14	-	14 ^h	2.5	1d	
MKL	>26	-	13 ^h	1.7	1d	
MKL	>50	-	12 ^h	0.75	1d	

ACE						
SIS	>10	07 ^h	11 ^h	11.6	1d	
SIS	>30	07 ^h	10 ^h	2.7	1d	
SOHO						
EPHIN (INT)	>50	07 ^h	9 ^h	0.16	1d	

Differential fluxes of protons for the event of 2005 January 15

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	08 ^h	19 ^h	16	1d	
LION	2-6	08 ^h	17 ^h	2.5	1d	
EPHIN	4-8	09 ^h	16 ^h	7.2	1d	
EPHIN	8-25	07 ^h	12 ^h	0.5	1d	
EPHIN	25-41	07 ^h	11 ^h	0.031	1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	


References:

Kuwabara T., J.W. Bieber, J. Clem, et.al., 2006.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 January 15

2005 January 15			•	AR10720			To event 457
H α	6563 Å	0556	0559	0817	N12E05	SF	UZ
1 – 12	keV	0554	0638	0717		M8.6	2.9E-1
12-25	keV	053608	053810	054124		18610	RHESSI
12-25	keV	062656	062758	063804		4957986	RHESSI
50-100	keV	063804	064022	065740		6039624	RHESSI
25-50	keV	065740	065822	072352		3280681	RHESSI
50-150	keV	0624		0632			SONG F
80-225	keV						SONG F

15.4	GHz	2211.0	2248.0	0037.0		4.26	
8.8	GHz	2230.0	2248.0	0057.0	U2.7 P8.8	4.32	
5	GHz	2207.0	2250.0	0124.0		4.18	
2.7	GHz	2224.0	2306.0	0102.0		3.81	
1.4	GHz	2227.0	2307.0	0043.0		4.18	
610	MHz	2230.0	2331.0	0045.0		4.87	
410	MHz	2230.0	2329.0	0109.0		5.11	
245	MHz	2236.0	2329.0	0047.0		4.56	
DS II		2224		2258	25-410	3	
DS IV		2233		>2400	18-1600	3	
DS III	GG	2235		2251	18-260	3	
DS III	N	2300		>2400	20-200	3	
DS III	S,C	<0000		0401	18-200	3	
DS UNCLF		2253		2300	20-45	3	
CME		0630	2049 km/s	-30.7 km/s ²	360°	359°	



Particle event: To($E_p > 10$ MeV) – 16d00^h

Tmax($E_p > 10$ MeV) – 16d18^h, Jmax ($E_p > 10$ MeV) – 330 /cm².s.sr

Duration of the event – 1 day

Quasimaximal energy of protons in the event – $E_{qm} = 330$ MeV

Sources: • solar flare 15d21^h54^m, 3B/X2.6, N14W08*, AR10720


3B/M1.0, N14W08*, AR10720

Main X-ray burst 1–8 Å: onset – 15d22^h25^m, max – 15d23^h02^m, $\Phi = 0.63$ J/m²



CME: 15d23^h06^m, $V = 2861$ km/s, $\Delta\phi = 360^\circ$, $dA = 323^\circ$;

*– One flare event with two X-ray bursts

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 January 16



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 January 16

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	00 ^h	18 ^h	890	1.5d	
EPS	>10	00 ^h	18 ^h	330	1.5d	
EPS	>30	00 ^h	15 ^h	45.5	1.5d	
EPS	>50	00 ^h	15 ^h	10.8	1.5d	
EPS	>60	00 ^h	15 ^h	4.1	1.5d	
EPS	>100	00 ^h	15 ^h	0.52	1.5d	
METEOR						
CBM	>5	00 ^h	16 ^h	107	1.5d	
CBM	>15	00 ^h	16 ^h	59	1.5d	
CBM	>25	00 ^h	16 ^h	27	1.5d	
CBM	>40	00 ^h	16 ^h	9,4	1.5d	
BP	>90	00 ^h	16 ^h	0,29	1.5d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	00 ^h	18 ^h	3150	1.5d	
MEPED	>0.8	00 ^h	18 ^h	3000	1.5d	
MEPED	>2.5	00 ^h	16 ^h	1100	1.5d	
MEPED	>6.9	00 ^h	16 ^h	580	1.5d	
MEPED	>16	00 ^h	16 ^h	190	1.5d	
MEPED	>36	00 ^h	16 ^h	12	1.5d	
MEPED	>70	00 ^h	13 ^h	1	1.5d	
MEPED	>140	00 ^h	-	-	-	

CORONAS F						
MKL	>1	-	18 ^h	3400	1.5d	
MKL	>14	-	17 ^h	120	1.5d	
MKL	>26	-	17 ^h	80	1.5d	
MKL	>50	-	17 ^h	20	1.5d	
ACE						
SIS	>10	00 ^h	17 ^h	342	1.5d	
SIS	>30	15d 23 ^h	14 ^h	65	1.5d	
SOHO						
EPHIN (INT)	>50	07 ^h	14 ^h	5.0	1.5d	

Differential fluxes of protons for the event of 2005 January 16


S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	03 ^h	20 ^h	231	1.5d	
LION	2-6	03 ^h	20 ^h	48	1.5d	
EPHIN	4-8	02 ^h	14 ^h	116	1.5d	
EPHIN	8-25	01 ^h	14 ^h	16.7	1.5d	
EPHIN	25-41	01 ^h	14 ^h	1.5	1.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 January 16

2005	January 15	•	AR10720	To event 458			
H α	6563 Å	2154	2249	>0009	N14W08	3B	UZ
1 – 12	keV	2201	2208	2216		M1.0*	6.9E-03
1 – 12	keV	2225	2302	2331		X2.6*	6.3E-01
12-25	keV	221504	221506	222032		78718	RHESSI
300-800	keV	222032	224958	230256		44250936	RHESSI
100-300	keV	230256	230626	231740		22362556	RHESSI
12-25	keV	234832	234930	000308		5883697	RHESSI
50-150	keV	2238		2255			SONG F
225-750	keV	225-750					SONG F

15.4	GHz	2211.0	2248.0	0037.0		4.26	
8.8	GHz	2230.0	2248.0	0057.0	U2.7 P8.8	4.32	
5	GHz	2207.0	2250.0	0124.0		4.18	
2.7	GHz	2224.0	2306.0	0102.0		3.81	
1.4	GHz	2227.0	2307.0	0043.0		4.18	
610	MHz	2230.0	2331.0	0045.0		4.87	
410	MHz	2230.0	2329.0	0109.0		5.11	
245	MHz	2236.0	2329.0	0047.0		4.56	
DS II	25-410	2224		2258		3	
DS IV	18-1600	2233		>2400		3	
DS III	18-260	2235		2251	GG	3	
DS III	20-200	2300		>2400	N	3	
DS III	18-200	<0000		0401	S,C	3	
DS UNCLF		2253		2300		3	
CME	WL	2306	2861 km/s	-127.4km/s ²	360°	323°	

*– One flare event with two X-ray bursts



Particle event: To($E_p > 10$ MeV) – 17d13^h

Tmax($E_p > 10$ MeV) – 17d17^h, Jmax ($E_p > 10$ MeV) – $3.8 \cdot 10^3$ /cm² .s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – $E_{qm} = 750$ MeV


Sources: • solar flare 17d06^h09^m, X3.8/3N, N14W24, AR10720

Main X-ray burst 1-8 Å onset – 17d06^h59^m, max – 17d09^h52^m, $\Phi = 0.84$ J/m²



CME: 17d09^h30^m, $V = 2094$ km/s, $\Delta\phi = 360^\circ$, $dA = 334^\circ$

CME: 17d09^h54^m, $V = 2547$ km/s, $\Delta\phi = 360^\circ$, $dA = 309^\circ$

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 January 17



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 January 17

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES 10						
EPS	>5	13 ^h	17 ^h	5310	2d	
EPS	>10	13 ^h	17 ^h	3820	2d	
EPS	>30	12 ^h	17 ^h	1090	2d	
EPS	>50	12 ^h	17 ^h	330	2d	
EPS	>60	09 ^h	17 ^h	150	2d	
EPS	>100	09 ^h	16 ^h	26	2d	
METEOR						
CBM	>5	10 ^h	22 ^h	950	2d	
CBM	>15	10 ^h	20 ^h	660	2d	
CBM	>25	10 ^h	18 ^h	460	2d	
CBM	>40	10 ^h	18 ^h	240	2d	
BP	>90	10 ^h	17 ^h	13	2d	
ChD	>600	-	-	-	-	
POES 16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	16 ^h	4060	2d	
MEPED	>16	-	16 ^h	2210	2d	
MEPED	>36	-	16 ^h	880	2d	
MEPED	>70	-	15 ^h	62	2d	
MEPED	>140	-	15 ^h	2.35	2d	

CORONAS F						
MKL	>1	-	20 ^h	8150	2d	
MKL	>14	-	19 ^h	1950	2d	
MKL	>26	-	18 ^h	890	2d	
MKL	>50	-	17 ^h	180	2d	
ACE						
SIS	>10	13 ^h	16 ^h	3040	2d	
SIS	>30	11	16 ^h	1020	2d	
SOHO						
EPHIN (INT)	>50	09 ^h	17 ^h	288	2d	
BALLOONS						
Mi	>138		18d06 ^h	1.6		After
Mi	>172		18d06 ^h	0.87		maximum
Mi	>203		18d06 ^h	0.32		- “ -
NM						
Network	>500	-	16 ^h	0.235	-	

Differential fluxes of protons for the event of 2005 January 17

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	11 ^h	18 ^h	196	2d	
LION	2-6	11 ^h	18 ^h	67.2	2d	
EPHIN	4-8	10 ^h	17 ^h	163	2d	
EPHIN	8-25	10 ^h	17 ^h	57.7	2d	
EPHIN	25-41	10 ^h	17 ^h	14.4	2d	


References:

Miroshnichenko L.I. and J. Perez-Peraza, 2008.
Lario D., R.B. Decker, and A. Aran, 2008.
Lario D, A. Aran, R.B. Decker, 2009.
Veslovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 January 17


2005 January 17			•	AR10720		To event 459	
H α	6563 Å	<0906	0951	1157	N14W24	3N	FU
1 – 12	keV	0659	0952	1007		X3.8	8.4E-1
25-50	keV	065956	070258	070920		264551	RHESSI
12-25	keV	070920	071334	072628		718159	RHESSI
50-100	keV	080232	080510	081028		22362556	RHESSI
12-25	keV	234832	234930	000308		7958930	RHESSI
100-300	keV	093536	094942	103852		53152512	RHESSI
50-150	keV	0952		1000			SONG F
150-500	keV						SONG F
500-1300	keV						SONG F
2-6	MeV						SONG F

15.4	GHz	0744.0	0943.0	1207.0		4.23	
8.8	GHz	0755.0	0926.0	1102.0		4.23	
5	GHz	0756.0	0929.0	1207.0	U1.4 P5	4.36	
2.7	GHz	0804.0	0931.0	1207.0		4.08	
1.4	GHz	0853.0	0931.0	1102.0		3.66	
900	MHz	0839.0	0926.7	>1054.0		3.75	
610	MHz	<0845.0	0926.0	>1209.0		4.08	
410	MHz	0812.0	0957.0	1207.0		4.56	
245	MHz	0755.0	1005.0	1207.0		4.59	
204	MHz	0943.1	0943.9	0945.7		4.98	
DS II	26-95	0916		0924		2	
DS II	25-65	0944		0947		3	
DS IV	100-4000	0837		1120	P,S,C	3	
DS IV	25-180	0900		1524		2	
DS I	45-270	<0650		>1200	S,C	2	
DS III	25-270	0923		0934	GG,P	2	
DS III	25-180	0941		0946		3	
DS CONT	25-180	0657		0900		1	
DS CONT	25-270	~0840		>1200	P	2	
DS DCIM	800-2000	0841		1132	GG	3	
CME	WL	0930	2094 km/s	-118.8 km/s ²	360°	334°	
CME	WL	0954	2547 km/s	-159.1 km/s ²	360°	309°	





Particle event: To(Ep>10MeV) – 20d06^hTmax₁(Ep>10MeV) – 20d10^h, Jmax₁(Ep>10MeV) – $1.1 \cdot 10^3$ /cm².s.srTmax₂(Ep>10MeV) – 21d17^h, Jmax₂(Ep>10MeV) – 134 /cm².s.sr

Duration of the event – 4 days


Quasimaximal energy of protons in the event – Eqm₁ = 3840 MeV– Eqm₂ = 1520 MeV**Sources:** • solar flare 20d06^h36^m, X7.1/2B, N12W58, AR10720Main X-ray burst 1-8 Å: onset – 20d06^h36^m, max – 20d07^h01^m, $\Phi = 1.3$ J/m²CME: 20d06^h54^m, V = 882 km/s, $\Delta\phi = 360^\circ$, dA = 288°▲ SC 21d17^h11^m**Particle fluxes and associated phenomena**

Time profiles of the proton fluxes for the event of 2005 January 20



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 January 20

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm².s.sr) ⁻¹	Dura- tion	Comments
GOES 10						
EPS	>5	06 ^h	12 ^h /21d17 ^h	1490/1300	4d	
EPS	>10	06 ^h	10 ^h /21d17 ^h	1310/134	4d	
EPS	>30	06 ^h	08 ^h / -	1130/ -	4d	
EPS	>50	06 ^h	07 ^h / -	940/ -	4d	
EPS	>60	06 ^h	07 ^h / -	820/ -	4d	
EPS	>100	06 ^h	07 ^h / -	490/ -	4d	
METEOR						
CBM	>5	>03 ^h	<21d05 ^h	>140/ -	5d	
CBM	>15	>03 ^h	<21d04 ^h	>81/ -	5d	
CBM	>25	>03 ^h	<21d04 ^h	>45/ -	5d	
CBM	>40	>03 ^h	<21d04 ^h	>26/ -	3d	
BP	>90	>03 ^h	<21d04 ^h	>4.4/ -	2d	
ChD	>600	-	-	-	-	
POES 16						
MEPED	>0.24	-	- /21d18 ^h	- /70800	4d	
MEPED	>0.8	-	- /21d18 ^h	- /35000	4d	
MEPED	>2.5	-	07 ^h /21d18 ^h	11920/11000	4d	
MEPED	>6.9	-	07 ^h /21d18 ^h	10610/1050	4d	
MEPED	>16	-	08 ^h /21d17 ^h	3830/110	4d	
MEPED	>36	-	08 ^h /21d17 ^h	1810/30	4d	
MEPED	>70	-	08 ^h /21d17 ^h	520/5	4d	
MEPED	>140	-	08 ^h /21d17 ^h	110/2	4d	
CORONAS F						
MKL	>1	-	9 ^h /21d17 ^h	5570/3800	4d	
MKL	>14	-	9 ^h /21d17 ^h	1850/16	4d	
MKL	>26	-	9 ^h /21d17 ^h	985/10	4d	
MKL	>50	-	9 ^h /21d17 ^h	385/5	4d	

ACE						
SIS	>10	06 ^h	08 ^h /21d16 ^h	3310/188	4d	
SIS	>30	06 ^h	08 ^h /21d16 ^h	1660/45.6	4d	
SOHO						
EPHIN (INT)	>50	06 ^h	07 ^h /21d16 ^h	1030/9	3d	
BALLOONS						
Mu	>306		07 ^h 45 ^m / -	32.7/ -		
Mu	>482		07 ^h 45 ^m / -	8.7/ -		
Mu	>912		07 ^h 45 ^m / -	1.15/ -		
NM						
Network	>433	-	07 ^h 30 ^m / -	73.6/ -	-	
Network	>1000	-	07 ^h 30 ^m / -	1.57/ -	-	
Network	>2000	-	07 ^h 30 ^m / -	0.065/ -	-	
Network	>3700	-	07 ^h 30 ^m / -	0.0038/ -	-	
Network	>5800	-	07 ^h 30 ^m / -	0.00044/ -	-	

Differential fluxes of protons for the event of 2005 January 20

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	08 ^h	12 ^h /21d17 ^h	30/2430	5d	
LION	2-6	08 ^h	12 ^h /21d17 ^h	26/580	5d	
EPHIN	4-8	08 ^h	11 ^h /21d16 ^h	79/800	4d	
EPHIN	8-25	08 ^h	11 ^h /21d16 ^h	24.6/13	4d	
EPHIN	25-41	07 ^h	08 ^h /21d15 ^h	6.9/0.45	4d	


References:

- Struminsky A., 2005.
 Bieber J.W., J. Clem, P. Evenson et al., 2005.
 Kuwabara T., J.W. Bieber, J. Clem et al., 2006.
 Struminsky A.B., 2006.
 Le G.M., Yu.H. Tang, and Y.B. Han, 2006.
 Kuznetsov S.N., V.G. Kurt, B.Yu. Yushkov et al., 2006.
 Lario D., R.B. Decker, and A. Aran, 2008.
 Miroshnichenko L.I. and J. Perez-Peraza, 2008.
 Perez-Peraza J., A. Gallegos-Cruz, L.I. Miroshnichenko et al., 2008.
 Grechnev V.V., V.G. Kurt, I.M. Chertok et al., 2008.
 Bombardieri D.J., M.L. Duldig, J.E. Humble et al., 2008.
 Lario D., A. Aran, R.B. Decker, 2009.
 Perez-Peraza J., A. Gallegos-Cruz, E.V. Vashenyuk et al., 2009.
 Troitskaia E.V., I.V. Arkhangelskaja, L.I. Miroshnichenko et al., 2009.
 Perez-Peraza J.A., V.M. Velasco-Herrera, J. Zapotitla et al., 2011.
 Miroshnichenko L.I., J.A. Pérez-Peraza, V.M. Velasco-Herrera et al., 2012.
 Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.
 Miroshnichenko L.I. and W.Q. Gan, 2012.
 Velinov P., A. Mishev, 2013.
 Bieber, J.W., J. Clem, P. Evenson et al., 2013.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2005 January 20**


2005 January 20 • AR10720 To event 460

H α	6563 Å	0641	0646	0854	N12W58	2B	UZ
1 – 12	keV	0636	0701	0726		X7.1	1.3E00
300-800	keV	062036	065110	072716		318884832	RHESSI
25-50	keV	075452	075534	084840		11856082	RHESSI
50-150	keV	0944		0956			SONG F
150-500	keV						SONG F
500-1300	keV						SONG F
90-150	MeV						SONG F
15.4	GHz	0642.0	0649.0	0822.0		4.76	
8.8	GHz	0635.0	0647.0	0806.0	U0.9 P8.8	4.79	
5	GHz	0638.0	0649.0	0844.0		4.38	
2.7	GHz	0639.0	0657.0	0840.0		3.95	
1.4	GHz	0639.0	0644.0	0752.0		3.41	
900	MHz	~0703.9	~0703.9	>0727.0		2.68	
610	MHz	0640.0	0643.0	0000.0		3.23	
410	MHz	0639.0	0643.0	0000.0		3.97	
245	MHz	0639.0	0707.0	0832.0		3.23	
DS II	SH	0644		~0700	S,H	3	
DS IV	FS	0636		>0800	F,S	2	
DS IV		0643		0728		3	
DS IV		0643		1055		2	
DS I	N,C	0755		~0950	N,C	2	
DS III	GG	0645		0701	GG	3	
DS III	G	0729		0738	G	3	
DS CONT	DC	0655		0657	DC	2	
DS CONT	DC	0702		0710	DC	2	
DS DCIM	S,C	0830		0856	S,C	2	
DS UNCLF	RS	0755		0755	RS	2	
CME	WL	0654	0882 km/s	16.0km/s ²	360°	288°	





Particle event: To($E_p > 10$ MeV) – 13d19^hTmax₁($E_p > 10$ MeV) – 14d03^h, Jmax₁($E_p > 10$ MeV) – 7.7 /cm².s.srTmax₂($E_p > 10$ MeV) – 14d14^h, Jmax₂($E_p > 10$ MeV) – 155 /cm².s.srTmax₃($E_p > 10$ MeV) – 15d03^h, Jmax₃($E_p > 10$ MeV) – $1.9 \cdot 10^3$ /cm².s.sr

Duration of the event – 2 days


Quasimaximal energy of protons in the event – Eqm₁ = 300 MeV– Eqm₂ = 85 MeV– Eqm₃ = 85 MeV**Sources:** ● solar flare 13d16^h13^m, M8.0/2B, N12E12, AR10759Main X-ray burst 1–8 Å: onset – 13d16^h13^m, max – 13d16^h57^m, $\Phi = 0.18$ J/m²CME: 13d17^h12^m, $V = 1689$ km/s, $\Delta\varphi = 360^\circ$, dA = 002°;▲ SC 15d02^h38^m;**Particle fluxes and associated phenomena**

Time profiles of the proton fluxes for the event of 2005 May 13



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 May 13

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	19 ^h	14d03 ^h /14d14 ^h /15d03 ^h	19.7/740/1·10 ⁴	2d	
EPS	>10	19 ^h	14d03 ^h /14d14 ^h /15d03 ^h	7.7/155/1.9·10 ³	2d	
EPS	>30	20 ^h	14d03 ^h /14d15 ^h /15d03 ^h	1.1/2.8/8.7	2d	
EPS	>50	20 ^h	14d03 ^h /14d15 ^h /15d02 ^h	0.4/0.4/0.52	2d	
EPS	>60	20 ^h	14d03 ^h / - /15d02 ^h	0.28/ - /0.3	2d	
EPS	>100	20 ^h	14d03 ^h / - / -	0.12/ - / -	2d	
METEOR						
CBM	>5	19 ^h	14d03 ^h / - / -	4,2/ - / -	1d	
CBM	>15	19 ^h	14d03 ^h / - / -	1,9/ - / -	1d	
CBM	>25	19 ^h	14d03 ^h / - / -	0,9/ - / -	1d	
CBM	>40	19 ^h	14d01 ^h / - / -	0,56/ - / -	1d	
BP	>90	19 ^h	14d01 ^h / - / -	0,1/ - / -	1d	
ChD	>600	-	-	-	-	
POES-16						
MEPED	>0.24	-	- /14d14 ^h /15d03 ^h	- /4900/110000	2d	
MEPED	>0.8	-	- /14d14 ^h /15d03 ^h	- /2000/40000	2d	
MEPED	>2.5	-	- /14d14 ^h /15d03 ^h	- /1100/20000	2d	
MEPED	>6.9	-	- /14d14 ^h /15d03 ^h	- /500/8000	2d	

MEPED	>16	-	- /14d14 ^h /15d03 ^h	- /35/330	2d	
MEPED	>36	-	- /14d14 ^h /15d03 ^h	- /2.3/3.5	2d	
MEPED	>70	-	- /14d14 ^h / -	- /-/-	2d	
MEPED	>140	-	- /14d14 ^h / -	- /-/-	2d	
CORONAS F						
MKL	>1.	-	- /14d14 ^h /15d02 ^h	-/6300/8300	2d	
MKL	>14	-	- /14d14 ^h /15d02 ^h	-/25///110	2d	
MKL	>26	-	- /14d14 ^h /15d02 ^h	-/13/33	2d	
MKL	>50	-	- /14d14 ^h /15d02 ^h	-/3/3//	2d	
ACE						
SIS	>10	19 ^h	- /14d14 ^h /15d02 ^h	- /262/2·10 ³	3d	
SIS	>30	19 ^h	- /14d14 ^h /15d02 ^h	- /5/10	2d	
SOHO						
EPHIN (INT)	>50	21 ^h	14d04 ^h /14d13 ^h /15d03 ^h	0.17/0.4/1.1	1d	


Differential fluxes of protons for the event of 2005 May 13

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	21 ^h	14d02 ^h /14d14 ^h /15d03 ^h	6.2/148/4.3·10 ³	4d	
LION	2-6	18 ^h	14d02 ^h /14d14 ^h /15d03 ^h	2.4/57/1.5·10 ³	4d	
EPHIN	4-8	21 ^h	14d02 ^h /14d14 ^h /15d03 ^h	8.5/304/2.7·10 ³	4d	
EPHIN	8-25	20 ^h	14d02 ^h /14d14 ^h /15d03 ^h	0.62/24.1/173	4d	
EPHIN	25-41	18 ^h	14d02 ^h /14d14 ^h /15d03 ^h	0.025/0.15/0.29	4d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 May.13

2005	May 13	•	AR10759	To event 461			
Hα	6563 Å	1631	1641	1946	N12E12	2B	MZ
1 – 12	keV	1613	1657	1758		M8.0	1.8E-1
50-100	keV	163628	165458	171420		46101460	RHESSI

15.4	GHz	1640.0	1645.0	0000.0		2.76	
8.8	GHz	1639.0	1645.0	0000.0		2.98	
5	GHz	1634.0	1643.0	1735.0		3.23	
2.7	GHz	1633.0	1703.0	0000.0		3.46	
1.4	GHz	1633.0	1705.0	0000.0		3.73	
610	MHz	1633.0	1705.0	0000.0	P0.6	4.63	
410	MHz	1634.0	1657.0	1713.0		4.28	
245	MHz	1633.0	1648.0	0000.0		2.62	
DS II	40-350	1638		1650	F,S,H	2	
DS II	25-81	1641		1652		3	
DS IV	40-600	1633		~1828	F,S	3	
DS IV	30-180	1645		2057		3	
DS III	25-153	1641		1641		1	
DS DCIM	800-2000	1628		~1723	GG,F,S	3	
DS DCIM	2000-4500	1632		1639	G	2	
CME	WL	1712	1689 km/s	–	360°	002°	



Particle event: To($E_p > 10$ MeV) – 16d20^h

Tmax($E_p > 10$ MeV) – 17d04^h, Jmax ($E_p > 10$ MeV) – 41 /cm².s.sr

Duration of the event – 2 days


Quasimaximal energy of protons in the event – $E_{qm} = 510$ MeV

Sources: ■ solar flare 16d20^h01^m, M4.0/SF, N09W87, AR10775



Main X-ray burst 1-8 Å: onset – 16d20^h01^m, max – 16d20^h22^m, $\Phi = 0.062$ J/m²

CME: gap

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 June 16



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 June 16

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	20 ^h	17d04 ^h	57	2d	
EPS	>10	20 ^h	17d04 ^h	41	2d	
EPS	>30	20 ^h	17d02 ^h	16	2d	
EPS	>50	20 ^h	23 ^h	8.7	2d	
EPS	>60	20 ^h	23 ^h	6.5	2d	
EPS	>100	20 ^h	23 ^h	2.6	2d	
POES-16						
MEPED	>0.24	20 ^h	-	-	-	
MEPED	>0.8	20 ^h	-	-	-	
MEPED	>2.5	20 ^h	-	-	-	
MEPED	>6.9	20 ^h	23 ^h	110	2d	
MEPED	>16	20 ^h	23 ^h	36	2d	
MEPED	>36	20 ^h	23 ^h	25	2d	
MEPED	>70	20 ^h	23 ^h	4.5	2d	
MEPED	>140	20 ^h	23 ^h	0.8	2d	
ACE						
SIS	>10	20 ^h	17d02 ^h	84.5	2d	
SIS	>30	20 ^h	17d02 ^h	40	1d	
SOHO						
EPHIN (INT)	>50	21 ^h	17d01 ^h	2.8	1d	

Differential fluxes of protons for the event of 2005 June 16

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	22 ^h	17d04 ^h	2.3	3d	
LION	2-6	22 ^h	17d04 ^h	0.9	3d	

Particle event: To($E_p > 10$ MeV) – 10d03^h

$T_{\max 1}(E_p > 10 \text{ MeV}) - 10\text{d}05^{\text{h}}$, $J_{\max 1}(E_p > 10 \text{ MeV}) - 1.1 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

$T_{\max 2}(E_p > 10 \text{ MeV}) - 10\text{d}12^{\text{h}}$, $J_{\max 2}(E_p > 10 \text{ MeV}) - 1.9 / \text{cm}^2 \cdot \text{s} \cdot \text{sr}$

Duration of the event – 2.5 days

Quasimaximal energy of protons in the event – $E_{qm1} = 75 \text{ MeV}$

– $E_{qm2} = 70 \text{ MeV}$


Sources: • solar flare 09d21^h47^m, M2.8/1N, N11W27, AR10786

Main X-ray burst 1-8 Å: onset – 09d21^h47^m, max – 09d22^h06^m, $\Phi = 0.029 \text{ J/m}^2$



CME: 09d22^h30^m, $V = 1540 \text{ km/s}$, $\Delta\phi = 360^\circ$, $dA = 328^\circ$;

▲ SC 10d03^h37^m;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 July 10



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 July 10

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	03 ^h	05 ^h /12 ^h	2.4/6.5	2.5d	
EPS	>10	03 ^h	05 ^h /12 ^h	1.1/1.9	2.5d	
EPS	>30	03 ^h	04 ^h /12 ^h	0.2/0.15	2.5d	
EPS	>50	-	04 ^h /12 ^h	0.06/0.03	2d	
EPS	>60	-	04 ^h /12 ^h	0.02/0.01	2d	
EPS	>100	-	-	-	-	
ACE						
SIS	>10	01 ^h	05 ^h /11 ^h	1.1/3	1d	
SIS	>30	02 ^h	05 ^h /12 ^h	0.2/0.3	1d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	


Differential fluxes of protons for the event of 2005 July 10

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	05 ^h	06 ^h /11 ^h	2.5/2.4	2.5d	
LION	2-6	05 ^h	06 ^h /11 ^h	0.18/0.5	2.5d	
EPHIN	4-8	04 ^h	- /11 ^h	- /1.1	2.5d	
EPHIN	8-25	02 ^h	- /11 ^h	- /0.12	2.5d	
EPHIN	25-41	01 ^h	- /11 ^h	- /0.31	2.5d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2005 July 10

2005 July 09 • AR10786 To event 463

H α	6563 Å	2156	2204	2238	N11W27	1N	FU
1 – 12	keV	2147	2206	2219		M2.8	2.9E-2
15.4	GHz	2158.0	2207.0	0000.0		2.85	
8.8	GHz	2155.0	2206.0	0000.0		3.00	
5	GHz	2152.0	2206.0	0000.0		3.23	
2.8	GHz	2147.0	2209.0	2209.0	P0.6 U1.4 P2.8	3.44	
1.4	GHz	2156.0	2208.0	2217.0		2.85	
610	MHz	2149.0	2214.0	2220.0		4.08	
410	MHz	2150.0	2208.0	2217.0		3.97	
245	MHz	2153.0	2208.0	2218.0		3.08	
DS II	23-90	2159		2205	S,H	3	
DS II	20-180	2203		2217	S,H	2	
DS IV	25-2000	2152		2223		2	
DS IV	20-180	2155		>2400	F,S	2	
DS III	20-80	2203		2209	GG	3	
DS UNCLF	130-180	2211		2215		2	
CME	WL	2230	1540 km/s	-168.5 km/s	360°	328°	



Particle event: To($E_p > 10$ MeV) – 13d17^h

Tmax($E_p > 10$ MeV) – 15d04^h, Jmax ($E_p > 10$ MeV) – 9.7 /cm².s.sr

Duration of the event – 1 day


Quasimaximal energy of protons in the event – $E_{qm} = 115$ MeV

Sources: • solar flare 13d14^h01^m, M5.0/SF, N10W80, AR10786



Main X-ray burst 1-8 Å: onset – 13d14^h01^m, max – 13d14^h49^m, $\Phi = 0.2$ J/m²

CME: 13d14^h30^m, $V = 1423$ km/s, $\Delta\phi = 360^\circ$, dA = 303°;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 July 13



Arrows on the profiles of the events indicate the time of the proton flux maxima,
taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 July 13

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	17 ^h	14d05 ^h	43	1d	
EPS	>10	17 ^h	14d04 ^h	9.7	1d	
EPS	>30	17 ^h	14d02 ^h	0.6	1d	
EPS	>50	17 ^h	14d01 ^h	0.12	1d	
EPS	>60	17 ^h	14d01 ^h	0.06	1d	
EPS	>100	17 ^h	14d01 ^h	0.01	1d	
POES-16						
MEPED	>0.24	-	14d02 ^h	400	1d	
MEPED	>0.8	-	14d02 ^h	320	1d	
MEPED	>2.5	-	14d02 ^h	150	1d	
MEPED	>6.9	-	14d02 ^h	40	1d	
MEPED	>16	-	14d02 ^h	4.5	1d	
MEPED	>36	-	14d02 ^h	1.2	1d	
MEPED	>70	-	14d02 ^h	0.2	1d	
MEPED	>140	-	14d02 ^h	-	-	
ACE						
SIS	>10	17 ^h	14d04 ^h	15	1d	
SIS	>30	18 ^h	14d05 ^h	1.2	1d	
SOHO						
EPHIN (INT)	>50	17 ^h	14d02 ^h	0.05	1d	

Differential fluxes of protons for the event of 2005 July 13

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	21 ^h	14d05 ^h	14.5	1d	
LION	2-6	17 ^h	14d05 ^h	3.4	1d	


EPHIN	4-8	17 ^h	14d05 ^h	9.8	1d	
EPHIN	8-25	17 ^h	14d03 ^h	0.74	1d	
EPHIN	25-41	17 ^h	14d02 ^h	0.02	1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Malandraki O.E., N. Agueda, A. Papaioannou et al., 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 July 13

2005		July 13		•		AR10786		To event 464	
H α	6563 Å	1422	1437	1458	N10W80		SF	F	
1 – 12	keV	1401	1449	1538			M5.0	2.0E-1	
100-300	keV	141300	143522	144404			20887532	RHESSI	
12-25	keV	144404	151422	151740			490896	RHESSI	
15.4	GHz	1401.0	1417.0	1505.0	P0.2 U1.4 [P5-15]		3.73		
8.8	GHz	1401.0	1417.0	0000.0			3.71		
5	GHz	1358.0	1418.0	1447.0			3.76		
2.7	GHz	1401.0	1419.0	1443.0			3.30		
1.4	GHz	1358.0	1420.0	1440.0			2.32		
610	MHz	1345.0	1401.0	1412.0			3.43		
410	MHz	1351.0	1403.0	1421.0			3.72		
245	MHz	1357.0	~1403.0	1422.0			3.82		
127	MHz	1411.6	1422.3	1424.6			2.80		
DS III	30-180	1406		1411			2		
DS V	25-180	1401		1409			2		
CME	WL	1430	1423 km/s	-14.1 km/s	360°		303°		



Particle event: To(Ep>10MeV) – 14d14^h

Tmax(Ep>10MeV) – 15d03^h, Jmax (Ep>10MeV) – 130 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – E_{qm} = 185 MeV

Sources: ■ solar flare 14d10^h16^m, X1.2/..., n11w90*, AR10786

○ solar flare 14d05^h57^m, M9.1/1N, N11W73, AR10786

○ solar flare 14d22^h50^m, M1.1/..., N09W90, AR10786


Main X-ray burst 1-8 Å: onset – 14d10^h16^m, max – 14d10^h55^m, Φ = 0.39 J/m²

CME: 14d10^h54^m, V = 2115 km/s, Δφ = 360°, dA = 296°;



▲ SC 17d01^h34^m;

* – probable localization of the flare event

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 July 14



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 July 14

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm².s.sr)⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	14 ^h	15d04 ^h	250	3d	
EPS	>10	14 ^h	15d03 ^h	130	3d	
EPS	>30	14 ^h	15d00 ^h	12.3	3d	
EPS	>50	14 ^h	15d05 ^h	2.08	3d	
EPS	>60	14 ^h	15d05 ^h	0.63	3d	
EPS	>100	-	15d05 ^h	0.06	3d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	15d00 ^h	350	3d	
MEPED	>16	-	15d05 ^h	65	3d	
MEPED	>36	-	15d05 ^h	8.4	3d	
MEPED	>70	-	15d05 ^h	0.6	3d	
ACE						
SIS	>10	13 ^h	15d01 ^h	141	3d	
SIS	>30	12 ^h	15d02 ^h	16.4	3d	
SOHO						
EPHIN (INT)	>50	14 ^h	15d01 ^h	0.26	2d	

Differential fluxes of protons for the event of 2005 July 14

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm².s.sr.MeV)⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	15 ^h	15d02 ^h	14.4	3d	
LION	2-6	15 ^h	15d01 ^h	5.4	3d	
EPHIN	4-8	15 ^h	15d04 ^h	20	3d	
EPHIN	8-25	15 ^h	15d04 ^h	4.1	3d	
EPHIN	25-41	15 ^h	15d03 ^h	0.3	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Chernetsky VA., M.A. Livshits, L.K. Kashapova et al., 2009.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 July 14

2005 July 14





AR10786

To event 465

H α	6563 Å	No Flare Patrol			n13w80		
1 – 12	keV	1016	1055	1129		X1.2	3.9E-1
25-50	keV	101508	102742	112956		4398593	RHESSI
25-50	keV	105952	110402	112548		14154225	RHESSI
15.4	GHz	1019.0	1034.0	1144.0	[U1.4]/15	3.79	
8.8	GHz	1020.0	1034.0	1145.0		3.72	
5	GHz	1020.0	1035.0	1143.0		3.43	
2.7	GHz	1021.0	1033.0	0000.0		2.32	
1.4	GHz	1029.0	1033.0	0000.0		1.49	
610	MHz	1033.0	1033.0	0000.0		2.57	
410	MHz	1025.0	1032.0	0000.0		2.00	
245	MHz	1026.0	1032.0	0000.0		2.86	
DS IV	200-4000	1023		1212	P,C	3	
DS IV	25-180	1025		1146		2	
DS III	25-47	1019		1019		1	
DS DCIM	2000-4500	1018		1128	GG	3	
DS DCIM	800-2000	1021		1149	GG,S,P	3	
9.1	GHz	1018.7	1056.3			3.48	
2.7	GHz	1021.0	1059.0	1136.0		3.53	
1.4	GHz	1029.0	1107.0	0000.0		4.11	
610	MHz	1047.0	1110.0	0000.0	P0.6	5.08	
410	MHz	1025.0	1111.0	1222.0		4.72	
245	MHz	1026.0	1111.0	1211.0		3.73	
CME	WL	1054	2115 km/s	198.0km/s ²	360°	296°	

* – probable localization of the flare event






2005 July 14

Ø

AR10758

To event 465

H α	6563 Å	0723	0724	0732	N11W73	1N	FM
1 – 12	keV	0557	0725	0743		M9.1	8.4E-2
12-25	keV	061224	061926	070720		1402202	RHESSI
6-12	keV	070720	070926	071348		63166	RHESSI
6-12	keV	071348	071550	071644		25014	RHESSI
50-150	keV	0723		0724			SONG F
2-6	MeV						SONG F
15.4	GHz	0722.0	0723.0	0723.0	[U0.9-2.7]/15	3.00	
8.8	GHz	0722.0	0723.0	0723.0		2.91	
5	GHz	0722.0	0723.0	0723.0		2.69	
2.7	GHz	0722.0	0723.0	0723.0		2.34	
1.4	GHz	0722.0	0723.0	0723.0		2.46	
900	MHz	0722.3	0722.8	0730.9		2.26	
610	MHz	0723.0	0723.0	~0723.0		3.04	
410	MHz	0722.0	0722.0	0724.0		2.45	
245	MHz	0722.0	0723.0	0724.0		2.11	
DS III	25-700	0722		0727	G	3	
DS III	25-600	0729		0738	G	1	
DS V	25-180	0722		0726		2	
DS DCIM	200-4000	0722		0737	P,C	3	
DS DCIM	800-2000	0735		0737	G	2	
CME	WL	0754	0752 km/s	-2.6 km/s ²	103°	237°	




2005 July 14

Ø

AR10758

To event 465

H α	6563 Å	No Flare Patrol			n13w89		
1 – 12	keV	2250	2257	2309		M1.1	4.8E-3
12-25	keV	225104	225822	230920		652561	RHESSI
5	GHz	2254.0	2254.0	2255.0		1.89	
2.7	GHz	2254.0	2255.0	2255.0	U0.6 P2.7	2.00	
1.4	GHz	2254.0	2255.0	2255.0		1.76	
610	MHz	<2253.0	2253.0	>2253.0		1.38	
410	MHz	<2253.0	2253.0	>2253.0		1.94	
245	MHz	2256.0	2256.0	2257.0		1.97	
DS III	18-180	2253		2307	GG	3	
CME	WL	2330	0724 km/s	0.2 km ² /s	033°	270°	



Particle event: To(Ep>10MeV) – 17d14^h

Tmax₁(Ep>10MeV) – 17d18^h, Jmax₁(Ep>10MeV) – 12 /cm².s.sr

Tmax₂(Ep>10MeV) – 17d22^h, Jmax₂(Ep>10MeV) – 19 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 85 MeV


– Eqm₂ = 85 MeV

Sources: ☐ solar flare 17d10^h25^m, B1.1/..., N13W9, AR10786 - 3d behind W limb;



Main X-ray burst 1-8 Å: onset – 17d10^h25^m, max – 17d10^h32^m, Φ = 0.00013 J/m²

CME: 17d11^h30^m, V = 1527 km/s, Δφ = 360°, dA=316°;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 July 17



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 July 17

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	14 ^h	18 ^h /22 ^h	23/41	3d	
EPS	>10	14 ^h	18 ^h /22 ^h	12/19	3d	
EPS	>30	13 ^h	18 ^h /22 ^h	2.6/2.7	3d	
EPS	>50	13 ^h	18 ^h /22 ^h	0.6/0.54	2d	
EPS	>60	13 ^h	18 ^h /22 ^h	0.16/0.16	2d	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	-	18 ^h /-	230/-	3d	
MEPED	>0.8	-	18 ^h /-	150/-	3d	
MEPED	>2.5	-	18 ^h /-	70/-	3d	
MEPED	>6.9	-	18 ^h /-	50/-	3d	
MEPED	>16	-	18 ^h /-	9/-	3d	
MEPED	>36	-	18 ^h /-	4/-	2d	
MEPED	>70	-	18 ^h /-	0.3/-	2d	
MEPED	>140	-	-	-	-	
ACE						
SIS	>10	14 ^h	- /21 ^h	- /30	2.5d	
SIS	>30	13 ^h	- /20 ^h	- /6	2d	
SOHO						
EPHIN (INT)	>50	15 ^h	19 ^h /22 ^h	0.1/0.08	1d	

Differential fluxes of protons for the event of 2005 July 17

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	12 ^h	19 ^h /23 ^h	4.9/5.85	4d	
LION	2-6	12 ^h	19 ^h /23 ^h	0.9/1.3	4d	
EPHIN	4-8	12 ^h	19 ^h /23 ^h	2.4/4	3d	
EPHIN	8-25	15 ^h	19 ^h /23 ^h	0.33/0.6	3d	
EPHIN	25-41	15 ^h	19 ^h /23 ^h	0.05/0.054	3d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 July 17

2005 July 17



AR10786

To event 466

H α	6563 Å	No Flare			n13w90		
1 – 12	keV	1025	1032	1045		B1.1	1.3E-4
6-12	keV	102652	102758	103008		23160	RHESSI
CME	WL	1130	1527 km/s	59.2 km/s ²	360°	316°	

Particle event: To($E_p > 10$ MeV) – 25d21^h

Tmax₁($E_p > 10$ MeV) – 28d14^h, Jmax₁($E_p > 10$ MeV) – 30 /cm².s.sr

Tmax₂($E_p > 10$ MeV) – 29d14^h, Jmax₂($E_p > 10$ MeV) – 36 /cm².s.sr

Duration of the event – 6 days

Quasimaximal energy of protons in the event – Eqm₁ = 85 MeV

– Eqm₂ = 85 MeV

Sun source: □ solar flare 27d04^h 33^m, M3.7/..., n10w90* AR10792, 1d behind E-limb;

○ solar flare 28d21^h 39^m, M4.8/..., N09E82 AR10792


Main X-ray burst 1-8 Å: onset – 26d04^h 33^m, max – 26d05^h 02^m, $\Phi = 0.0002$ J/m²

CME: 26d04^h 54^m, V=1458 km/s, $\Delta\phi = 360^\circ$, dA = 90°.



▲ SC 27d19^h 39^m;

* – probable localization of the flare even

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 July 25



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 July 25

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES 10						
EPS	>5	21 ^h	28d14 ^h /29d14 ^h	73.5/104	6d	
EPS	>10	21 ^h	28d14 ^h /29d14 ^h	30/36	6d	
EPS	>30	21 ^h	28d14 ^h /29d14 ^h	2.7/2.4	6d	
EPS	>50	21 ^h	28d14 ^h /29d13 ^h	0.5/0.44	6d	
EPS	>60	21 ^h	28d15 ^h /29d13 ^h	0.25/0.2	6d	
EPS	>100	-	-	-	-	
POES 16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	- /29d12 ^h	- /250	6d	
MEPED	>2.5	-	- /29d12 ^h	- /170	6d	
MEPED	>6.9	-	28d16 ^h /29d12 ^h	140/120	6d	
MEPED	>16	-	28d16 ^h /29d12 ^h	10/10	6d	
MEPED	>36	-	28d16 ^h /29d12 ^h	1.5/2.5	5d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
ACE						
SIS	>10	21 ^h	28d13 ^h /29d14 ^h	42.3/48.4	6d	
SIS	>30	21 ^h	28d13 ^h /29d06 ^h	4/3.2	6d	
SOHO						
EPHIN (INT)	>50	24d00 ^h	28d14 ^h /29d10 ^h	0.33/0.32	3d	

Differential fluxes of protons for the event of 2005 July 25

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	21 ^h	28d14 ^h /29d14 ^h	4.8/14.5	6d	
LION	2-6	21 ^h	28d14 ^h /29d14 ^h	2/4.6	6d	
EPHIN	4-8	21 ^h	28d14 ^h /29d14 ^h	13.1/26.7	6d	
EPHIN	8-25	21 ^h	28d14 ^h /29d13 ^h	2.4/3.45	6d	
EPHIN	25-41	24d00 ^h	28d14 ^h /29d13 ^h	0.12/0.12	6d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 July 25

2005 July 27

Ø



AR10792

To event 467

H α	6563 Å	No Flare			n10e90*		
EPL	6563 Å	0439		0511	N11E90		
1 – 12	keV	0433	0502	0530		M3.7	7.9E-2
50-100	keV	04:49:56	04:57:34	05:42:12		200402302	RHESSI
12-25	keV	14:44:04	15:14:22	15:17:40		490896	RHESSI
15.4	GHz	0450.0	0457.0	0530.0		2.79	
8.8	GHz	0447.0	0456.0	0535.0		3.08	
5	GHz	0447.0	0457.0	0457.0	U1.4 P5	3.18	
2.7	GHz	0445.0	0457.0	0457.0		2.86	
1.4	GHz	0445.0	0457.0	0457.0		1.99	
610	MHz	0444.0	0454.0	0512.0		2.26	
410	MHz	0450.0	0453.0	0506.0		2.49	
245	MHz	0445.0	0458.0	0503.0		2.74	
DS II	20-130	0443		0503	S,H	3	
DS IV	25-180	0446		0524		1	
DS IV	200-4000	0452		0523	P	3	
DS III	20-180	0445		0535	S,C	1	
DS UNCLF	40-60	0454		0458		2	

2.8	GHz	0430.0	0507.3	0610.0	U0.9 / 2.8	2.97	
1.4	GHz	0444.0	0507.0	0531.0		2.43	
900	MHz	0439.3	0508.5			2.32	
410	MHz	0444.0	0522.0	0535.0		2.84	
DS DCIM	800-2000	0510		0542	G	2	
CME	WL	0454	1787km/s	-75.4 km/s	360°	084°	

* – probable localization of the flare event



2005


July 28

Ø

AR10792

To event 467

H α	6563 Å	2151	2200	2221	N08E84	SF	FH
1 – 12	keV	2139	2208	2224		M4.8	8.1E-2
12-25	keV	221244	221342	231452		2447943	RHESSI
15.4	GHz	2142.0	2154.0	2210.0	U0.6-1.4 / 15	3.34	
8.8	GHz	2142.0	2153.0	2220.0		3.23	
5	GHz	2142.0	2154.0	2220.0		2.90	
2.8	GHz	2147.0	2154.0	2209.0		2.05	
1.4	GHz	2152.0	2204.0	2204.0		1.53	
610	MHz	2201.0	2201.0	~2201.0		1.54	
410	MHz	2201.0	2201.0	~2201.0		1.85	
245	MHz	2158.0	2204.0	2205.0		1.76	
DS II	110-170	2153		2157	UE	1	
DS IV	30-220	2158		2214		1	
DS III	20-180	2157		2221	S,C	1	
DS III	27-90	2206		2206	B	2	
DS III	20-90	2218		2218	G	3	
CME	WL	2206	1478 km/s				



Particle event: To(Ep>10MeV) – 31d22^h

Tmax₁(Ep>10MeV) – 01d05^h, Jmax₁(Ep>10MeV) – 21 /cm².s.sr

Tmax₂(Ep>10MeV) – 02d01^h, Jmax₂(Ep>10MeV) – 6 /cm².s.sr

Duration of the event – 5 days

Quasimaximal energy of protons in the event – Eqm₁ = 65 MeV

– Eqm₂ = 65 MeV


Sources: ☉ solar flare 30d06^h17^m, X1.3/2B, N12E61, AR10792

☾ solar flare 01d13^h00^m, M1.0/1F, N14E29, AR10792



Main X-ray burst 1-8 Å: onset – 30d16^h17^m, max – 30d06^h35^m, Φ = 0.23 J/m²

CME: 30d06^h50^m, V = 1968 km/s, Δφ = 360°, dA = 050°.

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 July 31



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 July 31

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	22 ^h	01d06 ^h /02d02 ^h	122/49.3	6d	
EPS	>10	22 ^h	01d05 ^h /02d01 ^h	21/6	5d	
EPS	>30	22 ^h	01d02 ^h /02d00 ^h	0.5/0.15	2d	
EPS	>50	22 ^h	01d02 ^h / -	0.03/ -	1d	
EPS	>60	22 ^h	-	-	-	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	-	01d06 ^h /02d04 ^h	1230/2620	6d	
MEPED	>0.8	-	01d06 ^h /02d04 ^h	910/1710	6d	
MEPED	>2.5	-	01d06 ^h /02d02 ^h	640/390	6d	
MEPED	>6.9	-	01d05 ^h /02d02 ^h	150/90	5d	
MEPED	>16	-	01d05 ^h /02d02 ^h	4.5/1.5	5d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
ACE						
SIS	>10	22 ^h	01d05 ^h /02d02 ^h	31.3/7.6	6d	
SIS	>30	-	01d05 ^h /02d01 ^h	0.6/0.3	2d	
SOHO						
EPHIN (INT)	>50	24d00 ^h	01d05 ^h /02d03 ^h	0.2/0.17	1.5d	

Differential fluxes of protons for the event of 2005 July 31

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						

LION	0.75-2	22 ^h	01d07 ^h /02d04 ^h	78/80.5	6d	
LION	2-6	22 ^h	01d07 ^h /02d03 ^h	16.2/9.7	6d	
EPHIN	4-8	21 ^h	01d07 ^h /02d04 ^h	68.3/25.1	6d	
EPHIN	8-25	21 ^h	01d07 ^h /02d02 ^h	3.1/0.82	6d	
EPHIN	25-41	24d00 ^h	01d05 ^h /02d02 ^h	0.03/0.006	6d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2005 July 31**


2005 July 30

•

AR10792

To event 468

H α	6563 Å	0608	0625	0823	N11E59	2N	UZ
1 – 12	keV	0617	0635	0701		X1.3	2.3E-1
100-300	keV	062740	063042	071348		82189624	RHESSI
6-12	keV	081736	081902	081908		48672	RHESSI
12-25	keV	091808	082118	082312		131232	RHESSI
15.4	GHz	0621.0	0625.0	0704.0		3.36	
8.8	GHz	0620.0	0632.0	0715.0		3.51	
5	GHz	0619.0	0632.0	0754.0		3.61	
2.7	GHz	0619.0	0636.0	0754.0		3.32	
1.4	GHz	0621.0	0625.0	0716.0		3.75	
900	MHz	0543.5	0626.1	0848.0	[P1]	3.91	
610	MHz	0621.0	0626.0	0720.0		3.38	
410	MHz	0621.0	0626.0	0719.0		3.23	
245	MHz	0621.0	0632.0	0641.0		3.00	
127	MHz	<0630.0	0633.4	>0642.0		3.66	
DS II	20-180	0626		0640	S,H	3	
DS IV	200-4000	0621		0847	P,S	3	
DS IV	25-180	0634		0734		2	
DS III	18-180	0626		0634	GG	3	
DS DCIM	2000-4500	0619		0803	GG	3	
CME	WL	0650	1968 km/s	-102.6 km/s ²	360°	050°	




2005 August 01

Ø

AR10792

To event 468

H α	6563 Å	1315	1340	1436	N14E29	1F	EF
1 – 12	keV	1300	1351	1429		M1.0	2.9E-2
6-12	keV	130244	130250	130820		3064	RHESSI
12-25	keV	130820	131718	132356		77104	RHESSI
25-50	keV	132356	132950	133448		263876	RHESSI
15.4	GHz	1329.0	1343.0	1416.0		2.26	
8.8	GHz	1328.0	1343.0	1411.0		2.56	
5	GHz	1328.0	1343.0	1416.0	U1.4 P5	2.67	
2.7	GHz	1329.0	1347.0	1416.0		2.46	
1.4	GHz	1341.0	1347.0	0000.0		2.11	
610	MHz	1342.0	1343.0	1349.0		2.53	
410	MHz	1341.0	1350.0	1418.0		2.53	
245	MHz	1345.0	1349.0	1418.0		2.76	
127	MHz	1355.0	1405.6	1425.0		2.90	
DS II	200-500	1349		1359		3	
DS IV	25-180	1349		1743		1	
DS DCIM	200-1600	1409		1419	P	3	
CME	WL	1430	0984 km/s	17.9 km/s ²	093°	068°	



Particle event: To(Ep>10MeV) – 22d03^h

Tmax(Ep>10MeV) – 22d07^h, Jmax (Ep>10MeV) – 5.4 /cm².s.sr

Duration of the event – 0.7 day


Quasimaximal energy of protons in the event – Eqm = 80 MeV

Sources: • solar flare 22d00^h44^m, M2.6/1F, S09W48, AR10798



Main X-ray burst 1-8 Å: onset – 22d00^h44^m, max – 22d01^h33^m, $\Phi = 0.096$ J/m²

CME: 22d01^h32^m, V=1194 km/s, $\Delta\phi = 360^\circ$, dA= 220°

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 August 22



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 August 22

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	03 ^h	08 ^h	20.1	0.7d	
EPS	>10	03 ^h	07 ^h	5.4	0.7d	
EPS	>30	03 ^h	05 ^h	0.48	0.7d	
EPS	>50	03 ^h	05 ^h	0.12	0.7d	
EPS	>60	03 ^h	05 ^h	0.06	0.7d	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	-	-	-	
MEPED	>16	-	07 ^h	2.5	1d	
MEPED	>36	-	-	-	-	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
ACE						
SIS	>10	03 ^h	06 ^h	11.4	0.7d	
SIS	>30	03 ^h	05 ^h	1.1	0.7d	
SOHO						
EPHIN (INT)	>50	-	-	-	-	

Differential fluxes of protons for the event of 2005 August 22

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	03 ^h	10 ^h /	7.4	0.7d	
LION	2-6	03 ^h	07 ^h /	1.15	0.7d	
EPHIN	4-8	03 ^h	09 ^h	5.8	0.7d	
EPHIN	8-25	03 ^h	06 ^h	0.53	0.7d	
EPHIN	25-41	03 ^h	06 ^h	0.02	0.7d	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 August 22

2005 August 22

•


AR10798

To event 469

H α	6563 Å	0104	0117	0300	S09W48	1F	FH
1 – 12	keV	0044	0133	0218		M2.6	9.6E-3
25-50	keV	010148	012134	020240		31078596	RHESSI
6-12	keV	023752	023846	031408		707931	RHESSI
15.4	GHz	0053.0	0122.0	0000.0		2.88	
8.8	GHz	0054.0	0124.0	0301.0		3.11	
5	GHz	0051.0	0124.0	0301.0		3.30	
2.7	GHz	0051.0	0124.0	0245.0		3.28	
1.4	GHz	0053.0	0137.0	0219.0		3.34	
610	MHz	0050.0	0138.0	0257.0		3.97	
410	MHz	0051.0	0138.0	0000.0		4.08	
245	MHz	0054.0	0140.0	0000.0	[0.2 \ 15]	4.30	
DS II	90-430	0054		0110		1	
DS II	30-60	0102		0108	S,H	3	
DS IV	50-1000	0101		0315		1	
DS IV	25-180	0103		0316	F,S	3	
DS III	20-90	0101		0122	GG	3	
DS III	20-50	0130		0135	G	3	

410	MHz	0051.0	0204.0	0402.0		4.38	
245	MHz	0054.0	0205.0	0000.0		4.89	
CME	WL	0132	1194 km/s	-17.8	360°	220°	

/



Particle event: To(Ep>10MeV) – 22d19^h

Tmax₁(Ep>10MeV) – 23d02^h, Jmax₁(Ep>10MeV) – 280 /cm².s.sr

Tmax₂(Ep>10MeV) – 23d10^h, Jmax₂(Ep>10MeV) – 290 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 330 MeV

– Eqm₂ = 290 MeV


Sources: ● solar flare 22d16^h52^m, M5.6/1N, S12W60, AR10798

Main X-ray burst 1-8 Å: onset – 22d16^h52^m, max – 22d17^h27^m, Φ = 0.17 J/m²



CME: 22d17^h30^m, V=2378 km/s, Δφ = 360°, dA= 220°;

▲ SC 24d06^h13^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 August 22a



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 August 22a

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	19 ^h	23d02 ^h /23d10 ^h	540/880	3d	
EPS	>10	19 ^h	23d02 ^h /23d10 ^h	280/290	3d	
EPS	>30	19 ^h	23d02 ^h /23d10 ^h	24.9/16.7	2d	
EPS	>50	19 ^h	23d02 ^h /23d10 ^h	4.1/2.2	2d	
EPS	>60	19 ^h	23d02 ^h /23d10 ^h	1.6/1	2d	
EPS	>100	18 ^h	23d02 ^h /23d10 ^h	0.19/0.09	1d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	- /23d10 ^h	- /3120	3d	
MEPED	>2.5	-	- /23d10 ^h	- /2140	3d	
MEPED	>6.9	-	23d02 ^h /23d10 ^h	760/650	3d	
MEPED	>16	-	23d02 ^h /23d10 ^h	180/120	3d	
MEPED	>36	-	23d02 ^h / -	18.1/ -	2d	
MEPED	>70	-	23d02 ^h / -	1/ -	2d	
MEPED	>140	-	-	-	-	

ACE						
SIS	>10	19 ^h	23d02 ^h /23d10 ^h	350/308	4 d	
SIS	>30	19 ^h	23d02 ^h /23d10 ^h	28.5/19	4 d	
SOHO						
EPHIN (INT)	>50	19 ^h	23d01 ^h /23d09 ^h	1.05/1	2d	

Differential fluxes of protons for the event of 2005 August 22a



S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	19 ^h	23d01 ^h /23d09 ^h	51/75.4	3d	
LION	2-6	19 ^h	23d01 ^h /23d09 ^h	13.3/33.7	3d	
EPHIN	4-8	19 ^h	23d01 ^h /23d09 ^h	66.8/200	3d	
EPHIN	8-25	19 ^h	23d01 ^h /23d09 ^h	13.3/24	3d	
EPHIN	25-41	19 ^h	23d01 ^h /23d09 ^h	0.62/0.68	3d	

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 August 22a

2005 August 22 • AR10798 To event 470

H α	6563 Å	1652	1711	1845	S12W60	1N	FU
1 – 12	keV	1646	1727	1802		M5.6	1.7E-1
100-300	keV	170116	171610	174320		67140128	RHESSI
6-12	keV	183712	183838	191308		301914	RHESSI
15.4	GHz	1707.0	1711.0	2006.0		3.92	
8.8	GHz	1707.0	1711.0	2014.0		3.99	
5	GHz	1649.0	1711.0	1717.0		4.08	
2.7	GHz	1649.0	1713.0	1717.0		3.85	
1.4	GHz	1651.0	1714.0	1717.0	P5	3.11	
DS IV	100-4000	1652		1737		3	
DS IV	30-180	1654		2145	P	3	

8.8	GHz	1654.0	1721.0	1915.0		3.59	
5	GHz	1654.0	1721.0	1936.0	U1.4 P5	4.00	
2.7	GHz	1654.0	1722.0	1937.0		3.91	
1.4	GHz	1654.0	1722.0	1921.0		3.41	
610	MHz	1655.0	1720.0	1907.0		3.66	
410	MHz	1655.0	1728.0	2035.0		4.18	
245	MHz	1656.0	1741.0	2035.0		4.52	
DS III	25-87	1720		1721		1	
CME	WL	1730	2378 km/s	108.0km/s	360°	227°	



Particle event: To(Ep>10MeV) – 07d21^h

Tmax₁(Ep>10MeV) – 08d20^h, Jmax₁(Ep>10MeV) – 70 /cm².s.sr

Tmax₂(Ep>10MeV) – 10d11^h, Jmax₂(Ep>10MeV) – 1000 /cm².s.sr

Duration of the event – 5 days

Quasimaximal energy of protons in the event – Eqm₁ = 800 MeV

– Eqm₂ = 555 MeV

Sources: ● solar flare 07d17^h17^m, X17.0/3B, S06E89, AR10808

Ø solar flare 08d20^h52^m, X5.4/2B, S12E62, AR10808

Ø solar flare 09d09^h42^m, X3.6/1N, S11E64, AR10808

Ø solar flare 09d19^h13^m, X6.2/2B, S12E67*, AR10808


Main X-ray burst 1-8 Å: onset – 07d17^h17^m, max – 07d17^h40^m, Φ = 2.6 J/m²

CME: gap



▲ SC 09d14^h01^m; ▲ SC 11d01^h14^m

* – localisation on the X-ray burst

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 September 07



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 September 07

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm².s.sr)⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	21 ^h	08d20 ^h /10d12 ^h	123/2000	5d	
EPS	>10	21 ^h	08d20 ^h /10d11 ^h	70/1000	5d	
EPS	>30	21 ^h	08d20 ^h /10d09 ^h	26.7/170	5d	
EPS	>50	21 ^h	08d20 ^h /10d06 ^h	15.3/47.4	5d	
EPS	>60	21 ^h	08d19 ^h /10d05 ^h	12.2/27.4	5d	
EPS	>100	21 ^h	08d19 ^h /10d03 ^h	6.8/6.9	5d	
POES-16						
MEPED	>0.24	-	- /10d12 ^h	- /2920	5d	
MEPED	>0.8	-	- /10d12 ^h	- /2810	5d	
MEPED	>2.5	-	- /10d10 ^h	- /2230	5d	
MEPED	>6.9	-	08d20 ^h /10d09 ^h	210/2140	5d	
MEPED	>16	-	08d20 ^h /10d06 ^h	95/460	5d	
MEPED	>36	-	08d20 ^h /10d04 ^h	16/150	5d	
MEPED	>70	-	08d20 ^h /10d04 ^h	10/10.3	5d	
MEPED	>140	-	08d20 ^h /10d04 ^h	3/1.6	5d	
ACE						
SIS	>10	20 ^h	08d21 ^h /10d11 ^h	233/1080	5d	
SIS	>30	20 ^h	08d21 ^h /10d11 ^h	94/216	5d	
SOHO						
EPHIN (INT)	>50	20 ^h	08d20 ^h /10d04 ^h	7.6/47.5	5d	

BALLOONS						
Mi	>134	-	08d09 ^h 25 ^m / -	1.5/ -	-	
Mi	>216	-	08d09 ^h 25 ^m / -	0.42/ -	-	
Mi	>492	-	08d09 ^h 25 ^m / -	0.09/ -	-	
Mi	>136	-	- /09d10 ^h 30 ^m	- /6.9	-	Before max2
Mi	>218	-	- /09d10 ^h 30 ^m	- /0.93	-	- “ -
Mi	>316	-	- /09d10 ^h 30 ^m	- /0.13	-	- “ -

Differential fluxes of protons for the event of 2005 September 07

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	20 ^h	08d19 ^h /10d09 ^h	7.4/304	5d	
LION	2-6	20 ^h	08d19 ^h /10d10 ^h	3.3/71.3	5d	
EPHIN	4-8	20 ^h	08d19 ^h /10d09 ^h	13/252	5d	
EPHIN	8-25	20 ^h	08d19 ^h /10d08 ^h	2.4/30.7	5d	
EPHIN	25-41	20 ^h	08d19 ^h /10d08 ^h	0.27/2.9	5d	

References:

Kuwabara T., Bieber J.W., Clem J., et.al., 2006.
Struminsky A.B., 2011.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 September 07


2005 September 07

•

AR10808



To event 471

H _α	6563 Å	1724	1740	1847	S06E89	3B	MY
1 – 12	keV	1717	1740	1803		X17.0	2.6E00
6-12	keV	171104	171334	171752		585282	RHESSI
50-100	keV	174244	174346	174704		3848547	RHESSI
50-100	keV	181904	182002	182752		21973536	RHESSI
25-50	keV	182752	183230	185936		52805952	RHESSI
15.4	GHz	1723.0	~1736.0	1902.0	[0.2 / 15]	4.57	
8.8	GHz	1723.0	~1736.0	1902.0		4.36	
5	GHz	<1725.0	~1736.0	>1902.0		4.38	
2.7	GHz	<1728.0	~1736.0	>1910.0		4.43	
1.4	GHz	<1729.0	~1732.0	>1905.0		4.20	
610	MHz	1732.0	~1738.0	1853.0		3.76	
410	MHz	1733.0	~1736.0	1843.0		3.77	
245	MHz	1735.0	~1737.0	1841.0		3.51	
DS II	25-180	1742		1750		3	
DS IV	25-180	1750		1846		2	
CME	WL						gap




2005 September 08 Ø AR10808 To event 471

H _α	6563 Å	2052	2105	0042	S11E74	2B	MU
1 – 12	keV	2052	2106	2117		X5.4	3.8E-1
25-50	keV	213016	213118	222056		9633607	RHESSI
12-25	keV	223324	223334	223424		19872	RHESSI
12-25	keV	230632	230654	234616		457791	RHESSI
6-12	keV	234616	234654	000208		78348	RHESSI
15.4	GHz	2026.0	2107.0	2148.0		3.41	
8.8	GHz	2053.0	2105.0	2142.0		3.11	
5	GHz	2101.0	2105.0	0000.0		2.76	
2.7	GHz	2102.0	2104.0	0000.0		3.00	
1.4	GHz	2105.0	2106.0	2145.0		3.18	
610	MHz	2111.0	2111.0	~2111.0		1.75	
410	MHz	2114.0	2114.0	2121.0		2.04	
245	MHz	2121.0	2121.0	~2121.0		1.72	
DS IV	25-300	2116		2337		1	
DS III	30-180	2110		2131	N	1	
DS III	40-200	2110		2112	G	3	
9.5	GHz	2135.2	2139.2	2143.1		2.31	
5	GHz	2101.0	2135.0	2143.0		2.94	
1.4	GHz	<2104.0	2137.0	>2146.0	U0.4 P1.4	3.15	
610	MHz	2111.0	2135.0	2138.0		2.18	
410	MHz	2114.0	2137.0	2151.0		2.11	
245	MHz	2122.0	2134.0	2203.0		2.32	
DS IV	23-57	2132		>2400	F,S	2	
DS IV	38-180	2244		0344		2	
DS III	20-57	2130		>2400	N	1	
CME	WL						gap



2005 September 09 Ø AR10792 To event 471

H _α	6563 Å	<1020	–	1115	S12E62	2B	MU
1 – 12	keV	0942	0959	1008		X3.6	2.3E-1
25-50	keV	092524	094602	094636		279628	RHESSI
12-25	keV	101800	101854	103316		2530704	RHESSI
12-25	keV	110024	110042	110848		32808	RHESSI
12-25	keV	110848	110914	112040		33560	RHESSI
15.4	GHz	0953.0	0956.0	1024.0		3.18	
8.8	GHz	0952.0	0956.0	1019.0		2.97	
5	GHz	0953.0	0956.0	1019.0		2.72	
2.7	GHz	0954.0	0956.0	1012.0		2.43	
1.4	GHz	0957.0	1005.0	1011.0		3.38	
610	MHz	1000.0	1001.0	1011.0		2.58	
410	MHz	1006.0	1007.0	1007.0		1.85	
245	MHz	1006.0	1006.0	1025.0		2.45	
DS IV	100-4000	0954		1024	P	3	
DS III	50-180	1025		1025		2	
DS DCIM	2000-4500	0952		1026	GG	3	
DS DCIM	800-2000	0954		1026	GG,F,S	3	
CME	WL						gap



2005 September 09


Ø

AR10808

To event 471

H _α	6563 Å	No Flare			S12E67		
1 – 12	keV	1913	2004	2036		X6.2	1.7E00
12-25	keV	191908	192146	192240		604126	RHESSI
50-100	keV	205652	205658	205840		531774	RHESSI
15.4	GHz	1916.0	1947.0	2147.0		4.49	
8.8	GHz	1916.0	1948.0	0000.0		4.34	
5	GHz	1916.0	~1949.0	2143.0		4.15	
2.7	GHz	1921.0	~1949.0	2142.0		3.62	
1.4	GHz	1905.0	1931.0	0000.0		3.96	
610	MHz	1929.0	1945.0	0000.0		3.11	
410	MHz	1925.0	~1945.0	2036.0		3.00	
235	MHz	1931.9	~1931.9	2048.0		>3.47	
DS II	25-180	1934		1949		2	
DS IV	30-180	1934		2123		3	
CME	WL	1948	2257 km/s	-128.6 km/s ²	360°	220°	

* – localisation on the X-ray burst



Particle event: To(Ep>10MeV) – 14d00^h

Tmax₁(Ep>10MeV) – 14d15^h, Jmax₁(Ep>10MeV) – 160 /cm².s.sr

Tmax₂(Ep>10MeV) – 15d08^h, Jmax₂(Ep>10MeV) – 180 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – Eqm₁ = 85 MeV

– Eqm₂ = 85 MeV

Sources: ● solar flare 13d19^h19^m, X1.5/2B, S09E10*, AR10808

13d23^h15^m, X1.7/2B, S09E10*, AR10808

○ solar flare 15d08^h30^m, X1.1/2N, S11W15, AR10808


Main X-ray burst 1-8 Å: onset – 13d19^h19^m, max – 13d19^h27^m, Φ = 0.055 J/m²

CME: 13d20^h00^m, V = 1866 km/s, Δφ = 360°, dA = 149°



▲ SC 15d08^h35^m; Δ SC15d09^h04^m;

* – One solar flare event with two X-ray burst

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2005 September 14



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2005 September 14

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Duration	Comments
GOES-10						
EPS	>5	00 ^h	16 ^h /15d08 ^h	460/740	3d	
EPS	>10	00 ^h	15 ^h /15d08 ^h	160/180	3d	
EPS	>30	00 ^h	14 ^h /15d08 ^h	9.4/5.2	2d	
EPS	>50	00 ^h	14 ^h /15d08 ^h	1.2/0.5	2d	
EPS	>60	00 ^h	14 ^h /15d08 ^h	0.3/0.13	2d	
EPS	>100	-	-	-	-	
POES-16						
MEPED	>0.24	-	- /15d08 ^h	- /4580	3d	
MEPED	>0.8	-	14 ^h /15d08 ^h	1050/4210	3d	
MEPED	>2.5	-	14 ^h /15d08 ^h	750/2050	2d	
MEPED	>6.9	-	14 ^h /15d07 ^h	370/820	2d	
MEPED	>16	-	14 ^h /15d06 ^h	50/50	2d	
MEPED	>36	-	14 ^h /15d06 ^h	6.2/3.4	1d	
MEPED	>70	-	-	-	-	
MEPED	>140	-	-	-	-	
ACE						
SIS	>10	00 ^h	15 ^h /15d06 ^h	222/263	3d	
SIS	>30	00 ^h	15 ^h /15d06 ^h	13.7/7.2	2d	

SOHO						
EPHIN (INT)	>50	00 ^h	15 ^h /15d08 ^h	0.68/0.6	2d	

Differential fluxes of protons for the event of 2005 September 14

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Duration	Comments
SOHO						
LION	0.75-2	00 ^h	15 ^h /15d08 ^h	73/245	3d	
LION	2-6	00 ^h	15 ^h /15d08 ^h	23/59	3d	
EPHIN	4-8	00 ^h	15 ^h /15d08 ^h	126/281	3d	
EPHIN	8-25	00 ^h	15 ^h /15d08 ^h	17.7/23.5	3d	
EPHIN	25-41	00 ^h	15 ^h /15d08 ^h	0.58/0.32	2d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	

References:

Kuwabara T., Bieber J.W., Clem J., et al., 2006.
Struminsky A.B., 2011.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2005 September 14

2005 September 13



•

AR10808

To event 472

H α	6563 Å	1922	1923	0021	S09E10	2B	FU
1 – 12	keV	1919	1927	2057		X1.5	5.5E-1
1 – 12	keV	2315	2322	2330		X1.7	9.3E-2
25-50	keV	195252	195822	201344		21140692	RHESSI
12-25	keV	201344	201510	202224		5285486	RHESSI
100-300	keV	202224	203254	203316		6368830	RHESSI
50-100	keV	204308	204330	205716		5777847	RHESSI
12-25	keV	212900	213046	215352		1368541	RHESSI
12-25	keV	215352	215638	221632		912288	RHESSI
12-25	keV	230456	230522	231052		116139	RHESSI
6-12	keV	231052	321234	231500		47715	RHESSI
100-300	keV	231500	232150	000540		22934672	RHESSI
15.4	GHz	<1922.0	1923.0	>2131.0		4.23	
8.8	GHz	<1921.0	1923.0	>2129.0	[U1.4] P8.8	4.38	
5	GHz	1921.0	1923.0	0000.0		4.23	
2.7	GHz	1922.0	1923.0	2137.0		3.78	
1.4	GHz	<1922.0	2021.0	>2135.0		3.30	
610	MHz	1929.0	2015.0	2059.0		4.18	
410	MHz	1937.0	2029.0	2153.0		2.56	
245	MHz	1930.0	2043.0	2157.0		2.85	
DS IV	20-57	<2010		>2400		3	
DS III	25-180	1950		1951		2	
DS III	25-135	2027		0422	N	1	
DS III	25-57	2041		2041	B	3	
DS V	30-180	1950		1951		2	


15.4	GHz	2317.0	2318.0	0000.0	0.4 / 15	3.38	
8.8	GHz	2317.0	2320.0	0000.0		3.20	
8.8	GHz	2317.0	2320.0	2337.0		3.20	
5	GHz	2318.0	2320.0	2330.0		2.97	
2.8	GHz	2314.0	2320.5	2354.0		2.37	
410	MHz	2319.0	2320.0	2321.0		1.93	
CME	WL	2000	1866 km/s	11.5 km/s ²	360°	149°	
CME	WL	2336	0999 km/s	-13.9 km/s ²	046°	158°	



2005 September 15 Ø AR10808 To event 472

H α	6563 Å	0834	0837	0953	S11W15	2N	FZ
1 – 12	keV	0830	0838	0846		X1.1	5.6E-2
3-6	keV	08:40:28	08:41:10	08:42:24		1256426	RHESSI
6-12	keV	08:42:24	09:11:22	09:34:04		646872	RHESSI

15.4	GHz	0834.0	0836.0	0932.0	U0.9 / 15	3.61	
8.8	GHz	0834.0	0836.0	0922.0		3.51	
5	GHz	0834.0	0836.0	0932.0		3.46	
3	GHz	0834.5	0836.8			2.78	
1.4	GHz	0836.0	0836.0	0000.0		1.48	
900	MHz	0835.1	0838.1	0840.3		1.34	
245	MHz	0840.0	0840.0	~0840.0		1.72	
DS DCIM	2000-4500	0834		0849	G	3	
DS DCIM	1000-4000	0835		0842	C	3	
CME	WL						gap



Events in 2006

		Page
1.	Event 2006.12.05 – (2006-339) № 473	715
2.	Event 2006.12.06 – (2006-340) № 474	720
3.	Event 2006.12.13 – (2006-347) – GLE-70 № 475	725
4.	Event 2006.12.14 – (2006-348) № 476	731

Particle event: To(Ep>10MeV) – 05d15^h

Tmax(Ep>10MeV) – 05d20^h, Jmax(Ep>10MeV) – 2.5 /cm².s.sr

Duration of the event – 1 day


Quasimaximal energy of protons in the event – E_{qm} = 275 MeV

Sources: • solar flare 05d10^h18^m, X9.0/2N, S07E79, AR10930;



Main x-ray burst 1–8 Å: onset – 05d10^h18^m, max – 05d10^h35^m, Φ = 0.71 J/m²

CME: gap

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2006 December 05



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2006 December 05

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm².s.sr)⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	15 ^h	20 ^h	3.1	1d	
EPS	>10	15 ^h	20 ^h	2.5	1d	
EPS	>30	15 ^h	21 ^h	0.85	1d	
EPS	>50	15 ^h	21 ^h	0.3	1d	
EPS	>60	15 ^h	21 ^h	0.2	1d	
EPS	>100	16 ^h	22 ^h	0.09	1d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	-	-	-	
MEPED	>6.9	-	20 ^h	100	1d	
MEPED	>16	-	20 ^h	3	1d	
MEPED	>36	-	20 ^h	1.7	1d	
MEPED	>70	-	18 ^h	0.2	1d	
MEPED	>140	-	-	-	-	
ACE						
SIS	>10	14 ^h	20 ^h	5	1d	
SIS	>30	14 ^h	20 ^h	1.3	1d	
SOHO						
EPHIN (INT)	>50		-	-	-	

Differential fluxes of protons for the event of 2006 December 05

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	-	20 ^h	0.18	1d	
LION	2-6	-	20 ^h	0.08	1d	
EPHIN	4-8	-	19 ^h	0.56	1d	
EPHIN	8-25	08 ^h	19 ^h	0.17	1d	
EPHIN	25-41	08 ^h	19 ^h	0.02	1d	
EPHIN	41-53	- “ -	- “ -	- “ -	- “ -	
STEREO – A						
HET	13.6-29.5	12 ^h	23 ^h	0.35	1d	
HET	29.5-60	12 ^h	23 ^h	0.04	1d	
HET	60-100	12 ^h	23 ^h	0.001	1d	
STEREO - B						
HET	13.6-29.5	12 ^h	23 ^h	0.3	1d	
HET	29.5-60	12 ^h	23 ^h	0.04	1d	
HET	60-100	12 ^h	23 ^h	0.0015	1d	


References:

Struminsky A. and I. Zimovets I., 2008.
Cohen C.M.S., G.M. Mason, R.A. Mewaldt et al., 2009.
Damiani A., M. Storini, M. Santee et al., 2009.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2006 December 05

2006 December 05		•		AR10808		To event 473	
H α	6563 Å	1028	1038	1100	S07E79	2N	F
1 – 12	keV	1018	1035	1045		X9.0	7.1E-01
25-50	keV	103032	103938	114424		87206176	RHESSI

15.4	GHz	1025.0	1030.0	1105.0	P0.2 U0.6[P2.7 U 5]/15	4.04	
8.8	GHz	1025.0	1030.0	1058.0		3.83	
5	GHz	1025.0	1030.0	1057.0		3.61	
2.7	GHz	1024.0	1029.0	1056.0		4.08	
1.4	GHz	1027.0	1030.0	1037.0		3.77	
610	MHz	1027.0	1028.0	1044.0		2.64	
410	MHz	1026.0	1029.0	1059.0		2.99	
245	MHz	1027.0	1027.0	1105.0		5.43	
127	MHz	1026.0	1032.2	1042.0		2.83	
DS II	100-300	1028		1036	H	3	
DS II	38-73	1034		1039		3	
DS IV	25-180	1034		1054		2	
DS III	110-400	1026		1033	GG,RS	3	
DS III	25-180	1027		1034		2	
DS DCIM	100-4000	1025		1234	P	3	
CME	WL						gap



Particle event: To($E_p > 10$ MeV) – 06d10^h

Tmax($E_p > 10$ MeV) – 07d22^h, Jmax($E_p > 10$ MeV) – 1860 /cm².s.sr

Duration of the event – 5 days

Quasimaximal energy of protons in the event – $E_{qm} = 850$ MeV


Sources: ● solar flare 06d18^h29^m, X6.5/3B, S06E63 AR10930

Main x-ray burst 1–8 Å: onset – 06d 18^h 29^m, max – 06d18^h47^m, $\Phi = 0.480$ J/m²



CME: 06d20^h12^m, $V = \dots$ km/s, $\Delta\phi = 360^\circ$, $dA = 135^\circ$.

▲ SC 08d 0435

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2006 December 06



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2006 December 06

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	10 ^h	7d23 ^h	4460	6d	
EPS	>10	10 ^h	7d22 ^h	1860	6d	
EPS	>30	10 ^h	7d18 ^h	360	6d	
EPS	>50	10 ^h	7d16 ^h	112	5d	
EPS	>60	10 ^h	7d16 ^h	66	5d	
EPS	>100	10 ^h	7d12 ^h	18	4d	
POES-16						
MEPED	>0.24	-	-	-	-	
MEPED	>0.8	-	-	-	-	
MEPED	>2.5	-	7d23 ^h	7220	-	
MEPED	>6.9	-	7d23 ^h	6130	-	
MEPED	>16	-	7d20 ^h	1230	-	
MEPED	>36	-	7d18 ^h	310	-	
MEPED	>70	-	7d14 ^h	26	-	
MEPED	>140	-	7d13 ^h	6.5	-	
ACE						
SIS	>10	10 ^h	7d18 ^h	1990	6d	
SIS	>30	10 ^h	7d16 ^h	420	6d	
SOHO						
EPHIN (INT)	>50	10 ^h	7d20 ^h	86.8	5d	

BALLONS						
Mi	>345		8d(09 ^h 37 ^m -09 ^h 57 ^m)	0.42		12 hours after
Mi	>483		8d(09 ^h 37 ^m -09 ^h 57 ^m)	0.25		maximum
Mi	>642		8d(09 ^h 37 ^m -09 ^h 57 ^m)	0.14		

Differential fluxes of protons for the event of 2006 December 06

S/c, instruments	ΔE, MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	-	7d23 ^h	980	6d	
LION	2-6	-	7d23 ^h	310	6d	
EPHIN	4-8	-	7d22 ^h	950	6d	
EPHIN	8-25	10 ^h	7d22 ^h	90	6d	
EPHIN	25-41	10 ^h	7d21 ^h	4	6d	
STEREO – A						
HET	13.6-29.5	12 ^h	8d00 ^h	267	6.5d	
HET	29.5-60	12 ^h	8d00 ^h	27.5	6.5d	
HET	60-100	12 ^h	8d00 ^h	0.7	6.5d	
STEREO - B						
HET	13.6-29.5	12 ^h	8d00 ^h	342	6.5d	
HET	29.5-60	12 ^h	8d00 ^h	29.3	6.5d	
HET	60-100	12 ^h	8d00 ^h	0.83	6.5d	

References:


Struminsky A. and I. Zimovets, 2008.
Cohen C.M.S., G.M. Mason, R.A. Mewaldt et al., 2009.
Damiani A., M. Storini, M. Santee et al., 2009.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2006 December 06

2006 December 06 • AR10930 To event 474

Hα	6563 Å	1832	1845	2135	S06E63	3B	FZ
1 – 12	keV	1829	1847	1900		X6.5	4.8E-1
300 800	keV	183232	184526	192232		60269264	RHESSI
6-12	keV	200740	200810	201332		120125	RHESSI
25-50	keV	201332	201746	204628		1984508	RHESSI
6-12	keV	204628	204750	204956		22362	RHESSI
6-12	keV	204956	205042	205436		26997	RHESSI
15.4	GHz	1838.0	1847.0	1937.0	[U1.4 P8.8-15]	4.08	
8.8	GHz	1833.0	1855.0	1959.0		4.11	
5	GHz	1832.0	1855.0	1959.0		4.00	
2.7	GHz	1842.0	1855.0	0000.0		3.88	
1.4	GHz	1841.0	1847.0	0000.0		3.45	

610	MHz	1842.0	1843.0	0000.0		4.00	
410	MHz	1844.0	1845.0	0000.0		3.93	
245	MHz	1844.0	1844.0	1855.0		4.49	
DS II	25-180	1842		1859		3	
DS IV	30-180	1844		1902		3	
DS III	30-180	1842		1844		2	
9.5	GHz	1916.2	1917.2	1959.6		2.79	
1.4	GHz	1842.0	1917.0	0000.0		5.18	
610	MHz	1844.0	1918.0	0000.0		4.28	
CME	WL	2012	—	—	360°	135°	



Particle event: To(Ep>10MeV) – 13d03^h

Tmax(Ep>10MeV) – 13d09^h, Jmax (Ep>10MeV) – 660 /cm².s.sr

Duration of the event – 2 days

Quasimaximal energy of protons in the event – E_{qm} = 3440 MeV


Sources: ● solar flare 13d02^h14^m, X3.4/4B, S06W24, AR10930

Main X-ray burst 1–8 Å: onset – 13d02^h14^m, max – 13d02^h40^m, Φ = 0.51 J/m²



CME: 13d02^h54^m, V = 1774 km/s, Δφ = 360°, dA = 193°;

▲ SC 14d14^h14^m

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2006 December 13



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2006 December 13

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm ² .s.sr) ⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	03 ^h	09 ^h	750	2d	
EPS	>10	03 ^h	09 ^h	660	2d	
EPS	>30	02 ^h	07 ^h	360	2d	
EPS	>50	02 ^h	06 ^h	227	2d	
EPS	>60	02 ^h	05 ^h	180	2d	
EPS	>100	01 ^h	05 ^h	86	2d	
POES-16						
MEPED	>0.8		-	-	-	
MEPED	>2.5		-	-	-	
MEPED	>6.9		05 ^h	1120	2d	
MEPED	>16		06 ^h	630	2d	
MEPED	>36		06 ^h	410	2d	
MEPED	>70		06 ^h	110	2d	
MEPED	>140		05 ^h	40	2d	
ACE						
SIS	>10	03 ^h	06 ^h	1375	2d	
SIS	>30	03 ^h	06 ^h	812	2d	
SOHO						
EPHIN (INT)	>50	02 ^h	06 ^h	122	2d	

BALLOONS						
Mu	>245	-	08 ^h 24 ^m -09 ^h 11 ^m	13.9	-	
Mu	>428	-	08 ^h 24 ^m -09 ^h 11 ^m	1.3	-	
Mu	>968	-	08 ^h 24 ^m -09 ^h 11 ^m	0.36	-	
NM						
Network	>433	-	04 ^h	46.8		
Network	>1000	-	04 ^h	0.915	-	
Network	>2000	-	04 ^h	0.035	-	
Network	>3700	-	04 ^h	0.002	-	

Differential fluxes of protons for the event of 2006 December 13

S/c, instruments	ΔE , MeV	To	Tmax	Jmax, (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	04 ^h	08 ^h	6,5	2d	
LION	2-6	04 ^h	08 ^h	4	2d	
EPHIN	4-8	05 ^h	09 ^h	13.2	2d	
EPHIN	8-25	04 ^h	09 ^h	10.2	2d	
EPHIN	25-41	03 ^h	09 ^h	4	2d	
STEREO-A						
HET	13.6-29.5	03 ^h	12 ^h	44	2d	
HET	29.5-60	03 ^h	12 ^h	13.5	2d	
HET	60-100	03 ^h	12 ^h	0.9	2d	
STEREO-B						
HET	13.6-29.5	03 ^h	12 ^h	39.3	2d	
HET	29.5-60	03 ^h	12 ^h	13.1	2d	
HET	60-100	03 ^h	12 ^h	0.9	2d	
PAMELA						
Tracker	126-136	03 ^h	05 ^h	0.23	>1.5d	
Tracker	237-265	03 ^h	04 ^h	0.068	>1.5d	
Tracker	455-512	03 ^h	03 ^h	0.016	>1.5d	
Tracker	713-804	03 ^h	03 ^h	0.0057	>1.5d	
Tracker	1260-1420	03 ^h	03 ^h	0.00051	>1.5d	
Tracker	2230-2520	03 ^h	03 ^h	0.0000134	>1.5d	

References:

- Bieber J.H., J. Clem, P. Evenson et al., 2008.
Heber B., A. Struminsky, I. Zimovets et al., 2008.
Cohen C.M.S., G.M. Mason, R.A. Mewaldt et al., 2009.
De Simone N., O. Adriani, G.C. Barbarino et al., 2009.
Damiani A., M. Storini, M. Santee et al., 2009.
Adriani O., G.C. Barbarino, G.A. Bazilevskaya et al., 2011.
Wang X. and Y. Yan, 2012.
Mulligan T., J.B. Blake, R.A. Mewaldt, 2008.
Miroshnichenko L.I. and J. Perez-Peraza, 2008.
Struminsky A.B., I.V. Zimovec, 2009.
Li C., Dai Y., J.-C. Vial et al., 2009.
Luhmann J.G., S.A. Ledvina, D. Odstrcil et al., 2010.
Verkhoglyadova O.P., G. Li, G.P. Zank et al., 2010.
Maurchev E.A., Yu.V Balabin., E.V. Vashenyuk et al., 2013.

**Electromagnetic and other phenomena that are sources and/or accompanying for the event of
2006 December 13**

2006 December 13

•


AR10930

To event 475

H α	6563 Å	0220	0234	0618	S06W24	4B	EF
1 – 12	keV	0214	0240	0257		X3.4	5.1E-1
100-300	keV	022812	023858	033104		248619072	RHESSI
6-12	keV	040408	040518	050640		4424582	RHESSI
15.4	GHz	0221.0	0229.0	0343.0	U5 / 15	4.00	
8.8	GHz	0221.0	0229.0	0325.0		3.85	
5	GHz	0222.0	0233.0	0326.0		3.67	
2.7	GHz	0222.0	0223.0	0000.0		4.15	
1.4	GHz	0223.0	0225.0	0326.0		5.18	
610	MHz	0223.0	0224.0	0000.0		3.67	
245	MHz	0224.0	0225.0	0326.0		5.00	
DS II	25-300	0227		0244		3	
DS IV	25-2000	0225		0404		3	
DS III	18-1800	0224		0233	GG	3	
DS V	25-180	0224		0226		3	
2.7	GHz	0222.0	0302.0	0326.0		4.81	
610	MHz	0223.0	0306.0	0326.0		5.08	
410	MHz	0224.0	0306.0	0326.0		4.30	
DS III	160-1800	0301		0307	GG	2	
DS III	25-120	0303		0307	G	1	
15.4	GHz	0356.0	0357.0	0402.0		2.26	
8.8	GHz	0356.0	0357.0	0359.0		2.28	
5	GHz	0355.0	0401.0	0404.0		2.74	
2.7	GHz	0355.0	0357.0	0404.0		3.18	
1.4	GHz	0355.0	0402.0	0404.0	P1.4	5.04	
610	MHz	0355.0	0359.0	0404.0		3.53	
410	MHz	0355.0	0403.0	0404.0		3.00	
245	MHz	0403.0	0403.0	~0403.0		2.66	
CME	WL	0254	1774 km/s	-61.4 km/s ²	360°	193°	


Maximum 13d 0223 - 0233 UT

U5 / 15



Maximum 13d 0357 - 0403 UT

P1.4



Particle event: To($E_p > 10$ MeV) – 14d23^h

Tmax($E_p > 10$ MeV) – 15d00^h, Jmax ($E_p > 10$ MeV) – 160 /cm².s.sr

Duration of the event – 3 days

Quasimaximal energy of protons in the event – $E_{qm} \geq 500$ MeV


Sun source: ● solar flare 14d21^h07^m, X1.5/2B, S06W46, AR10930

Main X-ray burst 1-8 Å: onset – 14d21^h07^m, max – 14d22^h15^m, $\Phi = 0.12$ J/m²



CME: 14d22^h30^m, $V = 1042$ km/s, $\Delta\phi = 360^\circ$, dA = 248°;

▲ SC 16d17^h55^m;

Particle fluxes and associated phenomena




Time profiles of the proton fluxes for the event of 2006 December 14



Arrows on the profiles of the events indicate the time of the proton flux maxima, taken to build the proton spectra (see below)

Integral time-of-maximum proton spectrum



Integral fluxes of protons for the event of 2006 December 14

S/c, instruments	Ep, MeV	To	Tmax	Jmax, (cm².s.sr)⁻¹	Dura- tion	Comments
GOES-10						
EPS	>5	23 ^h	15d00 ^h	245	3d	
EPS	>10	23 ^h	15d00 ^h	160	3d	
EPS	>30	23 ^h	15d00 ^h	31	2d	
EPS	>50	22 ^h	15d00 ^h	10	2d	
EPS	>60	22 ^h	23 ^h	6.2	1d	
EPS	>100	22 ^h	23 ^h	1.9	1d	
POES-16						
MEPED	>0.24	-	15d02 ^h	920	3d	
MEPED	>0.8	-	15d02 ^h	410	3d	
MEPED	>2.5	-	15d01 ^h	250	3d	
MEPED	>6.9	-	15d01 ^h	230	3d	
MEPED	>16	-	15d01 ^h	65	2d	
MEPED	>36	-	15d01 ^h	31	2d	
MEPED	>70	-	15d00 ^h	5	1d	
MEPED	>140	-	15d00 ^h	1	1d	
ACE						
SIS	>10	23 ^h	15d01 ^h	49	2d	
SIS	>30	22 ^h	15d01 ^h	23	1d	
SOHO						
EPHIN (INT)	>50	22 ^h	15d01 ^h	1.5	1d	
BALLOONS						
Mi	>68	-	16d11 ^h 39 ^m	0.15	-	After maximum
Mi	>75	-	16d11 ^h 39 ^m	0.11	-	- " -
Mi	>88	-	16d11 ^h 39 ^m	0.07	-	- " -

Differential fluxes of protons for the event of 2006 December 14

S/c, instruments	ΔE , MeV	To	Tmax	J_{\max} , (cm ² .s.sr.MeV) ⁻¹	Dura- tion	Comments
SOHO						
LION	0.75-2	15d01 ^h	15d02 ^h	13.2	3d	
LION	2-6	15d01 ^h	15d02 ^h	1.9	3d	
EPHIN	4-8	15d00 ^h	15d01 ^h	7.8	3d	
EPHIN	8-25	15d00 ^h	15d01 ^h	0.43	3d	
EPHIN	25-41	23 ^h	15d01 ^h	0.17	3d	
STEREO - A						
HET	13.6-29.5	23 ^h	15d00 ^h	1.6	8d	
HET	29.5-60	23 ^h	15d00 ^h	0.3	8d	
HET	60-100	23 ^h	15d00 ^h	0.016	8d	
STEREO - B						
HET	13.6-29.5	23 ^h	15d00 ^h	3.1	8d	
HET	29.5-60	23 ^h	15d00 ^h	0.54	8d	
HET	60-100	23 ^h	15d00 ^h	0.023	8d	
PAMELA						
Tracker	126-136	23 ^h	23 ^h 30 ^m	0.03	<1d	
Tracker	237-265	23 ^h	23 ^h 30 ^m	0.0017	<1d	
Tracker	455-512	23 ^h	23 ^h 30 ^m	0.00013	<1d	


References:

Adriani O., G.C. Barbarino, G.A. Bazilevskaya et al., 2011.
Mulligan T., J.B. Blake, R.A. Mewaldt, 2008.
von Rosenvinge T.T., I.G. Richardson, D.V. Reames et al., 2009.
De Simone N., O. Adriani, G.C. Barbarino et al., 2009.
Veselovskij I.S., I.N. Myagkova, O.S. Yakovchuk, 2012.

Electromagnetic and other phenomena that are sources and/or accompanying for the event of 2006 December 14

2006	December 14	•	AR10930	To event 476			
H α	6563 Å	2206	2211	0011	S06W46	2B	EFZ
1 – 12	keV	2107	2215	2226		X1.5	1.2E-1
6-12	keV	213812	214106	214316		1525	RHESSI
6-12	keV	214316	214734	214824		1713	RHESSI
12-25	keV	233000	233302	233528		174608	RHESSI
15.4	GHz	2207.0	2209.0	0000.0		3.23	
8.8	GHz	2207.0	2211.0	2314.0	[U2.7] P8.8	3.40	
5	GHz	2206.0	2211.0	2316.0		3.18	
2.7	GHz	2207.0	2208.0	0000.0		2.79	
1.4	GHz	2207.0	2209.0	0000.0		4.04	
410	MHz	2207.0	2208.0	0000.0		2.68	
245	MHz	2209.0	2210.0	2227.0		3.08	

DS II	25-290	2209		2219		3	
DS IV	25-2000	2207		2312		2	
DS III	18-90	2219		2226	GG	3	
DS V	25-114	2218		2226		2	
2.7	GHz	2207.0	2228.0	2316.0		3.41	
1.4	GHz	2207.0	2307.0	2315.0	0.4 / 1.4	4.80	
610	MHz	2207.0	2302.0	2308.0		3.89	
410	MHz	2207.0	2247.0	2309.0		3.30	
DS III	20-1300	2244		2248	GG	1	
DS III	300-820	2301		2304	G	2	
CME	WL	2230	1714km/s	-0.4	360°	248°	



Summary Table of Solar Proton Events in the 23rd Cycle of Solar Activity (1997-2006)

Legend:

Event name-(yyyymmdd-doy-year month day-day of the year);

To – the hour of the event beginning (UT), the event date coincides with the date in the event name;

Tmax – time of the first (second and third if any) peak intensity, UT;

Jmax – the flux of protons with energy ≥ 10 MeV in the first (second and third if any) maximum;

γ – power-law index of the integral spectrum ($J(>E) \sim E^{-\gamma}$);

Eqm – quasimaximal energy of proton in this event;

b/s – back side flare event;

W_L-1d – 1 day behind the W-limb;

DSF– the length of the rejected filament in the degrees;

Legend for solar flares is commonly accepted.

The following table shows all 142 events included in the Catalog of solar proton events in the 23rd cycle of solar activity

Particles enhancements						Solar flares or CME – the particle sources			
Event name	T _o	T _{max}	J _{max} (pfu)	γ	Eqm, MeV	T _{Fl max} /T _{0 CME} / T _{0 SC} , (UT) *	Class of flare	Localizat ion	Active region
19971104-308	07 ^h	04d11 ^h 05d02 ^h 20	66 17.5	2.0 1.8	470 320	● 04d05 ^h 58 ^m	X2.1/3B	S14W34	8100
19971106-310	13 ^h	07d02 ^h	430	3.5	2900	● 06d11 ^h 55 ^m	2B/X9.4	S18W63	8100
19971113-317	23 ^h	14d04 ^h	1.3	1.7	170	☐ <13d22h26m	CME	b/s, W _L	8100, W _L -5d
19980420-110	11 ^h	21d06 ^h 21d12 ^h	860 1.6·10 ³	3.3 2.4	440 600	■ 20d10 ^h 21 ^m	M1.4/EPL	s20w90	8194, W _L -2d
19980430-120	02 ^h	30d15 ^h 01d15 ^h	1.3 1.2	2.9 2.6	65 75	● 29d16 ^h 37 ^m	M6.8/3B	S16E22	8210
19980502-122	14 ^h	02d16 ^h	130	1.8	800	● 02d13 ^h 42 ^m	X1.1/3B	S15W15	8210
19980506-126	08 ^h	06d09 ^h	120	2.6	575	● 06d08 ^h 09 ^m	X2.7/1N	S15W64	8210
19980509-129	06 ^h	09d13 ^h 09d23 ^h	4.7 8.9	2.3 2.3	230 230	■ 09d03 ^h 40 ^m	M7.7/ -	s15w90	8210, W _L -1d
19980616-167	21 ^h	17d09 ^h 18d02 ^h	1.3 1.4	2.4 2.2	80 75	■ 16d18 ^h 42 ^m	M1.0/ -	s22w90	8232, W _L -1d
19980822-234	06 ^h	23d00 ^h 23d08 ^h	1.7 1.5	1.8 1.8	80 85	● 22d00 ^h 09 ^m	M9.0/2B	N42E51	8307
19980824-236	23 ^h	25d02 ^h 26d07 ^h	96 320	1.7 3.4	720 310	● 24d22 ^h 12 ^m	3B/X1.0	N35E09	8307
19980923-266	13 ^h	25d01 ^h	22	3.3	75	● 23d07 ^h 13 ^m	M7.1/3B	N19E09	8340
19980930-273	14 ^h	30d23 ^h	785	2.3	600	● 30d13 ^h 50 ^m	M2.8/2N	N23W78	8340
19981018-291	22 ^h	19d02 ^h 19d06 ^h	1.8 2.3	2.2 2.8	140 80	o 15d<10 ^h 05 ^m o 18d01 ^h 45 ^m	DSF/27° M2.4/2B	N19E10 N16W53	8358
19981106-310	03 ^h	06d12 ^h	4.6	3.0	75	● 05d19 ^h 55 ^m	M8.4/2B	N22W18	8375
19981107-311	12 ^h	07d14 ^h 08d02 ^h	2.8 6	2.4 2.1	75 80	● 07d11 ^h 06 ^m ▲ SC 08d04 ^h 51 ^m	M2.4/SN	N14W43	8375
19981114-318	06 ^h	14d12 ^h 15d06 ^h	250 10	2.4 2.9	580 190	☐ 14d05 ^h 08 ^m	C1.3/BSL	n28w90	8375, W _L -2d
19981122-326	07 ^h	22d09 ^h 22d14 ^h	1.1 0.6	1.5 1.6	285 160	● 22d06 ^h 42 ^m	X3.7/1N	S27W82	8384

Particles enhancements						Solar flares or CME – the particle sources			
Event name	T ₀	T _{max}	J _{max} (pfu)	γ	Eqm, MeV	T _{Fl max} /T _{0 CME} */ T _{0 SC} (UT)	Class of flare	Localizat ion	Active region
19981124-328	03 ^h	24d10 ^h	1.25	1.6	210	● 24d02 ^h 20 ^m	X1.0/SF	S30W81	8384
19990120-020	23 ^h	21d11 ^h 22d06 ^h	1.3 1.0	1.7 1.8	270 250	▣ 20d20 ^h 04 ^m	M5.2/...	n27e90	unknown, E _L
19990122-022	02 ^h	22d16 ^h 22d14 ^h	3 5	1.7 3.1	85 85	unknown Ø 22d17 ^h 24 ^m	M1.4/SF	N19W44	unknown 8440
19990424-114	15 ^h	24d21 ^h 25d06 ^h	3.7 4.3	2.2 2.3	85 210	▣ 24d<13 ^h 31 ^m	CME	b/s, W _L	8517, W _L -3d
19990504-124	08 ^h	05d21 ^h 06d06 ^h	3.7 4.0	2.8 3.2	75 65	⊙ 03d06 ^h 02 ^m	M4.4/2N	N15E32	8525
19990509-129	19 ^h	09d21 ^h	1.2	1.5	75	■ 09d18 ^h 07 ^m	M7.6/ ...	n23w90	8526, W _L -1d
19990527-147	12 ^h	27d13 ^h	2.75	1.6	275	▣ < 27d11 ^h 06 ^m ○ 26d19 ^h 32 ^m	CME M1.2/2N	b/s, W _L N17E46	unknown 8552
19990601-152	20 ^h	02d09 ^h 02d21 ^h	23 13	1.9 2.3	350 240	▣ 01d19 ^h 04 ^m	C1.2/...	n25w90	unknown
19990604-155	08 ^h	04d12 ^h	20	2.3	300	● 04d07 ^h 03 ^m	M3.9/2N	N18W72	8552
19990611-162	01 ^h	11d03 ^h	2.2	1.7	240	▣ 11d01 ^h 10 ^m	C1.0/...	b/s W event	unknown
19990625-176	10 ^h	26d12 ^h	1.7	3.5	40	Ø 26d05 ^h 12 ^m	M2.3/2B	N24E02	8598
19991117-321	19 ^h	19d02 ^h 19d23 ^h	1 0.4	2.55 3.1	50 25	● 17d09 ^h 57 ^m	M7.4/2B	N17E21	8766
20000218-049	06 ^h	18d12 ^h	1.7	2.0	290	⊙ 17d20 ^h 45 ^m	M1.3/2N	S29E07	8872
20000404-095	17 ^h	05d02 ^h 06d06 ^h	25 4	3.1 2.9	105 75	● 04d15 ^h 41 ^m	C9.7/2F	N16W66	8933
20000607-159	00 ^h	08d10 ^h	54	3.3	100	● 06d15 ^h 25 ^m	X2.3/3B	N20E18	9026
20000610-162	17 ^h	10d20 ^h	24	2.2	390	● 10d17 ^h 02 ^m	M5.2/3B	N22W39	9026
20000617-169	07 ^h	18d06 ^h	1.7	2.1	110	● 17d02 ^h 37 ^m	M3.5/2B	N22W72	9033
20000625-177	10 ^h	26d07 ^h	1.5	2.6	70	⊙ 25d07 ^h 52 ^m	M1.9/2N	N16W55	9046
20000713-195	06 ^h	13d10 ^h	5	3.1	40	⊙ 12d18 ^h 47 ^m	M5.7/2F	N16W64	9070
20000714-196	10 ^h	14d18 ^h 15d13 ^h	7.2 10 ³ 1.8 10 ⁴	3.7 4.7	2160 630	● 14d10 ^h 24 ^m SC 15d14 ^h 37 ^m	X5.7/3B	N22W07	9077
20000716-198	11 ^h	16d12 ^h 17d02 ^h	100 37	2.5 2.3	370 320	⊙ 15d08 ^h 33 ^m Ø 16d23 ^h 37 ^m	M1.3/SF M1.4/2F	N16W12 N17W40	9077
20000722-204	12 ^h	22d14 ^h 22d20 ^h	13 6	1.8 2.6	340 80	● 22d11 ^h 34 ^m	M3.7/2N	N14W56	9085
20000728-210	02 ^h	28d06 ^h 28d12 ^h	5 13	2.1 2.8	140 105	⊙ 27d23 ^h 42 ^m	M1.2/SF	N11W78	9090
20000811-224	15 ^h	11d17 ^h	3.2	3.3	75	▣ <11d16 ^h 54 ^m	CME	b/s, E _L	unknown
20000813-226	01 ^h	13d06 ^h	1.2	3.35	70	● 12d09 ^h 56 ^m	M1.1/SN	S16W79	9119
20000912-256	14 ^h	13d02 ^h	180	2.6	350	● 12d12 ^h 00 ^m	2F/M1.0	S19W08	9163
20001016-290	08 ^h	16d11 ^h 16d17 ^h	3.5 9.8	1.7 2.6	310 140	▣ 16d07 ^h 28 ^m	M2.5/-	n05w90	9182
20001025-299	13 ^h	25d23 ^h	4.1	2.2	95	▣ 25d11 ^h 25 ^m	C4/...	... W90	unknown
20001031-305	07 ^h	01d03 ^h	2.1	2.3	70	○ 31d03 ^h 00 ^m	C6.0/1N	S20E80	9209
20001108-313	23 ^h	09d15 ^h	9.7 10 ³	3.7	650	● 08d23 ^h 28 ^m	3F/M7.4	N20W66	9213
20001124-329	07 ^h	24d21 ^h	65	1.7	460	● 24d05 ^h 02 ^m	X2.0/3B	N20W05	9236
20001126-331	03 ^h	26d20 ^h	670	2.9	400	● 25d01 ^h 31 ^m	M8.2/2N	N07E50	9240
20010128-028	18 ^h	29d01 ^h	29	2.1	325	● 28d16 ^h 40 ^m	1N/M1.5	S04W59	9313
20010226-057	09 ^h	26d20 ^h	1	2.4	65	▣ 26d07 ^h 41 ^m	C1.6/...	s04w90	9354
20010326-085	20 ^h	27d08 ^h	1.8	3.0	55	⊙ 26d13 ^h 26 ^m	M2.2/1F	N15E27	9393

Particles enhancements						Solar flares or CME – the particle sources			
Event name	T _o	T _{max}	J _{max} (pfu)	γ	Eqm, MeV	T _{Fl max} /T _{0 CME} / T _{0 SC} (UT) *	Class of flare	Localizat ion	Active region
20010329-088	13 ^h	29d19 ^h 31d00 ^h	7 22	2.3 3.1	215 115	● 29d10 ^h 15 ^m	1N/X1.7	N16W12	9393
20010402-092	23 ^h	03d07 ^h	112	2.9	575	■ 02d21 ^h 51 ^m	X>17.5/	n19w90	9393
20010409-099	17 ^h	09d20 ^h	2.2	1.2	390	● 09d15 ^h 34 ^m	M7.9/1B	S21W04	9415
20010410-100	08 ^h	11d01 ^h 11d20 ^h	50 280	2.2 2.8	350 260	● 10d05 ^h 26 ^m Ø 11d13 ^h 26 ^m	3N/X2.3 M2.3/1F	S23W09 S20W28	9415
20010412-102	12 ^h	12d17 ^h 13d10 ^h	4.3 8.7	1.45 1.5	410 275	● 12d10 ^h 28 ^m	X2.0/2B	S20W42	9415
20010415-105	14 ^h	15d16 ^h	270	4.2	3480	● 15d13 ^h 50 ^m	X14.4/2B	S20W84	9415
20010418-108	03 ^h	18d10 ^h	190	2.8	2100	▣ 18d02 ^h 14 ^m	C2.2	s20w90	9415
20010427-117	03 ^h	28d05 ^h	15	3.6	80	● 26d13 ^h 20 ^m	M7.8/2B	N17W31	9433
20010507-127	14 ^h	07d18 ^h 08d12 ^h	7.7 11.5	2.3 3.15	80 85	▣ <07d12 ^h 06 ^m	CME	b/s, W _L	9433, 5d-W _L
20010520-140	07 ^h	20d10 ^h	1.8	1.75	410	● 20d06 ^h 03 ^m	M6.4/	s18w90	9455
20010615-166	16 ^h	15d20 ^h 16d06 ^h	5 8.1	2.1 2.7	335 120	▣ <15d15 ^h 56 ^m	CME	b/s, W _L	unknown
20010809-221	20 ^h	09d11 ^h	6	3.3	85	☉ 09d11 ^h 22 ^m	C3.7/SF	N10E54	DSF
20010816-228	00 ^h	16d03 ^h 17d12 ^h	87 75	3.0 2.1	600 475	▣ <15d23 ^h 54 ^m	CME	b/s, W _L -	9557?, 5d-W _L
20010915-258	12 ^h	15d15 ^h	6	2.4	150	● 15d11 ^h 28 ^m	1N/M1.5	S21W49	9608
20010924-267	11 ^h	24d18 ^h 25d07 ^h 25d22 ^h	390 1.1·10 ³ 9.5·10 ³	3.8 2.5 3.2	470 580 580	● 24d10 ^h 38 ^m Ø 25d04 ^h 24 ^m SC 25d20 ^h 25 ^m	X2.6/2B M7.6/1N	S17E26 S18W01	9632 9628
20011001-274	14 ^h	01d23 ^h 02d07 ^h	370 1.3·10 ³	4.8 5.6	155 150	■ 01d05 ^h 15 ^m	M9.1/...	s18w80	9628
20011019-292	02 ^h	19d08 ^h 19d21 ^h	3.6 8	1.75 1.9	300 310	● 19d01 ^h 05 ^m Ø 19d16 ^h 13 ^m	X1.6/2B X1.6/2B	N16W18 N15W30	9691
20011022-295	16 ^h	22d21 ^h	17	1.5	425	● 22d17 ^h 59 ^m	X1.2/2B	S18E16	9672
20011028-301	02 ^h	28d07 ^h	1.1	2.6	60	○ 28d04 ^h 50 ^m	M1.3/1F	N12E40	9682
20011104-308	16 ^h	04d20 ^h 06d00 ^h	540 2.4·10 ⁴	3.2 4.5	750 685	Ø 04d06 ^h 43 ^m ● 04d16 ^h 20 ^m	C8.4/1N X1.0/3B	N14W57 N07W19	9682 9684
20011117-321	10 ^h	19d22 ^h	13	4.6	45	● 17d05 ^h 25 ^m	M2.8/2N	S13E42	9704
20011122-326	21 ^h	23d10 ^h 24d06 ^h	2.7·10 ³ 1.1·10 ⁴	4.2 4.5	390 350	Ø 22d20 ^h 36 ^m ● 22d<23 ^h 30 ^m	M3.8/2B M9.9/3B	S26W68 S13W38	9698 9704
20011226-360	05 ^h	26d11 ^h	336	2.55	800	● 26d05 ^h 40 ^m	M7.1/1B	N08W54	9742
20011229-363	05 ^h	29d08 ^h	40	2.7	195	■ 28d20 ^h 45 ^m	X3.4/-	s26e90	9767
20011230-364	20 ^h	31d02 ^h 31d16 ^h	25.5 75	2.6 3.1	190 170	▣ 29d21 ^h 27 ^m	M1.8/	s08w90	9748
20020110-010	02 ^h	11d01 ^h	70	3.0	85	☉ 09d18 ^h 01 ^m	M9.5/2B	N13W02	9773
20020115-015	07 ^h	15d18 ^h	7.5	2.15	80	▣ 14d06 ^h 27 ^m	M4.4/...	s23w90	9767, 1.5d-W _L
20020220-051	07 ^h	20d08 ^h	3.3	2.5	145	● 20d06 ^h 12 ^m	M5.1/1N	N12W72	9825
20020316-075	02 ^h	16d13 ^h 17d11 ^h	1.3 1.1	2.7 2.4	70 65	● 15d23h10m	M2.2/1F	S08W03	9866
20020318-077	00 ^h	18d15 ^h 19d06 ^h	14.5 20	2.65 2.6	155 145	☉ 17d19 ^h 31 ^m Ø 18d02 ^h 31 ^m	M4.0/SF M1.0/SN	S22E16 S16E27	9871
20020320-079	13 ^h	20d17 ^h	8	3.6	60	○ 20d08 ^h 33 ^m ▲ SC20d13 ^h 28 ^m	C1.9/SF	S19W41	9873

Particles enhancements						Solar flares or CME – the particle sources			
Event name	T ₀	T _{max}	J _{max} (pfu)	γ	Eqm, MeV	T _{Fl max} /T _{0 CME} / T _{0 SC} (UT) *	Class of flare	Localizat ion	Active region
20020322-081	12 ^h	22d20 ^h 23d13 ^h	1 9	3.7 3.85	45 50	■ 22d11 ^h 14 ^m SC 23d11 ^h 37 ^m	M1.6/...	s10w90	9866
20020417-107	10 ^h	17d16 ^h	21	2.75	120	● 17d08h24m	M2.6/2N	S14W36	9906
20020419-109	05 ^h	19d09 ^h 19d19 ^h	1 2.7	3.5 3.4	45 50	SC 19d08 ^h 36 ^m ○ 19d15h21m	- C2.5/SF	- S16W59	9906
20020421-111	01 ^h	21d03 ^h 21d09 ^h	915 1.7·10 ³	1.9 2.8	575 570	■ 21d01 ^h 51 ^m	X1.5/1F	S14W84	9906
20020522-142	07 ^h	23d10 ^h 23d16 ^h	260 87	3.75 3	125 175	⊙ 22d03 ^h 54 ^m	C5.0/SF	S22W53	DSF
20020707-188	13 ^h	07d20 ^h	26	2.8	85	■ 07d11 ^h 43 ^m	M1/...	s19w90	10017
20020716-197	12 ^h	16d22 ^h 17d14 ^h	27 85	3.5 3.4	125 135	● 15d20 ^h 08 ^m Ø 17d07 ^h 13 ^m	3B/X3.0 M8.5/1B	N19W01 N20W16	10030
20020719-200	05 ^h	19d11 ^h	3.6	2.15	215	□ 18 ^d 07 ^h 44 ^m	X1.8/2B	N19W33	10030
20020722-203	01 ^h	22d11 ^h	18.5	2.6	85	□ 20d21 ^h 30 ^m	X3.3/...	s13e90	10039
20020814-226	06 ^h	14d09 ^h 14d16 ^h	6.7 6.9	2.5 4	85 50	● 14d02 ^h 12 ^m	M2.3/1N	N10W54	10061
20020817-229	00 ^h	17d10 ^h	1.7	2.6	70	● 16d12 ^h 32 ^m	2N/M5.2	S14E20	10069
20020818-230	22 ^h	19d03 ^h 19d12 ^h	2.3 1.8	2.1 2.25	75 70	● 18d21 ^h 25 ^m	1N/M2.2	S10W20	10069
20020820-232	09 ^h	20d10 ^h	2.5	1.95	80	● 20d01 ^h 40 ^m	M5.0/1B	S10W35	10069
20020822-234	03 ^h	22d05 ^h	16	1.5	450	● 22d01 ^h 57 ^m	M5.4/2B	S07W62	10069
20020824-236	01 ^h	24d03 ^h	92	2.1	775	● 24d01 ^h 12 ^m	X3.1/1F	S02W81	10069
20020906-249	06 ^h	06d14 ^h 07d17 ^h	3 67	2.2 3.15	80 175	⊙ 05d17 ^h 06 ^m SC 07d16 ^h 38 ^m	C5.2/SF	N12E28	10102
20021109-313	17 ^h	10d02 ^h 10d13 ^h	150 40	3.5 3.1	160 145	● 09d13 ^h 23 ^m Ø 10d03 ^h 21 ^m	M4.6/2B M2.4/2N	S04W29 S12W37	10180
20030528-148	04 ^h	28d11 ^h 29d16 ^h	2 77	2.0 2.95	170 175	● 27d23 ^h 07 ^m Ø 29d01 ^h 05 ^m	X1.3/2B X1.3/2B	S06W20 S06W37	10365
20030531-151	03 ^h	31d06 ^h	15.6	1.5	415	● 31d02 ^h 24 ^m	M9.3/2B	S07W65	10365
20030618-169	08 ^h	20d06 ^h	10.2	2.7	180	⊙ 17d22 ^h 55 ^m	M6.8/...	s12e60	10368
20031026-299*	18 ^h	26d20 ^h 27d02 ^h	230 360	3.1 2.9	340 400	● 26d18 ^h 19 ^m	X1.2/1N	N02W38	10484
20031028-301	12 ^h	28d18 ^h 29d02 ^h	4.6·10 ³ 1.2·10 ⁴	2.9 3.9	3340 1025	● 28d11 ^h 10 ^m	X17.2/4B	S16E08	10486
20031029-302	22 ^h	29d23 ^h	2.2·10 ³	2.0	810	● 29d20 ^h 49 ^m	X10.0/2B	S15W02	10486
20031102-306	17 ^h	02d23 ^h	990	3.1	1700	● 02d17 ^h 25 ^m	X8.3/2B	S14W56	10486
20031104-308	22 ^h	05d07 ^h	126	2.7	445	● 04d19 ^h 50 ^m	X>17.5/3B	S19W83	10486
20031120-324	08 ^h	20d11 ^h	4.4	2.2	140	● 20d07 ^h 47 ^m SC 20d08 ^h 03 ^m	M9.6/2B	N01W08	10501
20031121-325	08 ^h	22d02 ^h	10.7	2.5	80	● 20d23 ^h 53 ^m	M5.8/2B	N02W17	10501
20031202-336	12 ^h	02d18 ^h	21	4.0	100	⊙ 02d09 ^h 47 ^m	C7.2/...	s19w90	10508
20040411-102	06 ^h	11d12 ^h 11d20 ^h	13 14.5	3.0 3.0	95 80	● 11d04 ^h 19 ^m Ø <11d11 ^h 54 ^m	C9.6/1F CME	S14W47 b/s, W _L	10588 unknown
20040722-204	17 ^h	22d20 ^h 23d10 ^h	0.9 2	2.0 2.0	70 155	⊙ 22d00 ^h 32 ^m	M9.1/SB	n06e25	10652

Particles enhancements						Solar flares or CME – the particle sources			
Event name	T ₀	T _{max}	J _{max} (pfu)	γ	Eqm, MeV	T _{Fl max} /T _{0 CME} / T _{0 SC} (UT) *	Class of flare	Localizat ion	Active region
20040723-205	16 ^h	23d19 ^h	1.8	2.25	75	☉ 22d22 ^h 58 ^m	2N/M1.6	N05E04	10652
20040725-207	17 ^h	25d21 ^h 26d23 ^h	27 430	2.8 3.8	140 155	● 25d15 ^h 14 SC26d22 ^h 49 ^m	M1.1/1F	N08W33	10652
20040801-214	01 ^h	01d21 ^h 02d02 ^h	5.2 4.8	2.25 2.3	80 75	☐ 31d05 ^h 16 ^m Ø 31d10 ^h 35 ^m	C8.4/... C5.3/	n02w90 n02w90	10652 >1.5d W _L
20040913-257	19 ^h	13d23 ^h 14d05 ^h	210 180	3.4 4.1	110 90	☉ 12d00 ^h 56 ^m Ø 12d01 ^h 39	M4.8/2N M3.2/SN	N04E42 S14W61	10672 10667
2004.09.19-263	18 ^h	20d01 ^h 21d02 ^h	46 10	2.2 2.2	390 100	● 19d17 ^h 12 ^m	M1.9/...	n03w60	10672
20041101-306	06 ^h	01d08 ^h	54	1.9	410	● 01d03 ^h 22 ^m	M1.1/1F	N15W41	10691
20041107-312	01 ^h	07d23 ^h 09d00 ^h	490 70	3.0 3.2	330 100	● 07d15 ^h 42 ^m Ø 08d15 ^h 49 ^m	X2.0/... M2.3/1N	n10w15 N08W35	10696
20041110-315	02 ^h	10d10 ^h 10d16 ^h 12d09 ^h	264 193 75	2.2 2.5 3.0	485 430 110	● 09d17 ^h 19 ^m Ø 10d02 ^h 13 ^m	M8.9/2N X2.5/3B	N07W51 N09W49	10969
20050115-015	07 ^h	15d11 ^h	7.4	1.7	300	● 15d06 ^h 38 ^m	M8.6/SF	N11E06	10720
20050116-016	00 ^h	16d18 ^h	330	2.9	330	● 15d23h02m	X2.6/3B	N14W08	10720
20050117-017	13 ^h	17d17 ^h	3.8·10 ³	2.8	750	● 17d09 ^h 52 ^m	X3.8/3N	N14W24	10720
20050120-020	06 ^h	20d10 ^h 21d17 ^h	1.1·10 ³ 134	3.9 2.2	3840 1520	● 20d07 ^h 01 ^m SC 21d17 ^h 11 ^m	X7.1/2B	N12W58	10720
20050513-133	19 ^h	14d03 ^h 14d14 ^h 15d03 ^h	7.7 155 1.9·10 ³	1.85 3.3 4.4	300 85 85	● 13d16 ^h 57 ^m SC 15d 02 ^h 38 ^m	M8/2B	N12E12	10759
20050616-167	20 ^h	17d04 ^h	41	1.8	510	■ 16d20 ^h 22 ^m	M4.0/SF	N09W87	10775
20050710-191	03 ^h	10d05 ^h 10d12 ^h	1.1 1.9	1.85 2.8	75 70	● 09d22 ^h 06 ^m	M2.8/1N	N11W27	10786
20050713-194	17 ^h	15d04 ^h	9.7	2.6	115	● 13d14 ^h 49 ^m	M5/SF	N10W80	10786
20050714-195	14 ^h	15d03 ^h	130	3.15	185	■ 14d10 ^h 55 ^m	X1.2/...	n11w90	10786
20050717-198	14 ^h	17d18 ^h 17d22 ^h	12 19	2.25 3.9	85 85	☐ 17d10 ^h 32 ^m	B1.1/...	n13w9	10786, 3d-W _L
20050725-206	21 ^h	28d14 ^h 29d14 ^h	30 36	2.8 2.8	85 85	☐ 27d04 ^h 33 ^m Ø28d22 ^h 08 ^m	M3.7/... M4.8/SF	n10e90 N08E84	10792, E _L 10792
20050731-212	22 ^h	01d05 ^h 02d01 ^h	21 6	4.1 3.3	65 65	☉ 30d 06 ^h 35 ^m Ø 01d13 ^h 51 ^m	X1.3/2B M1.0/1F	N12E61 N14E29	10792
20050822-234	03 ^h	22d07 ^h	5.4	2.4	80	● 22d01 ^h 33 ^m	M2.6/1F	S09W48	10798
20050822a-234	19 ^h	23d02 ^h 23d10 ^h	280 290	3.1 3.0	330 290	● 22d16 ^h 52 ^m	M5.6/1N	S12W60	10798
20050907-250	21 ^h	08d20 ^h 10d11 ^h	70 1000	2.3 3.0	800 555	● 07d17 ^h 40 ^m Ø09d20 ^h 04 ^m	X17.0/3B X6.2/2B	S06E89 S12E67	10808
20050914-257	00 ^h	14d15 ^h 15d08 ^h	160 180	3.5 3.75	85 85	● 13d19 ^h 27 ^m SC 08 ^h 35 ^m	X1.5/2B	S09E10	10808
20061205-339	15 ^h	05d20 ^h	2.5	1.8	275	● 05d10 ^h 35 ^m	X9/2N	S07E79	10930
20061206-340	10 ^h	7d22 ^h	1.9·10 ³	2.6	850	● 06d18 ^h 47 ^m	X6.5/3B	S06E63	10930
20061213-347	03 ^h	13d09 ^h	660	4.7	3440	● 13d02 ^h 40 ^m	X3.4/4B	S06W24	10930
20061214-348	23 ^h	15d00 ^h	160	2.5	≥500	● 14d22 ^h 15 ^m	X1.5/2B	S06W46	10930

*) T_{Fl max} – time of peak intensity for a flare (UT);
T_{0 CME} – time of liftoff of CME in sight of coronagraph;
T_{0 SC} – time of SC arrival to the Earth.